

Hsuan-Ying Huang

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2977564/publications.pdf>

Version: 2024-02-01

141
papers

5,133
citations

87888

38
h-index

110387

64
g-index

143
all docs

143
docs citations

143
times ranked

6751
citing authors

#	ARTICLE	IF	CITATIONS
1	Pulmonary "Inflammatory Leiomyosarcomas" Are Indolent Tumors With Diploid Genomes and No Convincing Rhabdomyoblastic Differentiation. <i>American Journal of Surgical Pathology</i> , 2022, 46, 424-433.	3.7	3
2	The emerging PRRX1-NCOA fibroblastic neoplasm: a combined reappraisal of published tumors and two new cases. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2022, 481, 111-116.	2.8	5
3	Molecular Characterization of Dermatofibrosarcoma Protuberans. <i>American Journal of Surgical Pathology</i> , 2022, 46, 942-955.	3.7	8
4	Adult NTRK-rearranged spindle cell neoplasms of the viscera: with an emphasis on rare locations and heterologous elements. <i>Modern Pathology</i> , 2022, 35, 911-921.	5.5	13
5	Characterization of Aberrations in DNA Damage Repair Pathways in Gastrointestinal Stromal Tumors: The Clinicopathologic Relevance of γ -H2AX and 53BP1 in Correlation with Heterozygous Deletions of CHEK2, BRCA2, and RB1. <i>Cancers</i> , 2022, 14, 1787.	3.7	1
6	Superficial ALK-rearranged myxoid spindle cell neoplasm with a novel FMR1-ALK fusion gene. <i>Modern Pathology</i> , 2022, 35, 438-441.	5.5	3
7	Fine-needle aspiration cytology of melanotic schwannoma in the submandibular gland. <i>Diagnostic Cytopathology</i> , 2021, 49, 142-145.	1.0	1
8	Clinicopathological and molecular characterisation of <i>USP6</i> -rearranged soft tissue neoplasms: the evidence of genetic relatedness indicates an expanding family with variable bone-forming capacity. <i>Histopathology</i> , 2021, 78, 676-689.	2.9	17
9	RSF1 requires CEBP β and hSNF2H to promote IL-1 β -mediated angiogenesis: the clinical and therapeutic relevance of RSF1 overexpression and amplification in myxofibrosarcomas. <i>Angiogenesis</i> , 2021, 24, 533-548.	7.2	4
10	Toward a unifying entity that encompasses most, but perhaps not all, inflammatory leiomyosarcomas and histiocyte-rich rhabdomyoblastic tumors. <i>Modern Pathology</i> , 2021, 34, 1434-1438.	5.5	8
11	Infant malignant ectomesenchymoma masquerading as inguinal hernia in two patients. <i>Pediatrics and Neonatology</i> , 2021, 62, 324-326.	0.9	1
12	S100P as a marker for poor survival and advanced stage in gallbladder carcinoma. <i>Annals of Diagnostic Pathology</i> , 2021, 52, 151736.	1.3	1
13	Ossifying low grade endometrial stromal sarcoma with PHF1-BRD8 fusion. <i>Cancer Genetics</i> , 2021, 256-257, 81-85.	0.4	1
14	Myoepithelial and oral intracranial myxoid mesenchymal tumor-like neoplasms as diagnostic considerations of the ever-expanding extracranial myxocollagenous tumors harboring FET-CREB fusions. <i>Pathology Research and Practice</i> , 2021, 229, 153700.	2.3	4
15	Deep Convolutional Neural Networks Detect Tumor Genotype from Pathological Tissue Images in Gastrointestinal Stromal Tumors. <i>Cancers</i> , 2021, 13, 5787.	3.7	10
16	Sinonasal Angioleiomyoma. <i>Ear, Nose and Throat Journal</i> , 2020, 99, NP109-NP110.	0.8	3
17	GNA11 joins GNAQ and GNA14 as a recurrently mutated gene in anastomosing hemangioma. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 476, 475-481.	2.8	21
18	What is new about the molecular genetics in matrix-producing soft tissue tumors? -The contributions to pathogenetic understanding and diagnostic classification. <i>Virchows Archiv Fur Pathologische Anatomie Und Physiologie Und Fur Klinische Medizin</i> , 2020, 476, 121-134.	2.8	8

#	ARTICLE	IF	CITATIONS
19	Recurrent YAP1 and KMT2A Gene Rearrangements in a Subset of MUC4-negative Sclerosing Epithelioid Fibrosarcoma. <i>American Journal of Surgical Pathology</i> , 2020, 44, 368-377.	3.7	61
20	Dynamic contrast-enhanced magnetic resonance imaging in correlation with tongue cancer stages. <i>Acta Radiologica</i> , 2020, 62, 028418512097518.	1.1	1
21	Soft Tissue Special Issue: Giant Cell-Rich Lesions of the Head and Neck Region. <i>Head and Neck Pathology</i> , 2020, 14, 97-108.	2.6	25
22	First Glance of Molecular Profile of Atypical Cellular Angiofibroma/Cellular Angiofibroma with Sarcomatous Transformation by Next Generation Sequencing. <i>Diagnostics</i> , 2020, 10, 35.	2.6	5
23	Head and Neck Mesenchymal Neoplasms With GLI1 Gene Alterations. <i>American Journal of Surgical Pathology</i> , 2020, 44, 729-737.	3.7	46
24	Primary intradural extramedullary spinal mesenchymal chondrosarcoma: case report and literature review. <i>BMC Musculoskeletal Disorders</i> , 2019, 20, 408.	1.9	9
25	The histologic spectrum of soft tissue spindle cell tumors with <i>NTRK3</i> gene rearrangements. <i>Genes Chromosomes and Cancer</i> , 2019, 58, 739-746.	2.8	86
26	Clinicopathologic Characterization of GREB1-rearranged Uterine Sarcomas With Variable Sex-Cord Differentiation. <i>American Journal of Surgical Pathology</i> , 2019, 43, 928-942.	3.7	43
27	The expanding morphological and genetic spectrum of MYOD1 mutant spindle cell/sclerosing rhabdomyosarcomas: a clinicopathological and molecular comparison of mutated and non-mutated cases. <i>Histopathology</i> , 2019, 74, 933-943.	2.9	18
28	Histiocyte-rich rhabdomyoblastic tumor: rhabdomyosarcoma, rhabdomyoma, or rhabdomyoblastic tumor of uncertain malignant potential? A histologically distinctive rhabdomyoblastic tumor in search of a place in the classification of skeletal muscle neoplasms. <i>Modern Pathology</i> , 2019, 32, 446-457.	5.5	29
29	Solitary fibrous tumor: An evolving and unifying entity with unsettled issues. <i>Histology and Histopathology</i> , 2019, 34, 313-334.	0.7	35
30	USP6 gene rearrangement differentiates primary paranasal sinus solid aneurysmal bone cyst from other giant cell-rich lesions: report of a rare case. <i>Human Pathology</i> , 2018, 76, 117-121.	2.0	13
31	Re: Increased expression of interleukin-23 associated with progression of colorectal cancer. <i>Journal of Surgical Oncology</i> 2017;115(2):208-212. <i>Journal of Surgical Oncology</i> , 2018, 118, 723-723.	1.7	0
32	Multiple epithelioid hemangiomas in an unusual location. <i>Dermatologica Sinica</i> , 2018, 36, 258-259.	0.5	0
33	Hydroxysteroid 11-Beta Dehydrogenase 1 Overexpression with Copy-Number Gain and Missense Mutations in Primary Gastrointestinal Stromal Tumors. <i>Journal of Clinical Medicine</i> , 2018, 7, 408.	2.4	9
34	Overexpressed Fatty Acid Synthase in Gastrointestinal Stromal Tumors: Targeting a Progression-Associated Metabolic Driver Enhances the Antitumor Effect of Imatinib. <i>Clinical Cancer Research</i> , 2017, 23, 4908-4918.	7.0	27
35	Primary paranasal sinus clear cell carcinoma with EWSR1-ATF1 fusion: report of 2 molecularly confirmed cases exhibiting unique histopathology. <i>Human Pathology</i> , 2017, 63, 139-143.	2.0	14
36	Comprehensive screening for MED12 mutations in gynaecological mesenchymal tumours identified morphologically distinctive mixed epithelial and stromal tumours. <i>Histopathology</i> , 2017, 70, 954-965.	2.9	7

#	ARTICLE	IF	CITATIONS
37	<i>TFE3</i> rearranged hepatic epithelioid hemangioendothelioma—a case report with immunohistochemical and molecular study. <i>Apmis</i> , 2017, 125, 849-853.	2.0	12
38	Increased expression of interleukin-23 associated with progression of colorectal cancer. <i>Journal of Surgical Oncology</i> , 2017, 115, 208-212.	1.7	19
39	ALK oncoproteins in atypical inflammatory myofibroblastic tumours: novel RRBP1-ALK fusions in epithelioid inflammatory myofibroblastic sarcoma. <i>Journal of Pathology</i> , 2017, 241, 316-323.	4.5	87
40	Myxoid Plexiform Fibrohistiocytic Tumor Masquerading as Ganglion Cyst: A Case Report and Literature Review. <i>Case Reports in Pathology</i> , 2017, 2017, 1-4.	0.3	1
41	PLCB4 copy gain and PLC γ 4 overexpression in primary gastrointestinal stromal tumors: Integrative characterization of a lipid-catabolizing enzyme associated with worse disease-free survival. <i>Oncotarget</i> , 2017, 8, 19997-20010.	1.8	20
42	Increased expression of SKP2 is an independent predictor of locoregional recurrence in cervical cancer <i>via</i> promoting DNA-damage response after irradiation. <i>Oncotarget</i> , 2016, 7, 44047-44061.	1.8	21
43	Novel PAX3-NCOA1 Fusions in Biphenotypic Sinonasal Sarcoma With Focal Rhabdomyoblastic Differentiation. <i>American Journal of Surgical Pathology</i> , 2016, 40, 51-59.	3.7	102
44	Clinicopathological and genetic heterogeneity of the head and neck solitary fibrous tumours: a comparative histological, immunohistochemical and molecular study of 36 cases. <i>Histopathology</i> , 2016, 68, 492-501.	2.9	49
45	<i>BCOR</i> – <i>CCNB3</i> positive soft tissue sarcoma with round cell and spindle cell histology: a series of four cases highlighting the pitfall of mimicking poorly differentiated synovial sarcoma. <i>Histopathology</i> , 2016, 69, 792-801.	2.9	41
46	<i>NAB2</i> – <i>STAT6</i> gene fusion and STAT6 immunoeexpression in extrathoracic solitary fibrous tumors: the association between fusion variants and locations. <i>Pathology International</i> , 2016, 66, 288-296.	1.3	42
47	The clinicopathological significance of <i>NAB2</i> – <i>STAT6</i> gene fusions in 52 cases of intrathoracic solitary fibrous tumors. <i>Cancer Medicine</i> , 2016, 5, 159-168.	2.8	61
48	Characterization of FN1–FGFR1 and novel FN1–FGF1 fusion genes in a large series of phosphaturic mesenchymal tumors. <i>Modern Pathology</i> , 2016, 29, 1335-1346.	5.5	139
49	Genomewide copy number analysis of Malignant mesothelioma identified chromosomal instability in the aggressive subgroup. <i>Modern Pathology</i> , 2016, 29, 1070-1082.	5.5	28
50	Angiosarcoma arising in a chronically lymphedematous leg: An unusual presentation of Stewart-Treves syndrome. <i>Dermatologica Sinica</i> , 2016, 34, 62-63.	0.5	0
51	Transcriptomic reappraisal identifies <i>MGLL</i> overexpression as an unfavorable prognosticator in primary gastrointestinal stromal tumors. <i>Oncotarget</i> , 2016, 7, 49986-49997.	1.8	12
52	Cutaneous syncytial myoepithelioma: A case report with emphasis on the differential diagnosis of problematic dermal tumors. <i>Oncology Letters</i> , 2015, 9, 2275-2277.	1.8	15
53	Expression of retinoic acid-binding proteins and retinoic acid receptors in sebaceous cell carcinoma of the eyelids. <i>BMC Ophthalmology</i> , 2015, 15, 142.	1.4	5
54	Cytopathologic features of epithelioid inflammatory myofibroblastic sarcoma with correlation of histopathology, immunohistochemistry, and molecular cytogenetic analysis. <i>Cancer Cytopathology</i> , 2015, 123, 495-504.	2.4	21

#	ARTICLE	IF	CITATIONS
55	Galectin-1 Dysregulation Independently Predicts Disease Specific Survival in Bladder Urothelial Carcinoma. <i>Journal of Urology</i> , 2015, 193, 1002-1008.	0.4	23
56	Skp2-Dependent Ubiquitination and Activation of LKB1 Is Essential for Cancer Cell Survival under Energy Stress. <i>Molecular Cell</i> , 2015, 57, 1022-1033.	9.7	97
57	NAB2-STAT6 fusion types account for clinicopathological variations in solitary fibrous tumors. <i>Modern Pathology</i> , 2015, 28, 1324-1335.	5.5	133
58	Phosphorylated p70S6K expression is an independent prognosticator for patients with esophageal squamous cell carcinoma. <i>Surgery</i> , 2015, 157, 570-580.	1.9	39
59	The cAMP responsive element binding protein 1 transactivates epithelial membrane protein 2, a potential tumor suppressor in the urinary bladder urothelial carcinoma. <i>Oncotarget</i> , 2015, 6, 9220-9239.	1.8	30
60	<i>CEBPD</i> amplification and overexpression in urothelial carcinoma: a driver of tumor metastasis indicating adverse prognosis. <i>Oncotarget</i> , 2015, 6, 31069-31084.	1.8	35
61	Impact of late toxicities on quality of life for survivors of nasopharyngeal carcinoma. <i>BMC Cancer</i> , 2014, 14, 856.	2.6	34
62	Overexpression of REG4 confers an independent negative prognosticator in rectal cancers receiving concurrent chemoradiotherapy. <i>Journal of Surgical Oncology</i> , 2014, 110, 1002-1010.	1.7	13
63	Angiomatoid fibrous histiocytoma: clinicopathological and molecular characterisation with emphasis on variant histomorphology. <i>Journal of Clinical Pathology</i> , 2014, 67, 210-215.	2.0	53
64	Effect of Age-Related Cartilage Turnover on Serum C-Telopeptide of Collagen Type II and Osteocalcin Levels in Growing Rabbits with and without Surgically Induced Osteoarthritis. <i>BioMed Research International</i> , 2014, 2014, 1-9.	1.9	7
65	<i>AMACR</i> Amplification in Myxofibrosarcomas: A Mechanism of Overexpression That Promotes Cell Proliferation with Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2014, 20, 6141-6152.	7.0	32
66	Overexpression of CDC28 protein kinase regulatory subunit 1B confers an independent prognostic factor in nasopharyngeal carcinoma. <i>Apmis</i> , 2014, 122, 206-214.	2.0	10
67	Fibroblast growth factor receptor 2 overexpression is predictive of poor prognosis in rectal cancer patients receiving neoadjuvant chemoradiotherapy. <i>Journal of Clinical Pathology</i> , 2014, 67, 1056-1061.	2.0	23
68	TOP2A overexpression as a poor prognostic factor in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 179-187.	1.8	72
69	Deficiency in expression and epigenetic DNA Methylation of ASS1 gene in nasopharyngeal carcinoma: negative prognostic impact and therapeutic relevance. <i>Tumor Biology</i> , 2014, 35, 161-169.	1.8	54
70	Fibronectin overexpression is associated with latent membrane protein 1 expression and has independent prognostic value for nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2014, 35, 1703-1712.	1.8	40
71	Clinical Aggressiveness of Myxofibrosarcomas Associates with Down-Regulation of p12CDK2AP1: Prognostic Implication of a Putative Tumor Suppressor that Induces Cell Cycle Arrest and Apoptosis Via Mitochondrial Pathway. <i>Annals of Surgical Oncology</i> , 2014, 21, 711-720.	1.5	2
72	Downregulated <i>MTAP</i> expression in myxofibrosarcoma: A characterization of inactivating mechanisms, tumor suppressive function, and therapeutic relevance. <i>Oncotarget</i> , 2014, 5, 11428-11441.	1.8	21

#	ARTICLE	IF	CITATIONS
73	<i>AMACR</i> amplification and overexpression in primary imatinib-naïve gastrointestinal stromal tumors: a driver of cell proliferation indicating adverse prognosis. <i>Oncotarget</i> , 2014, 5, 11588-11603.	1.8	22
74	Nicotinamide N-methyltransferase overexpression is associated with Akt phosphorylation and indicates worse prognosis in patients with nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 3923-3931.	1.8	35
75	Clinicopathological significance of HuR expression in gallbladder carcinoma: with special emphasis on the implications of its nuclear and cytoplasmic expression. <i>Tumor Biology</i> , 2013, 34, 3059-3069.	1.8	18
76	Fatty acid synthase overexpression confers an independent prognosticator and associates with radiation resistance in nasopharyngeal carcinoma. <i>Tumor Biology</i> , 2013, 34, 759-768.	1.8	34
77	SKP2 overexpression is associated with a poor prognosis of rectal cancer treated with chemoradiotherapy and represents a therapeutic target with high potential. <i>Tumor Biology</i> , 2013, 34, 1107-1117.	1.8	34
78	Prognostic value of quality of life measured after treatment on subsequent survival in patients with nasopharyngeal carcinoma. <i>Quality of Life Research</i> , 2013, 22, 715-723.	3.1	21
79	E2F transcription factor 1 overexpression as a poor prognostic factor in patients with nasopharyngeal carcinomas. <i>Biomarkers and Genomic Medicine</i> , 2013, 5, 23-30.	0.2	0
80	<i>ASS1</i> as a Novel Tumor Suppressor Gene in Myxofibrosarcomas: Aberrant Loss via Epigenetic DNA Methylation Confers Aggressive Phenotypes, Negative Prognostic Impact, and Therapeutic Relevance. <i>Clinical Cancer Research</i> , 2013, 19, 2861-2872.	7.0	118
81	Dedifferentiated liposarcoma with homologous lipoblastic differentiation: expanding the spectrum to include low-grade tumours. <i>Histopathology</i> , 2013, 62, 702-710.	2.9	23
82	Overexpression of Rap-1A Indicates a Poor Prognosis for Oral Cavity Squamous Cell Carcinoma and Promotes Tumor Cell Invasion via Aurora-A Modulation. <i>American Journal of Pathology</i> , 2013, 182, 516-528.	3.8	37
83	EBV-encoded small RNA1 and nonresolving inflammation in rheumatoid arthritis. <i>Kaohsiung Journal of Medical Sciences</i> , 2013, 29, 606-610.	1.9	6
84	Loss of epithelial membrane protein-2 expression confers an independent prognosticator in gallbladder carcinoma. <i>Biomarkers and Genomic Medicine</i> , 2013, 5, 31-38.	0.2	1
85	Overexpression of thymidylate synthetase confers an independent prognostic indicator in nasopharyngeal carcinoma. <i>Experimental and Molecular Pathology</i> , 2013, 95, 83-90.	2.1	34
86	The E2F Transcription Factor 1 Transactivates Stathmin 1 in Hepatocellular Carcinoma. <i>Annals of Surgical Oncology</i> , 2013, 20, 4041-4054.	1.5	53
87	IGFBP-5 overexpression as a poor prognostic factor in patients with urothelial carcinomas of upper urinary tracts and urinary bladder. <i>Journal of Clinical Pathology</i> , 2013, 66, 573-582.	2.0	43
88	The Expression of Activin Receptor-Like Kinase 1 among Patients with Head and Neck Cancer. <i>Otolaryngology - Head and Neck Surgery</i> , 2013, 148, 965-973.	1.9	10
89	HuR cytoplasmic expression is associated with increased cyclin A expression and inferior disease-free survival in patients with gastrointestinal stromal tumours (GISTs). <i>Histopathology</i> , 2013, 63, 445-454.	2.9	7
90	Dedifferentiation in Gastrointestinal Stromal Tumor to an Anaplastic KIT-negative Phenotype. <i>American Journal of Surgical Pathology</i> , 2013, 37, 385-392.	3.7	90

#	ARTICLE	IF	CITATIONS
91	High Chromosomal Copy Number Alterations in Xp11 Translocation Renal Cell Carcinomas Detected by Array Comparative Genomic Hybridization Are Associated With Aggressive Behavior. <i>American Journal of Surgical Pathology</i> , 2013, 37, 1116-1119.	3.7	18
92	Rsf-1 expression in rectal cancer: with special emphasis on the independent prognostic value after neoadjuvant chemoradiation. <i>Journal of Clinical Pathology</i> , 2012, 65, 687-692.	2.0	24
93	Alpha-methylacyl coenzyme A racemase overexpression in gallbladder carcinoma confers an independent prognostic indicator. <i>Journal of Clinical Pathology</i> , 2012, 65, 309-314.	2.0	15
94	Expression of cyclin-dependent kinase 2-associated protein 1 confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>Journal of Clinical Pathology</i> , 2012, 65, 795-801.	2.0	8
95	Characterization of Gene Amplification-Driven SKP2 Overexpression in Myxofibrosarcoma: Potential Implications in Tumor Progression and Therapeutics. <i>Clinical Cancer Research</i> , 2012, 18, 1598-1610.	7.0	40
96	Associations of Rsf-1 overexpression with poor therapeutic response and worse survival in patients with nasopharyngeal carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 248-253.	2.0	32
97	Loss of epithelial membrane protein-2 expression confers an independent prognosticator in nasopharyngeal carcinoma: a cohort study. <i>BMJ Open</i> , 2012, 2, e000900.	1.9	11
98	Enhancer of Zeste Homolog 2 Overexpression in Nasopharyngeal Carcinoma: An Independent Poor Prognosticator That Enhances Cell Growth. <i>International Journal of Radiation Oncology Biology Physics</i> , 2012, 82, 597-604.	0.8	27
99	Rsf-1/HBXAP overexpression is independent of gene amplification and is associated with poor outcome in patients with urinary bladder urothelial carcinoma. <i>Journal of Clinical Pathology</i> , 2012, 65, 802-807.	2.0	15
100	Recurrent Amplification at 7q21.2 Targets CDK6 Gene in Primary Myxofibrosarcomas and Identifies CDK6 Overexpression as an Independent Adverse Prognosticator. <i>Annals of Surgical Oncology</i> , 2012, 19, 2716-2725.	1.5	23
101	The Skp2-SCF E3 Ligase Regulates Akt Ubiquitination, Glycolysis, Herceptin Sensitivity, and Tumorigenesis. <i>Cell</i> , 2012, 149, 1098-1111.	28.9	332
102	Peroxisome proliferator-activated receptors β /mitochondrial uncoupling protein 2 signaling protects against seizure-induced neuronal cell death in the hippocampus following experimental status epilepticus. <i>Journal of Neuroinflammation</i> , 2012, 9, 184.	7.2	54
103	HuR cytoplasmic expression is associated with increased cyclin A expression and poor outcome with upper urinary tract urothelial carcinoma. <i>BMC Cancer</i> , 2012, 12, 611.	2.6	37
104	EGFR Nuclear Import in Gallbladder Carcinoma: Nuclear Phosphorylated EGFR Upregulates iNOS Expression and Confers Independent Prognostic Impact. <i>Annals of Surgical Oncology</i> , 2012, 19, 443-454.	1.5	28
105	The expression and prognostic significance of hepatoma-derived growth factor in oral cancer. <i>Oral Oncology</i> , 2012, 48, 629-635.	1.5	31
106	Overexpression of a Chromatin Remodeling Factor, RSF-1/HBXAP, Correlates with Aggressive Oral Squamous Cell Carcinoma. <i>American Journal of Pathology</i> , 2011, 178, 2407-2415.	3.8	44
107	Rsf-1/HBXAP overexpression is associated with disease-specific survival of patients with gallbladder carcinoma. <i>Apmis</i> , 2011, 119, 808-814.	2.0	22
108	CKS1B Overexpression Implicates Clinical Aggressiveness of Hepatocellular Carcinomas but Not p27Kip1 Protein Turnover: an Independent Prognosticator with Potential p27Kip1-Independent Oncogenic Attributes?. <i>Annals of Surgical Oncology</i> , 2010, 17, 907-922.	1.5	26

#	ARTICLE	IF	CITATIONS
109	Prognostic Implication of Ezrin Overexpression in Myxofibrosarcomas. <i>Annals of Surgical Oncology</i> , 2010, 17, 3212-3219.	1.5	39
110	Deciphering the transcriptional complex critical for RhoA gene expression and cancer metastasis. <i>Nature Cell Biology</i> , 2010, 12, 457-467.	10.3	190
111	Prognostic implication of MET overexpression in myxofibrosarcomas: an integrative array comparative genomic hybridization, real-time quantitative PCR, immunoblotting, and immunohistochemical analysis. <i>Modern Pathology</i> , 2010, 23, 1379-1392.	5.5	31
112	Ezrin overexpression in gastrointestinal stromal tumors: an independent adverse prognosticator associated with the non-gastric location. <i>Modern Pathology</i> , 2009, 22, 1351-1360.	5.5	25
113	Effect of S-Phase Kinase-Associated Protein 2 Expression on Distant Metastasis and Survival in Nasopharyngeal Carcinoma Patients. <i>International Journal of Radiation Oncology Biology Physics</i> , 2009, 73, 202-207.	0.8	36
114	Homozygous Deletion of <i>MTAP</i> Gene as a Poor Prognosticator in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2009, 15, 6963-6972.	7.0	41
115	Quality of Life and Survival Outcome for Patients With Nasopharyngeal Carcinoma Receiving Three-Dimensional Conformal Radiotherapy vs. Intensity-Modulated Radiotherapy—A Longitudinal Study. <i>International Journal of Radiation Oncology Biology Physics</i> , 2008, 72, 356-364.	0.8	176
116	Flow Cytometric Analysis of DNA Ploidy and S-Phase Fraction in Primary Localized Myxofibrosarcoma: Correlations with Clinicopathological Factors, Skp2 Expression, and Patient Survival. <i>Annals of Surgical Oncology</i> , 2008, 15, 2239-2249.	1.5	10
117	Heat Shock Protein 90 Overexpression Independently Predicts Inferior Disease-Free Survival with Differential Expression of the α and β Isoforms in Gastrointestinal Stromal Tumors. <i>Clinical Cancer Research</i> , 2008, 14, 7822-7831.	7.0	40
118	Malignant Diffuse-type Tenosynovial Giant Cell Tumors. <i>American Journal of Surgical Pathology</i> , 2008, 32, 587-599.	3.7	57
119	Hamartoma of the Breast: An Underrecognized Disease?. <i>Tumori</i> , 2008, 94, 114-115.	1.1	5
120	Immunohistochemical expression of epidermal growth factor receptor and cyclooxygenase-2 in pediatric nasopharyngeal carcinomas: No significant correlations with clinicopathological variables and treatment outcomes. <i>International Journal of Pediatric Otorhinolaryngology</i> , 2007, 71, 447-455.	1.0	15
121	Skp2 is an independent prognosticator of gallbladder carcinoma among p27Kip1-interacting cell cycle regulators: an immunohistochemical study of 62 cases by tissue microarray. <i>Modern Pathology</i> , 2007, 20, 497-507.	5.5	43
122	Immunohistochemical Expression of p16INK4A, Ki-67, and Mcm2 Proteins in Gastrointestinal Stromal Tumors: Prognostic Implications and Correlations with Risk Stratification of NIH Consensus Criteria. <i>Annals of Surgical Oncology</i> , 2006, 13, 1633-1644.	1.5	46
123	Primary myxoid solitary fibrous tumor involving the seminal vesicle. <i>Pathology International</i> , 2006, 56, 642-644.	1.3	14
124	Primary lymphoepithelioma-like carcinoma of minor salivary gland: A case report with immunohistochemical and in situ hybridization studies. <i>Head and Neck</i> , 2006, 28, 182-186.	2.0	17
125	Prognostic factors of myxofibrosarcomas: Implications of margin status, tumor necrosis, and mitotic rate on survival. <i>Journal of Surgical Oncology</i> , 2006, 93, 294-303.	1.7	72
126	Skp2 Overexpression Is Highly Representative of Intrinsic Biological Aggressiveness and Independently Associated with Poor Prognosis in Primary Localized Myxofibrosarcomas. <i>Clinical Cancer Research</i> , 2006, 12, 487-498.	7.0	50

#	ARTICLE	IF	CITATIONS
127	Distant metastasis in retroperitoneal dedifferentiated liposarcoma is rare and rapidly fatal: a clinicopathological study with emphasis on the low-grade myxofibrosarcoma-like pattern as an early sign of dedifferentiation. <i>Modern Pathology</i> , 2005, 18, 976-984.	5.5	73
128	Nonrandom cell-cycle timing of a somatic chromosomal translocation: The t(X;17) of alveolar soft-part sarcoma occurs in G2. <i>Genes Chromosomes and Cancer</i> , 2005, 44, 170-176.	2.8	20
129	Ewing Sarcomas With <i>p53</i> Mutation or <i>p16/p14ARF</i> Homozygous Deletion: A Highly Lethal Subset Associated With Poor Chemoresponse. <i>Journal of Clinical Oncology</i> , 2005, 23, 548-558.	1.6	235
130	Low-grade myxofibrosarcoma: a clinicopathologic analysis of 49 cases treated at a single institution with simultaneous assessment of the efficacy of 3-tier and 4-tier grading systems. <i>Human Pathology</i> , 2004, 35, 612-621.	2.0	201
131	Immunohistochemical and Ultrastructural Comparative Study of External Lamina Structure in 31 Cases of Cellular, Classical, and Melanotic Schwannomas. <i>Applied Immunohistochemistry and Molecular Morphology</i> , 2004, 12, 50-58.	1.2	23
132	Molecular Detection of JAZF1-JJAZ1 Gene Fusion in Endometrial Stromal Neoplasms with Classic and Variant Histology. <i>American Journal of Surgical Pathology</i> , 2004, 28, 224-232.	3.7	117
133	Molecular Variability of TLS - CHOP Structure Shows No Significant Impact on the Level of Adipogenesis: A Comparative Ultrastructural and RT-PCR Analysis of 14 Cases of Myxoid/Round Cell Liposarcomas. <i>Ultrastructural Pathology</i> , 2003, 27, 217-226.	0.9	17
134	Sinonasal Smooth Muscle Cell Tumors. <i>Archives of Pathology and Laboratory Medicine</i> , 2003, 127, 297-304.	2.5	38
135	Epithelioid Variant of Pleomorphic Liposarcoma: A Comparative Immunohistochemical and Ultrastructural Analysis of Six Cases with Emphasis on Overlapping Features with Epithelial Malignancies. <i>Ultrastructural Pathology</i> , 2002, 26, 299-308.	0.9	37
136	Primary mesenchymal chondrosarcoma of the lung. <i>Annals of Thoracic Surgery</i> , 2002, 73, 1960-1962.	1.3	24
137	Solitary fibrous tumor of the abdominal wall: a report of two cases with immunohistochemical, flow cytometric, and ultrastructural studies and literature review. <i>Apmis</i> , 2002, 110, 253-262.	2.0	28
138	Superficial collagenous fibroma: Immunohistochemical, ultrastructural, and flow cytometric study of three cases, including one pemphigus vulgaris patient with a dermal mass. <i>Apmis</i> , 2002, 110, 283-289.	2.0	13
139	Primary Yolk Sac Tumor of the Urachus. <i>Archives of Pathology and Laboratory Medicine</i> , 2002, 126, 1106-1109.	2.5	24
140	Isolated intracranial Rosai-Dorfman disease: Case report and literature review. <i>Pathology International</i> , 1998, 48, 396-402.	1.3	35
141	Clinical significance of recombinant RANBP2-ALK oncogene to radiotherapy in a case of locally advanced rectal epithelioid inflammatory myofibroblastic sarcoma: case report and review of literature. <i>Therapeutic Radiology and Oncology</i> , 0, 4, 9-9.	0.2	0