

Peter G JÃ,rgensen

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/2978123/publications.pdf>

Version: 2024-02-01

63
papers

1,670
citations

361413

20
h-index

315739

38
g-index

64
all docs

64
docs citations

64
times ranked

2503
citing authors

#	ARTICLE	IF	CITATIONS
1	Accuracy, analysis time, and reproducibility of dedicated 4D echocardiographic left atrial volume quantification software. <i>International Journal of Cardiovascular Imaging</i> , 2022, 38, 1277-1288.	1.5	2
2	Acute changes in plasma glucose increases left ventricular systolic function in insulin-treated patients with type 2 diabetes and controls. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 1123-1131.	4.4	3
3	Potential role of conventional and speckle-tracking echocardiography in the screening of structural and functional cardiac abnormalities in elderly individuals: Baseline echocardiographic findings from the LOOP study. <i>PLoS ONE</i> , 2022, 17, e0269475.	2.5	2
4	Hypoglycaemia and rebound hyperglycaemia increase left ventricular systolic function in patients with type 1 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 2027-2037.	4.4	4
5	Diagnostic and prognostic value of the electrocardiogram in stable outpatients with type 2 diabetes. <i>Scandinavian Cardiovascular Journal</i> , 2022, 56, 256-263.	1.2	0
6	Sex differences in the association between myocardial function and prognosis in type 1 diabetes without known heart disease: the Thousand & 1 Study. <i>European Heart Journal Cardiovascular Imaging</i> , 2021, 22, 1017-1025.	1.2	4
7	Echocardiographic predictors of cardiovascular morbidity and mortality in women from the general population. <i>European Heart Journal Cardiovascular Imaging</i> , 2021, 22, 1026-1034.	1.2	10
8	Prognostic Value of Early Systolic Lengthening by Strain Imaging in Type 2 Diabetes. <i>Journal of the American Society of Echocardiography</i> , 2021, 34, 127-135.	2.8	10
9	Left ventricular systolic ejection time is an independent predictor of all-cause mortality in heart failure with reduced ejection fraction. <i>European Journal of Heart Failure</i> , 2021, 23, 240-249.	7.1	17
10	Relationship between peripheral neuropathy, diastolic function and adverse cardiovascular outcome in individuals with type 1 diabetes mellitus without known cardiovascular disease: Results from the Thousand & 1 Study. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 158-165.	4.4	4
11	Targeting epicardial adipose tissue with exercise, diet, bariatric surgery or pharmaceutical interventions: A systematic review and meta-analysis. <i>Obesity Reviews</i> , 2021, 22, e13136.	6.5	43
12	Prognostic value of left ventricular mitral annular longitudinal displacement obtained by tissue Doppler imaging in patients with heart failure with reduced ejection fraction. <i>Open Heart</i> , 2021, 8, e001494.	2.3	0
13	Layer-specific global longitudinal strain obtained by speckle tracking echocardiography for predicting heart failure and cardiovascular death following STEMI treated with primary PCI. <i>International Journal of Cardiovascular Imaging</i> , 2021, 37, 2207-2215.	1.5	5
14	Prognostic and comparative performance of cardiovascular risk markers in patients with type 2 diabetes. <i>Journal of Diabetes</i> , 2021, 13, 754-763.	1.8	2
15	The prognostic value of myocardial deformational patterns on all-cause mortality is modified by ischemic cardiomyopathy in patients with heart failure. <i>International Journal of Cardiovascular Imaging</i> , 2021, 37, 3137-3144.	1.5	3
16	Echocardiographic predictors of long-term adverse cardiovascular outcomes in participants with and without diabetes mellitus: A follow-up analysis of the Copenhagen City Heart Study. <i>Diabetic Medicine</i> , 2021, 38, e14627.	2.3	4
17	Prognostic value of right ventricular echocardiographic measures in patients with heart failure with reduced ejection fraction. <i>Journal of Clinical Ultrasound</i> , 2021, 49, 903-913.	0.8	7
18	Prognostic utility of diastolic dysfunction and speckle tracking echocardiography in heart failure with reduced ejection fraction. <i>ESC Heart Failure</i> , 2020, 7, 148-158.	3.1	11

#	ARTICLE	IF	CITATIONS
19	MR-proANP and incident cardiovascular disease in patients with type 2 diabetes with and without heart failure with preserved ejection fraction. <i>Cardiovascular Diabetology</i> , 2020, 19, 180.	6.8	7
20	The many-faced hurdle of cardiac involvement in diabetes: where to focus?. <i>Therapeutic Advances in Endocrinology and Metabolism</i> , 2020, 11, 204201882097044.	3.2	0
21	Hypoglycaemia and cardiac arrhythmias in diabetes. <i>Therapeutic Advances in Endocrinology and Metabolism</i> , 2020, 11, 204201882091180.	3.2	25
22	Epicardial adipose tissue: an emerging biomarker of cardiovascular complications in type 2 diabetes?. <i>Therapeutic Advances in Endocrinology and Metabolism</i> , 2020, 11, 204201882092882.	3.2	38
23	Cardiac adaptation in hibernating, free-ranging Scandinavian Brown Bears (<i>Ursus arctos</i>). <i>Scientific Reports</i> , 2020, 10, 247.	3.3	10
24	Cardiovascular prognostic value of echocardiography and N terminal pro B-type natriuretic peptide in type 1 diabetes: the Thousand & 1 Study. <i>European Journal of Endocrinology</i> , 2020, 182, 481-488.	3.7	10
25	Ratio of Transmitral Early Filling Velocity to Early Diastolic Strain Rate Predicts All-Cause Mortality in Heart Failure with Reduced Ejection Fraction. <i>Journal of Cardiac Failure</i> , 2019, 25, 877-885.	1.7	12
26	Echocardiography improves prediction of major adverse cardiovascular events in a population with type 1 diabetes and without known heart disease: the Thousand & 1 Study. <i>Diabetologia</i> , 2019, 62, 2354-2364.	6.3	23
27	Epicardial adipose tissue predicts incident cardiovascular disease and mortality in patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2019, 18, 114.	6.8	57
28	Usefulness of layer-specific strain in diagnosis of coronary artery disease in patients with stable angina pectoris. <i>International Journal of Cardiovascular Imaging</i> , 2019, 35, 1989-1999.	1.5	19
29	Epicardial and pericardial adipose tissues are associated with reduced diastolic and systolic function in type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 2006-2011.	4.4	44
30	An echocardiographic substrate for dyspnea identifies high risk patients with type 2 diabetes. <i>International Journal of Cardiology</i> , 2019, 289, 119-124.	1.7	3
31	Prognostic value of ratio of transmitral early filling velocity to early diastolic strain rate in patients with Type 2 diabetes. <i>European Heart Journal Cardiovascular Imaging</i> , 2019, 20, 1171-1178.	1.2	15
32	Utility of left atrial strain for predicting atrial fibrillation following ischemic stroke. <i>International Journal of Cardiovascular Imaging</i> , 2019, 35, 1605-1613.	1.5	27
33	Left ventricular concentric geometry predicts incident diabetes mellitus independent of established risk factors in the general population: the Copenhagen City Heart Study. <i>Cardiovascular Diabetology</i> , 2019, 18, 37.	6.8	5
34	Ratio of Transmitral Early Filling Velocity to Early Diastolic Strain Rate as a Predictor of Cardiovascular Morbidity and Mortality Following Acute Coronary Syndrome. <i>American Journal of Cardiology</i> , 2019, 123, 1776-1782.	1.6	7
35	Prevalence of heart failure and the diagnostic value of MR-proANP in outpatients with type 2 diabetes. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 736-740.	4.4	16
36	Predictive value of echocardiography in Type 2 diabetes. <i>European Heart Journal Cardiovascular Imaging</i> , 2019, 20, 687-693.	1.2	25

#	ARTICLE	IF	CITATIONS
37	Prognostic Value of Left Atrial Functional Measures in Heart Failure With Reduced Ejection Fraction. <i>Journal of Cardiac Failure</i> , 2019, 25, 87-96.	1.7	18
38	Post-systolic shortening predicts heart failure following acute coronary syndrome. <i>International Journal of Cardiology</i> , 2019, 276, 191-197.	1.7	14
39	Association between layer-specific global longitudinal strain and adverse outcomes following acute coronary syndrome. <i>European Heart Journal Cardiovascular Imaging</i> , 2018, 19, 1334-1342.	1.2	43
40	Layer-specific global longitudinal strain reveals impaired cardiac function in patients with reversible ischemia. <i>Echocardiography</i> , 2018, 35, 632-642.	0.9	17
41	Intestinal barrier integrity and inflammatory bowel disease: Stem cell-based approaches to regenerate the barrier. <i>Journal of Tissue Engineering and Regenerative Medicine</i> , 2018, 12, 923-935.	2.7	48
42	Second generation drug-eluting stents versus bare-metal stents for percutaneous coronary intervention of the proximal left anterior descending artery: An analysis of the BASKET-PROVE I and II trials. <i>Catheterization and Cardiovascular Interventions</i> , 2018, 91, 867-873.	1.7	4
43	Global longitudinal strain corrected by RR interval is a superior predictor of all-cause mortality in patients with systolic heart failure and atrial fibrillation. <i>ESC Heart Failure</i> , 2018, 5, 311-318.	3.1	18
44	Presence of micro- and macroalbuminuria and the association with cardiac mechanics in patients with type 2 diabetes. <i>European Heart Journal Cardiovascular Imaging</i> , 2018, 19, 1034-1041.	1.2	23
45	Echocardiographic Predictors of Mortality in Women With Heart Failure With Reduced Ejection Fraction. <i>Circulation: Cardiovascular Imaging</i> , 2018, 11, e008031.	2.6	20
46	Burden of Uncontrolled Metabolic Risk Factors and Left Ventricular Structure and Function in Patients With Type 2 Diabetes Mellitus. <i>Journal of the American Heart Association</i> , 2018, 7, e008856.	3.7	16
47	Left ventricular filling pressure by septal and lateral E/e ² equally predict cardiovascular events in the general population. <i>International Journal of Cardiovascular Imaging</i> , 2017, 33, 653-661.	1.5	11
48	Global Longitudinal Strain by Echocardiography Predicts Long-Term Risk of Cardiovascular Morbidity and Mortality in a Low-Risk General Population. <i>Circulation: Cardiovascular Imaging</i> , 2017, 10, .	2.6	270
49	Effect of exercise combined with glucagon-like peptide-1 receptor agonist treatment on cardiac function: randomized double-blind placebo-controlled clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2017, 19, 1040-1044.	4.4	26
50	Cardiac Adaptations to High-Intensity Aerobic Training in Premenopausal and Recent Postmenopausal Women: The Copenhagen Women Study. <i>Journal of the American Heart Association</i> , 2017, 6, .	3.7	18
51	Frequency of Cardiac Death and Stent Thrombosis in Patients With Chronic Obstructive Pulmonary Disease Undergoing Percutaneous Coronary Intervention (from the BASKET-PROVE I and II Trials). <i>American Journal of Cardiology</i> , 2017, 119, 14-19.	1.6	12
52	Echocardiographic quantification of systolic function during atrial fibrillation: probing the "ten heart cycles" rule. <i>Future Cardiology</i> , 2016, 12, 159-165.	1.2	6
53	Cholesterol remnants and triglycerides are associated with decreased myocardial function in patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2016, 15, 137.	6.8	25
54	Impact of type 2 diabetes and duration of type 2 diabetes on cardiac structure and function. <i>International Journal of Cardiology</i> , 2016, 221, 114-121.	1.7	39

#	ARTICLE	IF	CITATIONS
55	Abnormal echocardiography in patients with type 2 diabetes and relation to symptoms and clinical characteristics. <i>Diabetes and Vascular Disease Research</i> , 2016, 13, 321-330.	2.0	42
56	Predicting Paroxysmal Atrial Fibrillation in Cerebrovascular Ischemia Using Tissue Doppler Imaging and Speckle Tracking Echocardiography. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2016, 25, 350-359.	1.6	16
57	Global Longitudinal Strain Is a Superior Predictor of All-Cause Mortality in Heart Failure With Reduced Ejection Fraction. <i>JACC: Cardiovascular Imaging</i> , 2015, 8, 1351-1359.	5.3	288
58	Global Longitudinal Strain Is Not Impaired in Type 1 Diabetes Patients Without Albuminuria. <i>JACC: Cardiovascular Imaging</i> , 2015, 8, 400-410.	5.3	86
59	Diastolic myocardial dysfunction by tissue Doppler imaging predicts mortality in patients with cerebral infarction. <i>International Journal of Cardiovascular Imaging</i> , 2015, 31, 1413-1422.	1.5	11
60	Plasma pro-brain natriuretic peptide and electrocardiographic changes in combination improve risk prediction in persons without known heart disease. <i>International Journal of Cardiology</i> , 2015, 201, 104-109.	1.7	2
61	Low cardiac output as physiological phenomenon in hibernating, free-ranging Scandinavian brown bears (<i>Ursus arctos</i>) – an observational study. <i>Cardiovascular Ultrasound</i> , 2014, 12, 36.	1.6	20
62	Prevalence of systolic and diastolic dysfunction in patients with type 1 diabetes without known heart disease: the Thousand & 1 Study. <i>Diabetologia</i> , 2014, 57, 672-680.	6.3	71
63	Electrocardiographic Changes Improve Risk Prediction in Asymptomatic Persons Age 65 Years or Above Without Cardiovascular Disease. <i>Journal of the American College of Cardiology</i> , 2014, 64, 898-906.	2.8	32