

Michel Probst

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3060692/publications.pdf>

Version: 2024-02-01

176
papers

8,417
citations

41344

49
h-index

54911

84
g-index

179
all docs

179
docs citations

179
times ranked

7718
citing authors

#	ARTICLE	IF	CITATIONS
1	Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. <i>World Psychiatry</i> , 2017, 16, 308-315.	10.4	600
2	Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. <i>World Psychiatry</i> , 2016, 15, 166-174.	10.4	487
3	Metabolic Syndrome and Metabolic Abnormalities in Bipolar Disorder: A Meta-Analysis of Prevalence Rates and Moderators. <i>American Journal of Psychiatry</i> , 2013, 170, 265-274.	7.2	336
4	A systematic review of correlates of physical activity in patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2012, 125, 352-362.	4.5	255
5	Metabolic syndrome and metabolic abnormalities in patients with major depressive disorder: a meta-analysis of prevalences and moderating variables. <i>Psychological Medicine</i> , 2014, 44, 2017-2028.	4.5	223
6	A meta-analysis of cardio-metabolic abnormalities in drug naïve, first-episode and multi-episode patients with schizophrenia versus general population controls. <i>World Psychiatry</i> , 2013, 12, 240-250.	10.4	220
7	The Body Attitude Test for Patients with an Eating Disorder: Psychometric Characteristics of a New Questionnaire. <i>Eating Disorders</i> , 1995, 3, 133-144.	3.0	200
8	Prevalence and predictors of treatment dropout from physical activity interventions in schizophrenia: a meta-analysis. <i>General Hospital Psychiatry</i> , 2016, 39, 15-23.	2.4	172
9	Cardiorespiratory Fitness in Severe Mental Illness: A Systematic Review and Meta-analysis. <i>Sports Medicine</i> , 2017, 47, 343-352.	6.5	170
10	Physical activity and suicidal ideation: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2018, 225, 438-448.	4.1	140
11	TYPE 2 DIABETES IN PATIENTS WITH MAJOR DEPRESSIVE DISORDER: A META-ANALYSIS OF PREVALENCE ESTIMATES AND PREDICTORS. <i>Depression and Anxiety</i> , 2015, 32, 763-773.	4.1	138
12	The importance of self-determined motivation towards physical activity in patients with schizophrenia. <i>Psychiatry Research</i> , 2013, 210, 812-818.	3.3	133
13	Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. <i>Psychiatry Research</i> , 2013, 207, 25-32.	3.3	131
14	Considering a frame of reference for physical activity research related to the cardiometabolic risk profile in schizophrenia. <i>Psychiatry Research</i> , 2010, 177, 271-279.	3.3	125
15	International Organization of Physical Therapy in Mental Health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia. <i>Disability and Rehabilitation</i> , 2012, 34, 1-12.	1.8	122
16	Associations between sedentary behaviour and metabolic parameters in patients with schizophrenia. <i>Psychiatry Research</i> , 2012, 200, 73-78.	3.3	120
17	Systematic Review of the Benefits of Physical Therapy Within a Multidisciplinary Care Approach for People With Schizophrenia. <i>Physical Therapy</i> , 2012, 92, 11-23.	2.4	114
18	Relationships between obesity, functional exercise capacity, physical activity participation and physical self-perception in people with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2011, 123, 423-430.	4.5	113

#	ARTICLE	IF	CITATIONS
19	Physical activity and sedentary behavior in people with bipolar disorder: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2016, 201, 145-152.	4.1	109
20	A review of physical activity correlates in patients with bipolar disorder. <i>Journal of Affective Disorders</i> , 2013, 145, 285-291.	4.1	108
21	Promotion of cardiorespiratory fitness in schizophrenia: a clinical overview and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 131-143.	4.5	108
22	Global physical activity levels among people living with HIV: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2018, 40, 388-397.	1.8	100
23	Neurocognition in clinical high risk young adults who did or did not convert to a first schizophrenic psychosis: A meta-analysis. <i>Schizophrenia Research</i> , 2013, 149, 48-55.	2.0	97
24	Yoga in schizophrenia: a systematic review of randomised controlled trials. <i>Acta Psychiatrica Scandinavica</i> , 2012, 126, 12-20.	4.5	94
25	State anxiety, psychological stress and positive well-being responses to yoga and aerobic exercise in people with schizophrenia: a pilot study. <i>Disability and Rehabilitation</i> , 2011, 33, 684-689.	1.8	91
26	Lack of physical activity during leisure time contributes to an impaired health related quality of life in patients with schizophrenia. <i>Schizophrenia Research</i> , 2011, 129, 122-127.	2.0	91
27	Barriers to and Facilitators of Physical Activity Among Persons With Schizophrenia: A Survey of Physical Therapists. <i>Psychiatric Services</i> , 2014, 65, 693-696.	2.0	88
28	Adopting and maintaining physical activity behaviours in people with severe mental illness: The importance of autonomous motivation. <i>Preventive Medicine</i> , 2015, 81, 216-220.	3.4	86
29	Prevalence and Predictors of Type 2 Diabetes Mellitus in People With Bipolar Disorder. <i>Journal of Clinical Psychiatry</i> , 2015, 76, 1490-1499.	2.2	85
30	Can cognitive behavioural therapy based strategies be integrated into physiotherapy for the prevention of chronic low back pain? A systematic review. <i>Disability and Rehabilitation</i> , 2013, 35, 1-10.	1.8	80
31	A systematic review of physical therapy interventions for patients with anorexia and bulimia nervosa. <i>Disability and Rehabilitation</i> , 2014, 36, 628-634.	1.8	80
32	The relationship between chronic physical conditions, multimorbidity and anxiety in the general population: A global perspective across 42 countries. <i>General Hospital Psychiatry</i> , 2017, 45, 1-6.	2.4	80
33	Negative symptoms are associated with lower autonomous motivation towards physical activity in people with schizophrenia. <i>Comprehensive Psychiatry</i> , 2015, 56, 128-132.	3.1	77
34	The transcending benefits of physical activity for individuals with schizophrenia: A systematic review and meta-ethnography. <i>Psychiatry Research</i> , 2014, 220, 11-19.	3.3	75
35	The functional exercise capacity is correlated with global functioning in patients with schizophrenia. <i>Acta Psychiatrica Scandinavica</i> , 2012, 125, 382-387.	4.5	74
36	The Prevalence of Metabolic Syndrome in Alcohol Use Disorders: A Systematic Review and Meta-analysis. <i>Alcohol and Alcoholism</i> , 2016, 51, 515-521.	1.6	72

#	ARTICLE	IF	CITATIONS
37	Physical activity and sedentary behaviour in outpatients with schizophrenia: A systematic review and meta-analysis. <i>International Journal of Therapy and Rehabilitation</i> , 2013, 20, 588-595.	0.3	69
38	Drive for thinness, affect regulation and physical activity in eating disorders: A daily life study. <i>Behaviour Research and Therapy</i> , 2007, 45, 1717-1734.	3.1	68
39	What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. <i>Archives of Gerontology and Geriatrics</i> , 2014, 59, 195-203.	3.0	67
40	Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. <i>Disability and Rehabilitation</i> , 2018, 40, 1618-1629.	1.8	65
41	Neurobiological effects of physical exercise in schizophrenia: a systematic review. <i>Disability and Rehabilitation</i> , 2014, 36, 1749-1754.	1.8	63
42	Psychomotor Therapy and Psychiatry: What's in a Name?. <i>The Open Complementary Medicine Journal</i> , 2010, 2, 105-113.	1.5	57
43	A systematic review on physical therapy interventions for patients with binge eating disorder. <i>Disability and Rehabilitation</i> , 2013, 35, 2191-2196.	1.8	55
44	Markers of inflammation in schizophrenia: association vs. causation. <i>World Psychiatry</i> , 2014, 13, 189-192.	10.4	54
45	Reliability, minimal detectable changes, practice effects and correlates of the 6-min walk test in patients with schizophrenia. <i>Psychiatry Research</i> , 2011, 187, 62-67.	3.3	53
46	Body composition of anorexia nervosa patients assessed by underwater weighing and skinfold-thickness measurements before and after weight gain. <i>American Journal of Clinical Nutrition</i> , 2001, 73, 190-197.	4.7	52
47	Association of the metabolic syndrome with physical activity performance in patients with schizophrenia. <i>Diabetes and Metabolism</i> , 2011, 37, 318-323.	2.9	52
48	Physiotherapists can help implement physical activity programmes in clinical practice. <i>British Journal of Psychiatry</i> , 2014, 204, 164-164.	2.8	52
49	What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. <i>Psychiatria Danubina</i> , 2015, 27, 210-24.	0.4	51
50	Effects of progressive muscle relaxation on state anxiety and subjective well-being in people with schizophrenia: a randomized controlled trial. <i>Clinical Rehabilitation</i> , 2011, 25, 567-575.	2.2	50
51	Cardiometabolic effects of physical activity interventions for people with schizophrenia. <i>Physical Therapy Reviews</i> , 2009, 14, 388-398.	0.8	48
52	Progressive muscle relaxation in persons with schizophrenia: a systematic review of randomized controlled trials. <i>Clinical Rehabilitation</i> , 2013, 27, 291-298.	2.2	47
53	Could autonomous motivation hold the key to successfully implementing lifestyle changes in affective disorders? A multicentre cross sectional study. <i>Psychiatry Research</i> , 2015, 228, 100-106.	3.3	47
54	Evaluation of body experience questionnaires in eating disorders in female patients (AN/BN) and nonclinical participants. <i>International Journal of Eating Disorders</i> , 2008, 41, 657-665.	4.0	46

#	ARTICLE	IF	CITATIONS
55	A Systematic Review of Physical Activity Correlates in Alcohol Use Disorders. Archives of Psychiatric Nursing, 2015, 29, 196-201.	1.4	45
56	The significance of body size estimation in eating disorders: Its relationship with clinical and psychological variables. , 1998, 24, 167-174.		42
57	Self-determination and stage of readiness to change physical activity behaviour in schizophrenia. Mental Health and Physical Activity, 2014, 7, 171-176.	1.8	41
58	Investigating the benefits of sport participation for individuals with schizophrenia: a systematic review. Psychiatria Danubina, 2015, 27, 2-13.	0.4	41
59	Physical Therapists's Ability to Identify Psychological Factors and Their Self-Reported Competence to Manage Chronic Low Back Pain. Physical Therapy, 2018, 98, 471-479.	2.4	40
60	Physical Activity Correlates in Persons with Binge Eating Disorder: A Systematic Review. European Eating Disorders Review, 2014, 22, 1-8.	4.1	37
61	Health related quality of life, physical fitness and physical activity participation in treatment-seeking obese persons with and without binge eating disorder. Psychiatry Research, 2014, 216, 97-102.	3.3	37
62	Associations between physical activity and the built environment in patients with schizophrenia: a multi-centre study. General Hospital Psychiatry, 2013, 35, 653-658.	2.4	36
63	Cardiorespiratory fitness levels and moderators in people with HIV: A systematic review and meta-analysis. Preventive Medicine, 2016, 93, 106-114.	3.4	36
64	Health-related quality of life and aerobic fitness in people with schizophrenia. International Journal of Mental Health Nursing, 2015, 24, 394-402.	3.8	35
65	Understanding the role of physiotherapists in schizophrenia: an international perspective from members of the International Organisation of Physical Therapists in Mental Health (IOPTMH). Journal of Mental Health, 2014, 23, 125-129.	1.9	34
66	Changes in physical activity, physical fitness, self-perception and quality of life following a 6-month physical activity counseling and cognitive behavioral therapy program in outpatients with binge eating disorder. Psychiatry Research, 2014, 219, 361-366.	3.3	34
67	Autonomous motivation is associated with the maintenance stage of behaviour change in people with affective disorders. Psychiatry Research, 2016, 240, 267-271.	3.3	33
68	Sedentary behaviour and sleep problems among 42,489 community-dwelling adults in six low- and middle-income countries. Journal of Sleep Research, 2018, 27, e12714.	3.2	33
69	Physical activity as a vital sign in patients with schizophrenia: Evidence and clinical recommendations. Schizophrenia Research, 2016, 170, 336-340.	2.0	32
70	Variability in Affective Activation Predicts Non-suicidal Self-harm in Eating Disorders. European Eating Disorders Review, 2013, 21, 143-147.	4.1	31
71	Eurofit test battery in patients with schizophrenia or schizoaffective disorder: Reliability and clinical correlates. European Psychiatry, 2012, 27, 416-421.	0.2	29
72	Effects of a group physical activity program on physical fitness and quality of life in individuals with schizophrenia. Mental Health and Physical Activity, 2014, 7, 155-162.	1.8	29

#	ARTICLE	IF	CITATIONS
73	The benefits of walking for individuals with schizophrenia spectrum disorders: A systematic review. <i>International Journal of Therapy and Rehabilitation</i> , 2014, 21, 410-420.	0.3	29
74	An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. <i>BMC Psychiatry</i> , 2013, 13, 5.	2.6	28
75	Diabetes, physical activity participation and exercise capacity in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2013, 67, 451-456.	1.8	28
76	A Cognitive- Behavioral Therapeutic Program for Patients With Obesity and Binge Eating Disorder. <i>Behavior Modification</i> , 2012, 36, 670-686.	1.6	27
77	The prevalence of diabetes mellitus type 2 in people with alcohol use disorders: a systematic review and large scale meta-analysis. <i>Psychiatry Research</i> , 2016, 246, 394-400.	3.3	27
78	What are the top 10 physical activity research questions in schizophrenia?. <i>Disability and Rehabilitation</i> , 2016, 38, 2235-2243.	1.8	27
79	A comparison of physical fitness in patients with bipolar disorder, schizophrenia and healthy controls. <i>Disability and Rehabilitation</i> , 2016, 38, 2047-2051.	1.8	27
80	Is autonomous motivation the key to maintaining an active lifestyle in first-episode psychosis?. <i>Microbial Biotechnology</i> , 2018, 12, 821-827.	1.7	27
81	The value of social support to encourage people with schizophrenia to engage in physical activity: an international insight from specialist mental health physiotherapists. <i>Journal of Mental Health</i> , 2014, 23, 256-260.	1.9	26
82	Reliability and clinical correlates of the Astrand-Rhyming sub-maximal exercise test in patients with schizophrenia or schizoaffective disorder. <i>Psychiatry Research</i> , 2014, 220, 778-783.	3.3	26
83	The International Organization of Physical Therapists working in Mental Health (IOPTMH). <i>Mental Health and Physical Activity</i> , 2012, 5, 20-21.	1.8	25
84	Associations between perceived neighbourhood environmental attributes and self-reported sitting time in patients with schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 215, 33-38.	3.3	25
85	Health-related physical fitness in patients with bipolar disorder vs. healthy controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2015, 177, 22-27.	4.1	25
86	Physical activity is associated with the physical, psychological, social and environmental quality of life in people with mental health problems in a low resource setting. <i>Psychiatry Research</i> , 2017, 258, 250-254.	3.3	25
87	Impact of antipsychotic medication on physical activity and physical fitness in adolescents: An exploratory study. <i>Psychiatry Research</i> , 2016, 242, 192-197.	3.3	23
88	Association of lifestyle-related factors and psychological factors on quality of life in people with schizophrenia. <i>Psychiatry Research</i> , 2018, 267, 382-393.	3.3	23
89	Sedentary Behavior in People Living With HIV: A Systematic Review and Meta-Analysis. <i>Journal of Physical Activity and Health</i> , 2017, 14, 571-577.	2.0	22
90	Assessing positive body image, body satisfaction, weight bias, and appearance comparison in emerging adults: A cross-validation study across eight countries. <i>Body Image</i> , 2020, 35, 320-332.	4.3	22

#	ARTICLE	IF	CITATIONS
91	Attitudes of Flemish physiotherapy students towards mental health and psychiatry. <i>Physiotherapy</i> , 2010, 96, 44-51.	0.4	21
92	Quality Assessment of Physical Activity Recommendations Within Clinical Practice Guidelines for the Prevention and Treatment of Cardio-metabolic Risk Factors in People With Schizophrenia. <i>Community Mental Health Journal</i> , 2011, 47, 703-710.	2.0	21
93	Test-retest reliability, feasibility and clinical correlates of the Eurofit test battery in people with bipolar disorder. <i>Psychiatry Research</i> , 2015, 228, 620-625.	3.3	21
94	The functional exercise capacity and its correlates in obese treatment-seeking people with binge eating disorder: an exploratory study. <i>Disability and Rehabilitation</i> , 2015, 37, 777-782.	1.8	21
95	Cardiorespiratory fitness in outpatients with bipolar disorder versus matched controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2016, 199, 1-5.	4.1	21
96	Development and testing of a model for risk and protective factors for eating disorders and higher weight among emerging adults: A study protocol. <i>Body Image</i> , 2019, 31, 139-149.	4.3	21
97	SOCIAL ANXIETY IN PHYSICAL ACTIVITY PARTICIPATION IN PATIENTS WITH MENTAL ILLNESS: A CROSS-SECTIONAL MULTICENTER STUDY. <i>Depression and Anxiety</i> , 2013, 30, 757-762.	4.1	20
98	The Psychosocial Consequences of Sports Participation for Individuals with Severe Mental Illness: A Metasynthesis Review. <i>Advances in Psychiatry</i> , 2015, 2015, 1-8.	0.4	19
99	The functional exercise capacity in patients with bipolar disorder versus healthy controls: A pilot study. <i>Psychiatry Research</i> , 2015, 229, 194-199.	3.3	19
100	Physical fitness and physical activity levels in people with alcohol use disorder versus matched healthy controls: A pilot study. <i>Alcohol</i> , 2019, 76, 73-79.	1.7	19
101	Hyperactivity in anorexia nervosa: A case study using experience sampling methodology. <i>Eating Behaviors</i> , 2004, 5, 67-74.	2.0	18
102	Sitting time, physical fitness impairments and metabolic abnormalities in people with bipolar disorder: An exploratory study. <i>Psychiatry Research</i> , 2016, 242, 7-12.	3.3	18
103	Walking capacity is associated with health related quality of life and physical activity level in patients with schizophrenia: a preliminary report. <i>Actas Espanolas De Psiquiatria</i> , 2011, 39, 211-6.	0.1	18
104	Aerobic capacity is associated with global functioning in people with schizophrenia. <i>Journal of Mental Health</i> , 2015, 24, 214-218.	1.9	17
105	Physical activity as a vital sign in patients with bipolar disorder. <i>Psychiatry Research</i> , 2016, 246, 218-222.	3.3	17
106	Reliability and validity of 6MWT for outpatients with schizophrenia: A preliminary study. <i>Psychiatry Research</i> , 2016, 237, 37-42.	3.3	17
107	Physiotherapy and Mental Health. , 2017, , .		17
108	Body dissatisfaction moderates weight curves in the inpatient treatment of Anorexia Nervosa. <i>International Journal of Eating Disorders</i> , 2010, 43, 694-700.	4.0	16

#	ARTICLE	IF	CITATIONS
109	Addressing the disparity in physical health provision for people with schizophrenia: an important role for physiotherapists. <i>Physiotherapy</i> , 2014, 100, 185-186.	0.4	16
110	Associations Between Metabolic and Aerobic Fitness Parameters in Patients With Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 23-27.	1.0	16
111	Quality of life and physical activity levels in outpatients with schizophrenia. <i>Revista Brasileira De Psiquiatria</i> , 2016, 38, 157-160.	1.7	16
112	Comparison of clinical vignettes and standardized patients as measures of physiotherapists' activity and work recommendations in patients with non-specific low back pain. <i>Clinical Rehabilitation</i> , 2016, 30, 85-94.	2.2	16
113	Top 10 research questions to promote physical activity in bipolar disorders: A consensus statement from the International Organization of Physical Therapists in Mental Health. <i>Journal of Affective Disorders</i> , 2016, 195, 82-87.	4.1	16
114	Higher cardio-respiratory fitness is associated with increased mental and physical quality of life in people with bipolar disorder: A controlled pilot study. <i>Psychiatry Research</i> , 2017, 256, 219-224.	3.3	16
115	Physical activity and sleep problems in 38 low- and middle-income countries. <i>Sleep Medicine</i> , 2018, 48, 140-147.	1.6	16
116	Cross-cultural validation of the short form of the Physical Self Inventory (PSI-S).. <i>Sport, Exercise, and Performance Psychology</i> , 2018, 7, 60-79.	0.8	16
117	Associations between expiratory spirometry parameters and limitations in daily life activities in patients with schizophrenia. <i>General Hospital Psychiatry</i> , 2014, 36, 172-176.	2.4	15
118	Concurrent validity of the international physical activity questionnaire in outpatients with bipolar disorder: Comparison with the Sensewear Armband. <i>Psychiatry Research</i> , 2016, 237, 122-126.	3.3	15
119	A cross-country examination of emotional eating, restrained eating and intuitive eating: Measurement Invariance across eight countries. <i>Body Image</i> , 2020, 35, 245-254.	4.3	15
120	The Assessment, Benefits and Delivery of Physical Activity in People with Schizophrenia: A Survey of Members of the International Organization of Physical Therapists in Mental Health. <i>Physiotherapy Research International</i> , 2014, 19, 248-256.	1.5	14
121	Behavioural Regulation in Exercise Questionnaire in people with schizophrenia: construct validity of the Portuguese versions. <i>Disability and Rehabilitation</i> , 2018, 40, 2577-2584.	1.8	14
122	Prevalence of Primary and Secondary Exercise Dependence and Its Correlation with Drive for Thinness in Practitioners of Different Sports and Physical Activities. <i>International Journal of Mental Health and Addiction</i> , 2019, 17, 89-101.	7.4	14
123	Depressive symptoms and muscular fitness contribute independently to the ability to perform daily life activities in people with bipolar disorder. <i>Nordic Journal of Psychiatry</i> , 2016, 70, 477-482.	1.3	13
124	"The Awareness-Body-Chart: A new tool assessing body awareness. <i>PLoS ONE</i> , 2017, 12, e0186597.	2.5	13
125	Psychological Distress among Patients Attending Physiotherapy: A Survey-Based Investigation of Irish Physiotherapists' Current Practice and Opinions. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2020, 72, 239-248.	0.6	13
126	Assessing affective variability in eating disorders: Affect spins less in anorexia nervosa of the restrictive type. <i>Eating Behaviors</i> , 2013, 14, 263-268.	2.0	12

#	ARTICLE	IF	CITATIONS
127	Test-retest reliability and clinical correlates of the Eurofit test battery in people with alcohol use disorders. <i>Psychiatry Research</i> , 2019, 271, 208-213.	3.3	12
128	Psychometric properties of the Polish version of the Body Attitude Test. <i>Archives of Psychiatry and Psychotherapy</i> , 2014, 16, 39-46.	0.3	12
129	Body experience assessment in non-clinical male and female subjects. <i>Eating and Weight Disorders</i> , 2009, 14, e16-e21.	2.5	11
130	Dropout from physical activity interventions in children and adolescents with attention deficit hyperactivity disorder: A systematic review and meta-analysis. <i>Mental Health and Physical Activity</i> , 2016, 11, 46-52.	1.8	11
131	Physical activity correlates in people with mild cognitive impairment: findings from six low- and middle-income countries. <i>Public Health</i> , 2018, 156, 15-25.	2.9	11
132	Adherence to physical activity recommendations and physical and mental health risk in people with severe mental illness in Uganda. <i>Psychiatry Research</i> , 2018, 260, 236-240.	3.3	11
133	Dietary Intake, Adherence to Mediterranean Diet and Lifestyle-Related Factors in People with Schizophrenia. <i>Issues in Mental Health Nursing</i> , 2019, 40, 851-860.	1.2	11
134	Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis. <i>Microbial Biotechnology</i> , 2019, 13, 562-567.	1.7	11
135	Metabolic syndrome and lung function in schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 220, 58-62.	3.3	10
136	Validity of the 6min walk test in outpatients with bipolar disorder. <i>Psychiatry Research</i> , 2015, 230, 664-667.	3.3	10
137	Lower cardiorespiratory fitness is associated with more time spent sedentary in first episode psychosis: A pilot study. <i>Psychiatry Research</i> , 2017, 253, 13-17.	3.3	10
138	Psychomotor Therapy for Patients with Severe Mental Health Disorders. , 0, , .		10
139	Correlates of physical activity among community-dwelling individuals aged 65 years or older with anxiety in six low- and middle-income countries. <i>International Psychogeriatrics</i> , 2018, 30, 705-714.	1.0	10
140	Psychometric properties of measures of sociocultural influence and internalization of appearance ideals across eight countries. <i>Body Image</i> , 2020, 35, 300-315.	4.3	10
141	Cross-Country Measurement Invariance and Effects of Sociodemographic Factors on Body Weight and Shape Concern-Related Constructs in Eight Countries. <i>Body Image</i> , 2020, 35, 288-299.	4.3	10
142	Motives for physical activity in the adoption and maintenance of physical activity in men with alcohol use disorders. <i>Psychiatry Research</i> , 2018, 261, 522-526.	3.3	9
143	A quantitative assessment of the views of mental health professionals on exercise for people with mental illness: perspectives from a low-resource setting. <i>African Health Sciences</i> , 2019, 19, 2172.	0.7	9
144	Physiotherapy for people with mental health problems in Sub-Saharan African countries: a systematic review. <i>Archives of Physiotherapy</i> , 2018, 8, 2.	1.8	8

#	ARTICLE	IF	CITATIONS
145	Measuring perfectionism, impulsivity, self-esteem and social anxiety: Cross-national study in emerging adults from eight countries. <i>Body Image</i> , 2020, 35, 265-278.	4.3	8
146	Physical activity participation, functional exercise capacity and self-esteem in patients with schizophrenia. <i>International Journal of Therapy and Rehabilitation</i> , 2011, 18, 222-229.	0.3	7
147	The Functional Exercise Capacity Is Associated With Global Functioning in People With Bipolar Disorder. <i>Journal of Nervous and Mental Disease</i> , 2016, 204, 673-677.	1.0	7
148	Sleep quality in patients with schizophrenia: The relevance of physical activity. <i>Mental Health and Physical Activity</i> , 2018, 14, 140-145.	1.8	7
149	Functional exercise capacity in inpatients with alcohol use disorder versus healthy controls: A pilot study. <i>Alcohol</i> , 2020, 82, 47-52.	1.7	7
150	Testing of a model for risk factors for eating disorders and higher weight among emerging adults: Baseline evaluation. <i>Body Image</i> , 2022, 40, 322-339.	4.3	7
151	Considering the Role of Physical Therapists Within the Treatment and Rehabilitation of Individuals With Eating Disorders: An International Survey of Expert Clinicians. <i>Physiotherapy Research International</i> , 2016, 21, 237-246.	1.5	6
152	Interest, competence, appearance, fitness and social relatedness as motives for physical activity in Ugandan outpatients with psychosis. <i>Mental Health and Physical Activity</i> , 2017, 13, 94-99.	1.8	6
153	Autonomous motivation and quality of life as predictors of physical activity in patients with schizophrenia. <i>International Journal of Psychiatry in Clinical Practice</i> , 2018, 22, 184-190.	2.4	6
154	Top 10 research questions to promote physical activity research in people with binge eating disorder. <i>Eating Disorders</i> , 2016, 24, 326-337.	3.0	5
155	Exercise self-efficacy correlates in people with psychosis. <i>Psychiatry Research</i> , 2018, 262, 359-362.	3.3	5
156	Functional exercise capacity is associated with global functioning in patients with alcohol use disorder. <i>Archives of Psychiatric Nursing</i> , 2019, 33, 144-148.	1.4	5
157	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.7	5
158	Clinical correlates of global functioning in obese treatment seeking persons with binge eating disorder. <i>Psychiatria Danubina</i> , 2014, 26, 256-60.	0.4	5
159	Body composition in girls and young women with anorexia nervosa: Comparison of different equations. <i>International Journal of Eating Disorders</i> , 2008, 41, 180-183.	4.0	4
160	Reliability of the Detailed Assessment of Speed of Handwriting on Flemish Children. <i>Pediatric Physical Therapy</i> , 2014, 26, 318-324.	0.6	4
161	Gender differences in motives for physical activity across the stages of change in Ugandan outpatients with psychosis. <i>Schizophrenia Research</i> , 2018, 197, 568-569.	2.0	4
162	Associations between alliance, physiotherapists' confidence in managing the patient and patient-reported distress in chronic low back pain practice. <i>European Journal of Physiotherapy</i> , 2021, 23, 196-200.	1.3	4

#	ARTICLE	IF	CITATIONS
163	Barriers, attitudes, confidence, and knowledge of nurses regarding metabolic health screening and intervention in people with mental illness: a pilot study from Uganda. <i>African Health Sciences</i> , 2019, 19, 2546-2554.	0.7	4
164	Missing Data in Long-term Follow-up of Patients with Eating Disorders Using the Body Attitude Test. <i>European Eating Disorders Review</i> , 2013, 21, 224-229.	4.1	3
165	Changes in eating disorder characteristics over the years. <i>European Eating Disorders Review</i> , 2018, 26, 417-421.	4.1	3
166	Cross-linguistic validity of the French and Dutch versions of the Very Short form of the Physical Self-Inventory among adolescents. <i>Body Image</i> , 2015, 15, 35-39.	4.3	2
167	Reliability and Validity of the Turkish Version of Body Attitude Test in Women with Breast Cancer. <i>Current Psychology</i> , 2022, 41, 963-969.	2.8	2
168	Clinical Practice of Body Composition Assessment in Female Subjects with Anorexia Nervosa. , 2012, , 2783-2794.		2
169	A CONTRIBUTION TO DESIGNING EFFECTIVE AND ENJOYABLE PHYSICAL ACTIVITY PROGRAMS FOR INDIVIDUALS WITH SCHIZOPHRENIA. <i>European Journal of Adapted Physical Activity</i> , 2014, 7, 24-31.	0.5	2
170	Sedentary Behavior and Quality of Life in People with Psychotic Disorders from a Low Income Country: A Study from Uganda. <i>Community Mental Health Journal</i> , 2019, 55, 714-720.	2.0	1
171	Factor structure of the German version of the pain attitudes and beliefs scale for physiotherapists. <i>Physiotherapy Theory and Practice</i> , 2019, 35, 995-1003.	1.3	1
172	Lichaamsbeleving en bewegingsdrang. , 2013, , 234-246.		1
173	The effect of psychomotor therapy on mental health in in-patient schizophrenia treatment: A randomized, double-blind intervention study. <i>Acta Gymnica</i> , 2020, 50, 83-88.	1.1	1
174	Physical Activity and Women's Mental Health. <i>Lecture Notes in Computational Vision and Biomechanics</i> , 2018, , 15-26.	0.5	0
175	Relaxatietherapie in de GGZ. , 2013, , 247-262.		0
176	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.7	0