

Michel Probst

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3060692/publications.pdf>

Version: 2024-02-01

176
papers

8,417
citations

41344

49
h-index

54911

84
g-index

179
all docs

179
docs citations

179
times ranked

7718
citing authors

#	ARTICLE	IF	CITATIONS
1	Reliability and Validity of the Turkish Version of Body Attitude Test in Women with Breast Cancer. <i>Current Psychology</i> , 2022, 41, 963-969.	2.8	2
2	Testing of a model for risk factors for eating disorders and higher weight among emerging adults: Baseline evaluation. <i>Body Image</i> , 2022, 40, 322-339.	4.3	7
3	Associations between alliance, physiotherapists' confidence in managing the patient and patient-reported distress in chronic low back pain practice. <i>European Journal of Physiotherapy</i> , 2021, 23, 196-200.	1.3	4
4	Functional exercise capacity in inpatients with alcohol use disorder versus healthy controls: A pilot study. <i>Alcohol</i> , 2020, 82, 47-52.	1.7	7
5	Assessing positive body image, body satisfaction, weight bias, and appearance comparison in emerging adults: A cross-validation study across eight countries. <i>Body Image</i> , 2020, 35, 320-332.	4.3	22
6	Psychometric properties of measures of sociocultural influence and internalization of appearance ideals across eight countries. <i>Body Image</i> , 2020, 35, 300-315.	4.3	10
7	A cross-country examination of emotional eating, restrained eating and intuitive eating: Measurement Invariance across eight countries. <i>Body Image</i> , 2020, 35, 245-254.	4.3	15
8	Measuring perfectionism, impulsivity, self-esteem and social anxiety: Cross-national study in emerging adults from eight countries. <i>Body Image</i> , 2020, 35, 265-278.	4.3	8
9	Psychological Distress among Patients Attending Physiotherapy: A Survey-Based Investigation of Irish Physiotherapists' Current Practice and Opinions. <i>Physiotherapy Canada Physiotherapie Canada</i> , 2020, 72, 239-248.	0.6	13
10	Cross-Country Measurement Invariance and Effects of Sociodemographic Factors on Body Weight and Shape Concern-Related Constructs in Eight Countries. <i>Body Image</i> , 2020, 35, 288-299.	4.3	10
11	The effect of psychomotor therapy on mental health in in-patient schizophrenia treatment: A randomized, double-blind intervention study. <i>Acta Gymnica</i> , 2020, 50, 83-88.	1.1	1
12	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.7	0
13	Physical activity participation is associated with higher quality of life scores in men with alcohol use disorders: a study from Uganda. <i>African Health Sciences</i> , 2020, 20, 1407-1415.	0.7	5
14	Physical fitness and physical activity levels in people with alcohol use disorder versus matched healthy controls: A pilot study. <i>Alcohol</i> , 2019, 76, 73-79.	1.7	19
15	Dietary Intake, Adherence to Mediterranean Diet and Lifestyle-Related Factors in People with Schizophrenia. <i>Issues in Mental Health Nursing</i> , 2019, 40, 851-860.	1.2	11
16	Development and testing of a model for risk and protective factors for eating disorders and higher weight among emerging adults: A study protocol. <i>Body Image</i> , 2019, 31, 139-149.	4.3	21
17	A quantitative assessment of the views of mental health professionals on exercise for people with mental illness: perspectives from a low-resource setting. <i>African Health Sciences</i> , 2019, 19, 2172.	0.7	9
18	Functional exercise capacity is associated with global functioning in patients with alcohol use disorder. <i>Archives of Psychiatric Nursing</i> , 2019, 33, 144-148.	1.4	5

#	ARTICLE	IF	CITATIONS
19	Test-retest reliability and clinical correlates of the Eurofit test battery in people with alcohol use disorders. <i>Psychiatry Research</i> , 2019, 271, 208-213.	3.3	12
20	Sedentary Behavior and Quality of Life in People with Psychotic Disorders from a Low Income Country: A Study from Uganda. <i>Community Mental Health Journal</i> , 2019, 55, 714-720.	2.0	1
21	Validity and correlates of the International Physical Activity Questionnaire in first-episode psychosis. <i>Microbial Biotechnology</i> , 2019, 13, 562-567.	1.7	11
22	Factor structure of the German version of the pain attitudes and beliefs scale for physiotherapists. <i>Physiotherapy Theory and Practice</i> , 2019, 35, 995-1003.	1.3	1
23	Prevalence of Primary and Secondary Exercise Dependence and Its Correlation with Drive for Thinness in Practitioners of Different Sports and Physical Activities. <i>International Journal of Mental Health and Addiction</i> , 2019, 17, 89-101.	7.4	14
24	Barriers, attitudes, confidence, and knowledge of nurses regarding metabolic health screening and intervention in people with mental illness: a pilot study from Uganda. <i>African Health Sciences</i> , 2019, 19, 2546-2554.	0.7	4
25	Sleep quality in patients with schizophrenia: The relevance of physical activity. <i>Mental Health and Physical Activity</i> , 2018, 14, 140-145.	1.8	7
26	Motives for physical activity in the adoption and maintenance of physical activity in men with alcohol use disorders. <i>Psychiatry Research</i> , 2018, 261, 522-526.	3.3	9
27	Physical activity correlates in people with mild cognitive impairment: findings from six low- and middle-income countries. <i>Public Health</i> , 2018, 156, 15-25.	2.9	11
28	Autonomous motivation and quality of life as predictors of physical activity in patients with schizophrenia. <i>International Journal of Psychiatry in Clinical Practice</i> , 2018, 22, 184-190.	2.4	6
29	Physical Therapists' Ability to Identify Psychological Factors and Their Self-Reported Competence to Manage Chronic Low Back Pain. <i>Physical Therapy</i> , 2018, 98, 471-479.	2.4	40
30	Physical Activity and Women's Mental Health. <i>Lecture Notes in Computational Vision and Biomechanics</i> , 2018, , 15-26.	0.5	0
31	Adherence to physical activity recommendations and physical and mental health risk in people with severe mental illness in Uganda. <i>Psychiatry Research</i> , 2018, 260, 236-240.	3.3	11
32	Is autonomous motivation the key to maintaining an active lifestyle in first-episode psychosis?. <i>Microbial Biotechnology</i> , 2018, 12, 821-827.	1.7	27
33	Global physical activity levels among people living with HIV: a systematic review and meta-analysis. <i>Disability and Rehabilitation</i> , 2018, 40, 388-397.	1.8	100
34	Physical activity correlates in people living with HIV/AIDS: a systematic review of 45 studies. <i>Disability and Rehabilitation</i> , 2018, 40, 1618-1629.	1.8	65
35	Behavioural Regulation in Exercise Questionnaire in people with schizophrenia: construct validity of the Portuguese versions. <i>Disability and Rehabilitation</i> , 2018, 40, 2577-2584.	1.8	14
36	Physical activity and suicidal ideation: A systematic review and meta-analysis. <i>Journal of Affective Disorders</i> , 2018, 225, 438-448.	4.1	140

#	ARTICLE	IF	CITATIONS
37	Exercise self-efficacy correlates in people with psychosis. <i>Psychiatry Research</i> , 2018, 262, 359-362.	3.3	5
38	Gender differences in motives for physical activity across the stages of change in Ugandan outpatients with psychosis. <i>Schizophrenia Research</i> , 2018, 197, 568-569.	2.0	4
39	Correlates of physical activity among community-dwelling individuals aged 65 years or older with anxiety in six low- and middle-income countries. <i>International Psychogeriatrics</i> , 2018, 30, 705-714.	1.0	10
40	Changes in eating disorder characteristics over the years. <i>European Eating Disorders Review</i> , 2018, 26, 417-421.	4.1	3
41	Physical activity and sleep problems in 38 low- and middle-income countries. <i>Sleep Medicine</i> , 2018, 48, 140-147.	1.6	16
42	Sedentary behaviour and sleep problems among 42,489 community-dwelling adults in six low- and middle-income countries. <i>Journal of Sleep Research</i> , 2018, 27, e12714.	3.2	33
43	Association of lifestyle-related factors and psychological factors on quality of life in people with schizophrenia. <i>Psychiatry Research</i> , 2018, 267, 382-393.	3.3	23
44	Physiotherapy for people with mental health problems in Sub-Saharan African countries: a systematic review. <i>Archives of Physiotherapy</i> , 2018, 8, 2.	1.8	8
45	Cross-cultural validation of the short form of the Physical Self Inventory (PSI-S).. <i>Sport, Exercise, and Performance Psychology</i> , 2018, 7, 60-79.	0.8	16
46	Cardiorespiratory Fitness in Severe Mental Illness: A Systematic Review and Meta-analysis. <i>Sports Medicine</i> , 2017, 47, 343-352.	6.5	170
47	Higher cardio-respiratory fitness is associated with increased mental and physical quality of life in people with bipolar disorder: A controlled pilot study. <i>Psychiatry Research</i> , 2017, 256, 219-224.	3.3	16
48	Interest, competence, appearance, fitness and social relatedness as motives for physical activity in Ugandan outpatients with psychosis. <i>Mental Health and Physical Activity</i> , 2017, 13, 94-99.	1.8	6
49	Sedentary Behavior in People Living With HIV: A Systematic Review and Meta-Analysis. <i>Journal of Physical Activity and Health</i> , 2017, 14, 571-577.	2.0	22
50	Lower cardiorespiratory fitness is associated with more time spent sedentary in first episode psychosis: A pilot study. <i>Psychiatry Research</i> , 2017, 253, 13-17.	3.3	10
51	Sedentary behavior and physical activity levels in people with schizophrenia, bipolar disorder and major depressive disorder: a global systematic review and meta-analysis. <i>World Psychiatry</i> , 2017, 16, 308-315.	10.4	600
52	Physical activity is associated with the physical, psychological, social and environmental quality of life in people with mental health problems in a low resource setting. <i>Psychiatry Research</i> , 2017, 258, 250-254.	3.3	25
53	The relationship between chronic physical conditions, multimorbidity and anxiety in the general population: A global perspective across 42 countries. <i>General Hospital Psychiatry</i> , 2017, 45, 1-6.	2.4	80
54	Physiotherapy and Mental Health. , 2017, , .		17

#	ARTICLE	IF	CITATIONS
55	“The Awareness-Body-Chart: A new tool assessing body awareness. PLoS ONE, 2017, 12, e0186597.	2.5	13
56	Quality of life and physical activity levels in outpatients with schizophrenia. Revista Brasileira De Psiquiatria, 2016, 38, 157-160.	1.7	16
57	Sitting time, physical fitness impairments and metabolic abnormalities in people with bipolar disorder: An exploratory study. Psychiatry Research, 2016, 242, 7-12.	3.3	18
58	Autonomous motivation is associated with the maintenance stage of behaviour change in people with affective disorders. Psychiatry Research, 2016, 240, 267-271.	3.3	33
59	Physical activity and sedentary behavior in people with bipolar disorder: A systematic review and meta-analysis. Journal of Affective Disorders, 2016, 201, 145-152.	4.1	109
60	Cardiorespiratory fitness in outpatients with bipolar disorder versus matched controls: An exploratory study. Journal of Affective Disorders, 2016, 199, 1-5.	4.1	21
61	Depressive symptoms and muscular fitness contribute independently to the ability to perform daily life activities in people with bipolar disorder. Nordic Journal of Psychiatry, 2016, 70, 477-482.	1.3	13
62	Physical activity as a vital sign in patients with bipolar disorder. Psychiatry Research, 2016, 246, 218-222.	3.3	17
63	Dropout from physical activity interventions in children and adolescents with attention deficit hyperactivity disorder: A systematic review and meta-analysis. Mental Health and Physical Activity, 2016, 11, 46-52.	1.8	11
64	Cardiorespiratory fitness levels and moderators in people with HIV: A systematic review and meta-analysis. Preventive Medicine, 2016, 93, 106-114.	3.4	36
65	Diabetes mellitus in people with schizophrenia, bipolar disorder and major depressive disorder: a systematic review and large scale meta-analysis. World Psychiatry, 2016, 15, 166-174.	10.4	487
66	The prevalence of diabetes mellitus type 2 in people with alcohol use disorders: a systematic review and large scale meta-analysis. Psychiatry Research, 2016, 246, 394-400.	3.3	27
67	The Functional Exercise Capacity Is Associated With Global Functioning in People With Bipolar Disorder. Journal of Nervous and Mental Disease, 2016, 204, 673-677.	1.0	7
68	The Prevalence of Metabolic Syndrome in Alcohol Use Disorders: A Systematic Review and Meta-analysis. Alcohol and Alcoholism, 2016, 51, 515-521.	1.6	72
69	Impact of antipsychotic medication on physical activity and physical fitness in adolescents: An exploratory study. Psychiatry Research, 2016, 242, 192-197.	3.3	23
70	What are the top 10 physical activity research questions in schizophrenia?. Disability and Rehabilitation, 2016, 38, 2235-2243.	1.8	27
71	Comparison of clinical vignettes and standardized patients as measures of physiotherapists' activity and work recommendations in patients with non-specific low back pain. Clinical Rehabilitation, 2016, 30, 85-94.	2.2	16
72	Prevalence and predictors of treatment dropout from physical activity interventions in schizophrenia: a meta-analysis. General Hospital Psychiatry, 2016, 39, 15-23.	2.4	172

#	ARTICLE	IF	CITATIONS
73	Physical activity as a vital sign in patients with schizophrenia: Evidence and clinical recommendations. <i>Schizophrenia Research</i> , 2016, 170, 336-340.	2.0	32
74	Top 10 research questions to promote physical activity in bipolar disorders: A consensus statement from the International Organization of Physical Therapists in Mental Health. <i>Journal of Affective Disorders</i> , 2016, 195, 82-87.	4.1	16
75	Concurrent validity of the international physical activity questionnaire in outpatients with bipolar disorder: Comparison with the Sensewear Armband. <i>Psychiatry Research</i> , 2016, 237, 122-126.	3.3	15
76	Reliability and validity of 6MWT for outpatients with schizophrenia: A preliminary study. <i>Psychiatry Research</i> , 2016, 237, 37-42.	3.3	17
77	A comparison of physical fitness in patients with bipolar disorder, schizophrenia and healthy controls. <i>Disability and Rehabilitation</i> , 2016, 38, 2047-2051.	1.8	27
78	Top 10 research questions to promote physical activity research in people with binge eating disorder. <i>Eating Disorders</i> , 2016, 24, 326-337.	3.0	5
79	Considering the Role of Physical Therapists Within the Treatment and Rehabilitation of Individuals With Eating Disorders: An International Survey of Expert Clinicians. <i>Physiotherapy Research International</i> , 2016, 21, 237-246.	1.5	6
80	Validity of the 6min walk test in outpatients with bipolar disorder. <i>Psychiatry Research</i> , 2015, 230, 664-667.	3.3	10
81	TYPE 2 DIABETES IN PATIENTS WITH MAJOR DEPRESSIVE DISORDER: A META-ANALYSIS OF PREVALENCE ESTIMATES AND PREDICTORS. <i>Depression and Anxiety</i> , 2015, 32, 763-773.	4.1	138
82	Health-related quality of life and aerobic fitness in people with schizophrenia. <i>International Journal of Mental Health Nursing</i> , 2015, 24, 394-402.	3.8	35
83	The Psychosocial Consequences of Sports Participation for Individuals with Severe Mental Illness: A Metasynthesis Review. <i>Advances in Psychiatry</i> , 2015, 2015, 1-8.	0.4	19
84	A Systematic Review of Physical Activity Correlates in Alcohol Use Disorders. <i>Archives of Psychiatric Nursing</i> , 2015, 29, 196-201.	1.4	45
85	Cross-linguistic validity of the French and Dutch versions of the Very Short form of the Physical Self-Inventory among adolescents. <i>Body Image</i> , 2015, 15, 35-39.	4.3	2
86	Could autonomous motivation hold the key to successfully implementing lifestyle changes in affective disorders? A multicentre cross sectional study. <i>Psychiatry Research</i> , 2015, 228, 100-106.	3.3	47
87	Test-retest reliability, feasibility and clinical correlates of the Eurofit test battery in people with bipolar disorder. <i>Psychiatry Research</i> , 2015, 228, 620-625.	3.3	21
88	The functional exercise capacity in patients with bipolar disorder versus healthy controls: A pilot study. <i>Psychiatry Research</i> , 2015, 229, 194-199.	3.3	19
89	Health-related physical fitness in patients with bipolar disorder vs. healthy controls: An exploratory study. <i>Journal of Affective Disorders</i> , 2015, 177, 22-27.	4.1	25
90	Associations Between Metabolic and Aerobic Fitness Parameters in Patients With Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2015, 203, 23-27.	1.0	16

#	ARTICLE	IF	CITATIONS
91	Promotion of cardiorespiratory fitness in schizophrenia: a clinical overview and meta-analysis. <i>Acta Psychiatrica Scandinavica</i> , 2015, 132, 131-143.	4.5	108
92	Aerobic capacity is associated with global functioning in people with schizophrenia. <i>Journal of Mental Health</i> , 2015, 24, 214-218.	1.9	17
93	Adopting and maintaining physical activity behaviours in people with severe mental illness: The importance of autonomous motivation. <i>Preventive Medicine</i> , 2015, 81, 216-220.	3.4	86
94	Negative symptoms are associated with lower autonomous motivation towards physical activity in people with schizophrenia. <i>Comprehensive Psychiatry</i> , 2015, 56, 128-132.	3.1	77
95	The functional exercise capacity and its correlates in obese treatment-seeking people with binge eating disorder: an exploratory study. <i>Disability and Rehabilitation</i> , 2015, 37, 777-782.	1.8	21
96	Prevalence and Predictors of Type 2 Diabetes Mellitus in People With Bipolar Disorder. <i>Journal of Clinical Psychiatry</i> , 2015, 76, 1490-1499.	2.2	85
97	Investigating the benefits of sport participation for individuals with schizophrenia: a systematic review. <i>Psychiatria Danubina</i> , 2015, 27, 2-13.	0.4	41
98	What are the factors that influence physical activity participation in individuals with depression? A review of physical activity correlates from 59 studies. <i>Psychiatria Danubina</i> , 2015, 27, 210-24.	0.4	51
99	The Assessment, Benefits and Delivery of Physical Activity in People with Schizophrenia: A Survey of Members of the International Organization of Physical Therapists in Mental Health. <i>Physiotherapy Research International</i> , 2014, 19, 248-256.	1.5	14
100	Effects of a group physical activity program on physical fitness and quality of life in individuals with schizophrenia. <i>Mental Health and Physical Activity</i> , 2014, 7, 155-162.	1.8	29
101	Metabolic syndrome and metabolic abnormalities in patients with major depressive disorder: a meta-analysis of prevalences and moderating variables. <i>Psychological Medicine</i> , 2014, 44, 2017-2028.	4.5	223
102	The value of social support to encourage people with schizophrenia to engage in physical activity: an international insight from specialist mental health physiotherapists. <i>Journal of Mental Health</i> , 2014, 23, 256-260.	1.9	26
103	Physical Activity Correlates in Persons with Binge Eating Disorder: A Systematic Review. <i>European Eating Disorders Review</i> , 2014, 22, 1-8.	4.1	37
104	Reliability and clinical correlates of the Astrand-Rhyming sub-maximal exercise test in patients with schizophrenia or schizoaffective disorder. <i>Psychiatry Research</i> , 2014, 220, 778-783.	3.3	26
105	Self-determination and stage of readiness to change physical activity behaviour in schizophrenia. <i>Mental Health and Physical Activity</i> , 2014, 7, 171-176.	1.8	41
106	Reliability of the Detailed Assessment of Speed of Handwriting on Flemish Children. <i>Pediatric Physical Therapy</i> , 2014, 26, 318-324.	0.6	4
107	Understanding the role of physiotherapists in schizophrenia: an international perspective from members of the International Organisation of Physical Therapists in Mental Health (IOPTMH). <i>Journal of Mental Health</i> , 2014, 23, 125-129.	1.9	34
108	Barriers to and Facilitators of Physical Activity Among Persons With Schizophrenia: A Survey of Physical Therapists. <i>Psychiatric Services</i> , 2014, 65, 693-696.	2.0	88

#	ARTICLE	IF	CITATIONS
109	Physiotherapists can help implement physical activity programmes in clinical practice. <i>British Journal of Psychiatry</i> , 2014, 204, 164-164.	2.8	52
110	A systematic review of physical therapy interventions for patients with anorexia and bulimia nervosa. <i>Disability and Rehabilitation</i> , 2014, 36, 628-634.	1.8	80
111	Neurobiological effects of physical exercise in schizophrenia: a systematic review. <i>Disability and Rehabilitation</i> , 2014, 36, 1749-1754.	1.8	63
112	Markers of inflammation in schizophrenia: association vs. causation. <i>World Psychiatry</i> , 2014, 13, 189-192.	10.4	54
113	What are the factors associated with physical activity (PA) participation in community dwelling adults with dementia? A systematic review of PA correlates. <i>Archives of Gerontology and Geriatrics</i> , 2014, 59, 195-203.	3.0	67
114	The transcending benefits of physical activity for individuals with schizophrenia: A systematic review and meta-ethnography. <i>Psychiatry Research</i> , 2014, 220, 11-19.	3.3	75
115	Metabolic syndrome and lung function in schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 220, 58-62.	3.3	10
116	Health related quality of life, physical fitness and physical activity participation in treatment-seeking obese persons with and without binge eating disorder. <i>Psychiatry Research</i> , 2014, 216, 97-102.	3.3	37
117	Associations between perceived neighbourhood environmental attributes and self-reported sitting time in patients with schizophrenia: A pilot study. <i>Psychiatry Research</i> , 2014, 215, 33-38.	3.3	25
118	Associations between expiratory spirometry parameters and limitations in daily life activities in patients with schizophrenia. <i>General Hospital Psychiatry</i> , 2014, 36, 172-176.	2.4	15
119	Addressing the disparity in physical health provision for people with schizophrenia: an important role for physiotherapists. <i>Physiotherapy</i> , 2014, 100, 185-186.	0.4	16
120	Changes in physical activity, physical fitness, self-perception and quality of life following a 6-month physical activity counseling and cognitive behavioral therapy program in outpatients with binge eating disorder. <i>Psychiatry Research</i> , 2014, 219, 361-366.	3.3	34
121	The benefits of walking for individuals with schizophrenia spectrum disorders: A systematic review. <i>International Journal of Therapy and Rehabilitation</i> , 2014, 21, 410-420.	0.3	29
122	Psychometric properties of the Polish version of the Body Attitude Test. <i>Archives of Psychiatry and Psychotherapy</i> , 2014, 16, 39-46.	0.3	12
123	A CONTRIBUTION TO DESIGNING EFFECTIVE AND ENJOYABLE PHYSICAL ACTIVITY PROGRAMS FOR INDIVIDUALS WITH SCHIZOPHRENIA. <i>European Journal of Adapted Physical Activity</i> , 2014, 7, 24-31.	0.5	2
124	Clinical correlates of global functioning in obese treatment seeking persons with binge eating disorder. <i>Psychiatria Danubina</i> , 2014, 26, 256-60.	0.4	5
125	An impaired health related muscular fitness contributes to a reduced walking capacity in patients with schizophrenia: a cross-sectional study. <i>BMC Psychiatry</i> , 2013, 13, 5.	2.6	28
126	SOCIAL ANXIETY IN PHYSICAL ACTIVITY PARTICIPATION IN PATIENTS WITH MENTAL ILLNESS: A CROSS-SECTIONAL MULTICENTER STUDY. <i>Depression and Anxiety</i> , 2013, 30, 757-762.	4.1	20

#	ARTICLE	IF	CITATIONS
127	The importance of self-determined motivation towards physical activity in patients with schizophrenia. <i>Psychiatry Research</i> , 2013, 210, 812-818.	3.3	133
128	Neurocognition in clinical high risk young adults who did or did not convert to a first schizophrenic psychosis: A meta-analysis. <i>Schizophrenia Research</i> , 2013, 149, 48-55.	2.0	97
129	A systematic review on physical therapy interventions for patients with binge eating disorder. <i>Disability and Rehabilitation</i> , 2013, 35, 2191-2196.	1.8	55
130	Progressive muscle relaxation in persons with schizophrenia: a systematic review of randomized controlled trials. <i>Clinical Rehabilitation</i> , 2013, 27, 291-298.	2.2	47
131	A review of physical activity correlates in patients with bipolar disorder. <i>Journal of Affective Disorders</i> , 2013, 145, 285-291.	4.1	108
132	Assessing affective variability in eating disorders: Affect spins less in anorexia nervosa of the restrictive type. <i>Eating Behaviors</i> , 2013, 14, 263-268.	2.0	12
133	Relationships between physical fitness, physical activity, smoking and metabolic and mental health parameters in people with schizophrenia. <i>Psychiatry Research</i> , 2013, 207, 25-32.	3.3	131
134	Can cognitive behavioural therapy based strategies be integrated into physiotherapy for the prevention of chronic low back pain? A systematic review. <i>Disability and Rehabilitation</i> , 2013, 35, 1-10.	1.8	80
135	Associations between physical activity and the built environment in patients with schizophrenia: a multi-centre study. <i>General Hospital Psychiatry</i> , 2013, 35, 653-658.	2.4	36
136	A meta-analysis of cardio-metabolic abnormalities in drug naïve, first-episode and multi-episode patients with schizophrenia versus general population controls. <i>World Psychiatry</i> , 2013, 12, 240-250.	10.4	220
137	Missing Data in Long-term Follow-up of Patients with Eating Disorders Using the Body Attitude Test. <i>European Eating Disorders Review</i> , 2013, 21, 224-229.	4.1	3
138	Metabolic Syndrome and Metabolic Abnormalities in Bipolar Disorder: A Meta-Analysis of Prevalence Rates and Moderators. <i>American Journal of Psychiatry</i> , 2013, 170, 265-274.	7.2	336
139	Variability in Affective Activation Predicts Non-suicidal Self-injury in Eating Disorders. <i>European Eating Disorders Review</i> , 2013, 21, 143-147.	4.1	31
140	Diabetes, physical activity participation and exercise capacity in patients with schizophrenia. <i>Psychiatry and Clinical Neurosciences</i> , 2013, 67, 451-456.	1.8	28
141	Physical activity and sedentary behaviour in outpatients with schizophrenia: A systematic review and meta-analysis. <i>International Journal of Therapy and Rehabilitation</i> , 2013, 20, 588-595.	0.3	69
142	Lichaamsbeleving en bewegingsdrang. , 2013, , 234-246.		1
143	Relaxatietherapie in de GGZ. , 2013, , 247-262.		0
144	Systematic Review of the Benefits of Physical Therapy Within a Multidisciplinary Care Approach for People With Schizophrenia. <i>Physical Therapy</i> , 2012, 92, 11-23.	2.4	114

#	ARTICLE	IF	CITATIONS
145	A Cognitive- Behavioral Therapeutic Program for Patients With Obesity and Binge Eating Disorder. Behavior Modification, 2012, 36, 670-686.	1.6	27
146	International Organization of Physical Therapy in Mental Health consensus on physical activity within multidisciplinary rehabilitation programmes for minimising cardio-metabolic risk in patients with schizophrenia. Disability and Rehabilitation, 2012, 34, 1-12.	1.8	122
147	Eurofit test battery in patients with schizophrenia or schizoaffective disorder: Reliability and clinical correlates. European Psychiatry, 2012, 27, 416-421.	0.2	29
148	Associations between sedentary behaviour and metabolic parameters in patients with schizophrenia. Psychiatry Research, 2012, 200, 73-78.	3.3	120
149	The International Organization of Physical Therapists working in Mental Health (IOPTMH). Mental Health and Physical Activity, 2012, 5, 20-21.	1.8	25
150	A systematic review of correlates of physical activity in patients with schizophrenia. Acta Psychiatrica Scandinavica, 2012, 125, 352-362.	4.5	255
151	The functional exercise capacity is correlated with global functioning in patients with schizophrenia. Acta Psychiatrica Scandinavica, 2012, 125, 382-387.	4.5	74
152	Yoga in schizophrenia: a systematic review of randomised controlled trials. Acta Psychiatrica Scandinavica, 2012, 126, 12-20.	4.5	94
153	Clinical Practice of Body Composition Assessment in Female Subjects with Anorexia Nervosa. , 2012, , 2783-2794.		2
154	State anxiety, psychological stress and positive well-being responses to yoga and aerobic exercise in people with schizophrenia: a pilot study. Disability and Rehabilitation, 2011, 33, 684-689.	1.8	91
155	Lack of physical activity during leisure time contributes to an impaired health related quality of life in patients with schizophrenia. Schizophrenia Research, 2011, 129, 122-127.	2.0	91
156	Association of the metabolic syndrome with physical activity performance in patients with schizophrenia. Diabetes and Metabolism, 2011, 37, 318-323.	2.9	52
157	Reliability, minimal detectable changes, practice effects and correlates of the 6-min walk test in patients with schizophrenia. Psychiatry Research, 2011, 187, 62-67.	3.3	53
158	Effects of progressive muscle relaxation on state anxiety and subjective well-being in people with schizophrenia: a randomized controlled trial. Clinical Rehabilitation, 2011, 25, 567-575.	2.2	50
159	Physical activity participation, functional exercise capacity and self-esteem in patients with schizophrenia. International Journal of Therapy and Rehabilitation, 2011, 18, 222-229.	0.3	7
160	Relationships between obesity, functional exercise capacity, physical activity participation and physical self-perception in people with schizophrenia. Acta Psychiatrica Scandinavica, 2011, 123, 423-430.	4.5	113
161	Quality Assessment of Physical Activity Recommendations Within Clinical Practice Guidelines for the Prevention and Treatment of Cardio-metabolic Risk Factors in People With Schizophrenia. Community Mental Health Journal, 2011, 47, 703-710.	2.0	21
162	Walking capacity is associated with health related quality of life and physical activity level in patients with schizophrenia: a preliminary report. Actas Espanolas De Psiquiatria, 2011, 39, 211-6.	0.1	18

#	ARTICLE	IF	CITATIONS
163	Attitudes of Flemish physiotherapy students towards mental health and psychiatry. <i>Physiotherapy</i> , 2010, 96, 44-51.	0.4	21
164	Body dissatisfaction moderates weight curves in the inpatient treatment of Anorexia Nervosa. <i>International Journal of Eating Disorders</i> , 2010, 43, 694-700.	4.0	16
165	Considering a frame of reference for physical activity research related to the cardiometabolic risk profile in schizophrenia. <i>Psychiatry Research</i> , 2010, 177, 271-279.	3.3	125
166	Psychomotor Therapy and Psychiatry: Whatâ€™s in a Name?. <i>The Open Complementary Medicine Journal</i> , 2010, 2, 105-113.	1.5	57
167	Cardiometabolic effects of physical activity interventions for people with schizophrenia. <i>Physical Therapy Reviews</i> , 2009, 14, 388-398.	0.8	48
168	Body experience assessment in non-clinical male and female subjects. <i>Eating and Weight Disorders</i> , 2009, 14, e16-e21.	2.5	11
169	Body composition in girls and young women with anorexia nervosa: Comparison of different equations. <i>International Journal of Eating Disorders</i> , 2008, 41, 180-183.	4.0	4
170	Evaluation of body experience questionnaires in eating disorders in female patients (AN/BN) and nonclinical participants. <i>International Journal of Eating Disorders</i> , 2008, 41, 657-665.	4.0	46
171	Drive for thinness, affect regulation and physical activity in eating disorders: A daily life study. <i>Behaviour Research and Therapy</i> , 2007, 45, 1717-1734.	3.1	68
172	Hyperactivity in anorexia nervosa: A case study using experience sampling methodology. <i>Eating Behaviors</i> , 2004, 5, 67-74.	2.0	18
173	Body composition of anorexia nervosa patients assessed by underwater weighing and skinfold-thickness measurements before and after weight gain. <i>American Journal of Clinical Nutrition</i> , 2001, 73, 190-197.	4.7	52
174	The significance of body size estimation in eating disorders: Its relationship with clinical and psychological variables. , 1998, 24, 167-174.		42
175	The Body Attitude Test for Patients with an Eating Disorder: Psychometric Characteristics of a New Questionnaire. <i>Eating Disorders</i> , 1995, 3, 133-144.	3.0	200
176	Psychomotor Therapy for Patients with Severe Mental Health Disorders. , 0, , .		10