

Susan B Altenbach

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3080268/publications.pdf>

Version: 2024-02-01

48
papers

2,865
citations

147801

31
h-index

223800

46
g-index

48
all docs

48
docs citations

48
times ranked

2198
citing authors

#	ARTICLE	IF	CITATIONS
1	Proteomic Determination of Low-Molecular-Weight Glutenin Subunit Composition in Aroona Near-Isogenic Lines and Standard Wheat Cultivars. <i>International Journal of Molecular Sciences</i> , 2021, 22, 7709.	4.1	4
2	Exploiting the reference genome sequence of hexaploid wheat: a proteomic study of flour proteins from the cultivar Chinese Spring. <i>Functional and Integrative Genomics</i> , 2020, 20, 1-16.	3.5	42
3	Development of an Optimized MALDI-TOF-MS Method for High-Throughput Identification of High-Molecular-Weight Glutenin Subunits in Wheat. <i>Molecules</i> , 2020, 25, 4347.	3.8	10
4	Deciphering the immunogenic potential of wheat flour: a reference map of the salt-soluble proteome from the U.S. wheat Butte 86. <i>Proteome Science</i> , 2020, 18, 8.	1.7	6
5	Comparison of MALDI-TOF-MS and RP-HPLC as Rapid Screening Methods for Wheat Lines With Altered Gliadin Compositions. <i>Frontiers in Plant Science</i> , 2020, 11, 600489.	3.6	6
6	Reducing the Immunogenic Potential of Wheat Flour: Silencing of Alpha Gliadin Genes in a U.S. Wheat Cultivar. <i>Frontiers in Plant Science</i> , 2020, 11, 20.	3.6	25
7	Rapid evolution of α -gliadin gene family revealed by analyzing Gli-2 locus regions of wild emmer wheat. <i>Functional and Integrative Genomics</i> , 2019, 19, 993-1005.	3.5	28
8	Elimination of Omega-1,2 Gliadins From Bread Wheat (<i>Triticum aestivum</i>) Flour: Effects on Immunogenic Potential and End-Use Quality. <i>Frontiers in Plant Science</i> , 2019, 10, 580.	3.6	39
9	Dynamic Evolution of α -Gliadin Prolamin Gene Family in Homeologous Genomes of Hexaploid Wheat. <i>Scientific Reports</i> , 2018, 8, 5181.	3.3	68
10	Towards reducing the immunogenic potential of wheat flour: omega gliadins encoded by the D genome of hexaploid wheat may also harbor epitopes for the serious food allergy WDEIA. <i>BMC Plant Biology</i> , 2018, 18, 291.	3.6	23
11	LED Lighting "Modification of Growth, Metabolism, Yield and Flour Composition in Wheat by Spectral Quality and Intensity. <i>Frontiers in Plant Science</i> , 2018, 9, 605.	3.6	73
12	Gene Duplication and Evolution Dynamics in the Homeologous Regions Harboring Multiple Prolamin and Resistance Gene Families in Hexaploid Wheat. <i>Frontiers in Plant Science</i> , 2018, 9, 673.	3.6	84
13	Proteomic Profiling and Epitope Analysis of the Complex α -, β -, and γ -Gliadin Families in a Commercial Bread Wheat. <i>Frontiers in Plant Science</i> , 2018, 9, 818.	3.6	15
14	New insights into structural organization and gene duplication in a 1.75 Mb genomic region harboring the α -gliadin gene family in <i>Aegilops tauschii</i> , the source of wheat D genome. <i>Plant Journal</i> , 2017, 92, 571-583.	5.7	29
15	Improved Method for Reliable HMW-GS Identification by RP-HPLC and SDS-PAGE in Common Wheat Cultivars. <i>Molecules</i> , 2017, 22, 1055.	3.8	22
16	Proteomics of Wheat Flour., 2017, , 57-73.		3
17	Effects of post-anthesis fertilizer on the protein composition of the gluten polymer in a US bread wheat. <i>Journal of Cereal Science</i> , 2016, 68, 66-73.	3.7	8
18	Comprehensive identification of LMW-GS genes and their protein products in a common wheat variety. <i>Functional and Integrative Genomics</i> , 2016, 16, 269-279.	3.5	27

#	ARTICLE	IF	CITATIONS
19	RNA interference targeting rye secalins alters flour protein composition in a wheat variety carrying a 1BL.1RS translocation. <i>Journal of Cereal Science</i> , 2016, 68, 172-180.	3.7	9
20	Assessment of the Allergenic Potential of Transgenic Wheat (<i>Triticum aestivum</i>) with Reduced Levels of γ -Gliadins, the Major Sensitizing Allergen in Wheat-Dependent Exercise-Induced Anaphylaxis. <i>Journal of Agricultural and Food Chemistry</i> , 2015, 63, 9323-9332.	5.2	36
21	Specific Nongluten Proteins of Wheat Are Novel Target Antigens in Celiac Disease Humoral Response. <i>Journal of Proteome Research</i> , 2015, 14, 503-511.	3.7	60
22	Silencing of omega-5 gliadins in transgenic wheat eliminates a major source of environmental variability and improves dough mixing properties of flour. <i>BMC Plant Biology</i> , 2014, 14, 393.	3.6	56
23	Quantitative proteomic analysis of wheat grain proteins reveals differential effects of silencing of omega-5 gliadin genes in transgenic lines. <i>Journal of Cereal Science</i> , 2014, 59, 118-125.	3.7	35
24	Protein composition of wheat gluten polymer fractions determined by quantitative two-dimensional gel electrophoresis and tandem mass spectrometry. <i>Proteome Science</i> , 2014, 12, 8.	1.7	68
25	Comparative proteomic analysis of the effect of temperature and fertilizer on gliadin and glutenin accumulation in the developing endosperm and flour from <i>Triticum aestivum</i> L. cv. Butte 86. <i>Proteome Science</i> , 2013, 11, 8.	1.7	83
26	Farinin: Characterization of a Novel Wheat Endosperm Protein Belonging to the Prolamin Superfamily. <i>Journal of Agricultural and Food Chemistry</i> , 2013, 61, 2407-2417.	5.2	37
27	New insights into the effects of high temperature, drought and post-anthesis fertilizer on wheat grain development. <i>Journal of Cereal Science</i> , 2012, 56, 39-50.	3.7	103
28	Effect of cleavage enzyme, search algorithm and decoy database on mass spectrometric identification of wheat gluten proteins. <i>Phytochemistry</i> , 2011, 72, 1154-1161.	2.9	37
29	Deciphering the complexities of the wheat flour proteome using quantitative two-dimensional electrophoresis, three proteases and tandem mass spectrometry. <i>Proteome Science</i> , 2011, 9, 10.	1.7	199
30	Differential effects of a post-anthesis fertilizer regimen on the wheat flour proteome determined by quantitative 2-DE. <i>Proteome Science</i> , 2011, 9, 46.	1.7	61
31	The spectrum of low molecular weight alpha-amylase/protease inhibitor genes expressed in the US bread wheat cultivar Butte 86. <i>BMC Research Notes</i> , 2011, 4, 242.	1.4	82
32	Transformation of the US bread wheat "Butte 86"™ and silencing of omega-5 gliadin genes. <i>GM Crops</i> , 2011, 2, 66-73.	1.9	62
33	Integration of transcriptomic and proteomic data from a single wheat cultivar provides new tools for understanding the roles of individual alpha gliadin proteins in flour quality and celiac disease. <i>Journal of Cereal Science</i> , 2010, 52, 143-151.	3.7	39
34	Analysis of expressed sequence tags from a single wheat cultivar facilitates interpretation of tandem mass spectrometry data and discrimination of gamma gliadin proteins that may play different functional roles in flour. <i>BMC Plant Biology</i> , 2010, 10, 7.	3.6	45
35	Effect of high temperature on albumin and globulin accumulation in the endosperm proteome of the developing wheat grain. <i>Journal of Cereal Science</i> , 2009, 49, 12-23.	3.7	140
36	Expression of globulin-2, a member of the cupin superfamily of proteins with similarity to known food allergens, is increased under high temperature regimens during wheat grain development. <i>Journal of Cereal Science</i> , 2009, 49, 47-54.	3.7	24

#	ARTICLE	IF	CITATIONS
37	Genes encoding the PR-4 protein wheatwin are developmentally regulated in wheat grains and respond to high temperatures during grainfill. <i>Plant Science</i> , 2007, 173, 135-143.	3.6	26
38	Omega gliadin genes expressed in <i>Triticum aestivum</i> cv. Butte 86: Effects of post-anthesis fertilizer on transcript accumulation during grain development. <i>Journal of Cereal Science</i> , 2007, 46, 169-177.	3.7	36
39	Protein accumulation and composition in wheat grains: Effects of mineral nutrients and high temperature. <i>European Journal of Agronomy</i> , 2006, 25, 96-107.	4.1	201
40	Transcript profiles of genes expressed in endosperm tissue are altered by high temperature during wheat grain development. <i>Journal of Cereal Science</i> , 2004, 40, 115-126.	3.7	44
41	Effect of temperature on expression of genes encoding enzymes for starch biosynthesis in developing wheat endosperm. <i>Plant Science</i> , 2003, 164, 873-881.	3.6	276
42	Environmental Conditions During Wheat Grain Development Alter Temporal Regulation of Major Gluten Protein Genes. <i>Cereal Chemistry</i> , 2002, 79, 279-285.	2.2	59
43	Accumulation of a Brazil nut albumin in seeds of transgenic canola results in enhanced levels of seed protein methionine. <i>Plant Molecular Biology</i> , 1992, 18, 235-245.	3.9	186
44	Manipulation of methionine-rich protein genes in plant seeds. <i>Trends in Biotechnology</i> , 1990, 8, 156-160.	9.3	35
45	Enhancement of the methionine content of seed proteins by the expression of a chimeric gene encoding a methionine-rich protein in transgenic plants. <i>Plant Molecular Biology</i> , 1989, 13, 513-522.	3.9	148
46	Cloning and sequence analysis of a cDNA encoding a Brazil nut protein exceptionally rich in methionine. <i>Plant Molecular Biology</i> , 1987, 8, 239-250.	3.9	108
47	Properties, biosynthesis and processing of a sulfur-rich protein in Brazil nut (<i>Bertholletia excelsa</i>) Tj ETQq1 1 0.784314 rgBT /Overlock 10 0.2 56		
48	Endosperm and Amyloplast Proteomes of Wheat Grain. , 0, , 207-222.		2