René Gislum

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3081078/publications.pdf

Version: 2024-02-01

52 papers 1,200 citations

331538 21 h-index 32 g-index

54 all docs

54 docs citations

54 times ranked

1495 citing authors

#	Article	IF	CITATIONS
1	Designing and Testing a UAV Mapping System for Agricultural Field Surveying. Sensors, 2017, 17, 2703.	2.1	132
2	Methane production potential from Miscanthus sp.: Effect of harvesting time, genotypes and plant fractions. Biosystems Engineering, 2015, 133, 71-80.	1.9	65
3	Productivity and carbon footprint of perennial grass–forage legume intercropping strategies with high or low nitrogen fertilizer input. Science of the Total Environment, 2016, 541, 1339-1347.	3.9	64
4	Quantification of fructan concentration in grasses using NIR spectroscopy and PLSR. Field Crops Research, 2011, 120, 31-37.	2.3	62
5	Single seed near-infrared hyperspectral imaging in determining tomato (Solanum lycopersicum L.) seed quality in association with multivariate data analysis. Sensors and Actuators B: Chemical, 2016, 237, 1027-1034.	4.0	51
6	Use of Multispectral Imaging in Varietal Identification of Tomato. Sensors, 2015, 15, 4496-4512.	2.1	45
7	Predicting heavy metal contents by applying machine learning approaches and environmental covariates in west of Iran. Journal of Geochemical Exploration, 2022, 233, 106921.	1.5	45
8	Viability Prediction of Ricinus cummunis L. Seeds Using Multispectral Imaging. Sensors, 2015, 15, 4592-4604.	2.1	43
9	Comparing predictive ability of laser-induced breakdown spectroscopy to visible near-infrared spectroscopy for soil property determination. Biosystems Engineering, 2017, 156, 157-172.	1.9	43
10	Segmentation of lettuce in coloured 3D point clouds for fresh weight estimation. Computers and Electronics in Agriculture, 2018, 154, 373-381.	3.7	43
11	Classification of different tomato seed cultivars by multispectral visible-near infrared spectroscopy and chemometrics. Journal of Spectral Imaging, 0, , .	0.0	33
12	Optimal Sample Size for Predicting Viability of Cabbage and Radish Seeds Based on near Infrared Spectra of Single Seeds. Journal of Near Infrared Spectroscopy, 2011, 19, 451-461.	0.8	32
13	Predicting Soil Organic Carbon at Field Scale Using a National Soil Spectral Library. Journal of Near Infrared Spectroscopy, 2013, 21, 213-222.	0.8	32
14	Quantification of SOC and Clay Content Using Visible Near-Infrared Reflectance–Mid-Infrared Reflectance Spectroscopy With Jack-Knifing Partial Least Squares Regression. Soil Science, 2014, 179, 325-332.	0.9	32
15	Estimation of the Botanical Composition of Clover-Grass Leys from RGB Images Using Data Simulation and Fully Convolutional Neural Networks. Sensors, 2017, 17, 2930.	2.1	31
16	Visible–Nearâ€Infrared Spectroscopy Can Predict the Clay/Organic Carbon and Mineral Fines/Organic Carbon Ratios. Soil Science Society of America Journal, 2016, 80, 1486-1495.	1.2	29
17	Separation of viable and non-viable tomato (Solanum lycopersicum L.) seeds using single seed near-infrared spectroscopy. Computers and Electronics in Agriculture, 2017, 142, 348-355.	3.7	26
18	Development of NIR calibration models to assess year-to-year variation in total non-structural carbohydrates in grasses using PLSR. Chemometrics and Intelligent Laboratory Systems, 2012, 111, 34-38.	1.8	25

#	Article	IF	CITATIONS
19	Estimating the nitrogen nutrition index in grass seed crops using a UAV-mounted multispectral camera. International Journal of Remote Sensing, 2019, 40, 2467-2482.	1.3	25
20	The Use of Image-Spectroscopy Technology as a Diagnostic Method for Seed Health Testing and Variety Identification. PLoS ONE, 2016, 11, e0152011.	1.1	25
21	Validity of accessible critical nitrogen dilution curves in perennial ryegrass for seed production. Field Crops Research, 2009, 111, 152-156.	2.3	22
22	Use of partial least squares discriminant analysis on visibleâ€near infrared multispectral image data to examine germination ability and germ length in spinach seeds. Journal of Chemometrics, 2012, 26, 462-466.	0.7	22
23	Classification of Viable and Non-Viable Spinach (<i>Spinacia Oleracea</i> L.) Seeds by Single Seed near Infrared Spectroscopy and Extended Canonical Variates Analysis. Journal of Near Infrared Spectroscopy, 2011, 19, 171-180.	0.8	21
24	Prediction of biogas yield and its kinetics in reed canary grass using near infrared reflectance spectroscopy and chemometrics. Bioresource Technology, 2013, 146, 282-287.	4.8	21
25	Correlation of Deoxynivalenol Accumulation in Fusarium-Infected Winter and Spring Wheat Cultivars with Secondary Metabolites at Different Growth Stages. Journal of Agricultural and Food Chemistry, 2016, 64, 4545-4555.	2.4	21
26	The effect of harvest time, dry matter content and mechanical pretreatments on anaerobic digestion and enzymatic hydrolysis of miscanthus. Bioresource Technology, 2016, 218, 1008-1015.	4.8	17
27	Grain protein concentration and harvestable protein under future climate conditions. A study of 108 spring barley accessions. Journal of Experimental Botany, 2016, 67, 2151-2158.	2.4	16
28	Seed yield response to N fertilization and potential of proximal sensing in Italian ryegrass seed crops. Field Crops Research, 2017, 211, 37-47.	2.3	16
29	The Use of Multispectral Imaging and Single Seed and Bulk Near-Infrared Spectroscopy to Characterize Seed Covering Structures: Methods and Applications in Seed Testing and Research. Agriculture (Switzerland), 2021, 11, 301.	1.4	16
30	Path and Ridge Regression Analysis of Seed Yield and Seed Yield Components of Russian Wildrye (Psathyrostachys juncea Nevski) under Field Conditions. PLoS ONE, 2011, 6, e18245.	1.1	15
31	Effects of winter wheat N status on assimilate and N partitioning in the mechanistic agroecosystem model DAISY. Journal of Agronomy and Crop Science, 2020, 206, 784-805.	1.7	12
32	Changes in spatiotemporal protein and amino acid gradients in wheat caryopsis after N-topdressing. Plant Science, 2020, 291, 110336.	1.7	11
33	Are ant feces nutrients for plants? A metabolomics approach to elucidate the nutritional effects on plants hosting weaver ants. Metabolomics, 2015, 11, 1013-1028.	1.4	10
34	Characterisation of Castor (Ricinus communis L.) Seed Quality Using Fourier Transform Near-Infrared Spectroscopy in Combination with Multivariate Data Analysis. Agriculture (Switzerland), 2018, 8, 59.	1.4	10
35	Manual geo-rectification to improve the spatial accuracy of ortho-mosaics based on images from consumer-grade unmanned aerial vehicles (UAVs). Precision Agriculture, 2019, 20, 1199-1210.	3.1	10
36	A simulation of variable rate nitrogen application in winter wheat with soil and sensor information - An economic feasibility study. Agricultural Systems, 2021, 192, 103147.	3.2	10

#	Article	IF	Citations
37	Nonâ€destructive biomass estimation of <i>Oecophylla smaragdina</i> colonies: a model species for the ecological impact of ants. Insect Conservation and Diversity, 2015, 8, 464-473.	1.4	8
38	Preliminary Results of Clover and Grass Coverage and Total Dry Matter Estimation in Clover-Grass Crops Using Image Analysis. Journal of Imaging, 2017, 3, 59.	1.7	8
39	Robust Species Distribution Mapping of Crop Mixtures Using Color Images and Convolutional Neural Networks. Sensors, 2021, 21, 175.	2.1	8
40	Comparing methods for measuring the digestibility of miscanthus in bioethanol or biogas processing. GCB Bioenergy, 2017, 9, 168-175.	2.5	7
41	Cultivar and row distance interactions in perennial ryegrass. Acta Agriculturae Scandinavica - Section B Soil and Plant Science, 2009, 59, 335-341.	0.3	6
42	Assessment of grass lodging using texture and canopy height distribution features derived from UAV visual-band images. Agricultural and Forest Meteorology, 2021, 308-309, 108541.	1.9	6
43	Optimizing the number of consecutive seed harvests in red fescue (<i>Festuca rubra</i> L.) and perennial ryegrass (<i>Lolium perenne</i> L.) for yield, yield components and economic return. Acta Agriculturae Scandinavica - Section B Soil and Plant Science, 2013, 63, 1-10.	0.3	5
44	OPTIMUM DISTRIBUTION BETWEEN AUTUMN-APPLIED AND SPRING-APPLIED NITROGEN IN SEED PRODUCTION OF TALL FESCUE. Journal of Plant Nutrition, 2012, 35, 1394-1410.	0.9	4
45	Expression of starch-binding factor CBM20 in barley plastids controls the number of starch granules and the level of CO2 fixation. Journal of Experimental Botany, 2020, 71, 234-246.	2.4	3
46	Field scale agronomic and environmental consequences of overlapping N fertilizer application by disc spreaders. Field Crops Research, 2020, 255, 107901.	2.3	3
47	The use of linear mixed models for analysis of repeated measurements applied to water-soluble carbohydrates in perennial ryegrass for seed production. Acta Agriculturae Scandinavica - Section B Soil and Plant Science, 2009, 59, 151-156.	0.3	2
48	Single Seed NIR as a Fast Method to Predict Germination Ability in Pak Choi. NIR News, 2012, 23, 6-7.	1.6	2
49	Predicting seed yield in perennial ryegrass using repeated canopy reflectance measurements and PLSR. Acta Agriculturae Scandinavica - Section B Soil and Plant Science, 2009, 59, 414-423.	0.3	1
50	The Use of Remote Sensing to Determine Nitrogen Status in Perennial Ryegrass (Lolium perenne L.) for Seed Production. Nitrogen, 2021, 2, 229-243.	0.6	1
51	Agricultural Applications of NIR Reflectance and Transmittance. NIR News, 2009, 20, 4-4.	1.6	0
52	Potential Impact of Learning Management Zones for Site-Specific N Fertilisation: A Case Study for Wheat Crops. Nitrogen, 2022, 3, 387-403.	0.6	0