

Ildiko Van Rhijn

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3087915/publications.pdf>

Version: 2024-02-01

68
papers

2,910
citations

172457

29
h-index

182427

51
g-index

69
all docs

69
docs citations

69
times ranked

2643
citing authors

#	ARTICLE	IF	CITATIONS
1	Dual TCR- $\alpha\beta$ Expression on Mucosal-Associated Invariant T Cells as a Potential Confounder of TCR Interpretation. <i>Journal of Immunology</i> , 2022, 208, 1389-1395.	0.8	2
2	Atypical sideways recognition of CD1a by autoreactive $\alpha\beta$ T cell receptors. <i>Nature Communications</i> , 2022, 13, .	12.8	12
3	Synthetic mycobacterial diacyl trehaloses reveal differential recognition by human T cell receptors and the C-type lectin Mincle. <i>Scientific Reports</i> , 2021, 11, 2010.	3.3	7
4	Human skin is colonized by T cells that recognize CD1a independently of lipid. <i>Journal of Clinical Investigation</i> , 2021, 131, .	8.2	31
5	CD1 and MR1 recognition by human $\alpha\beta$ T cells. <i>Molecular Immunology</i> , 2021, 133, 95-100.	2.2	4
6	Multimodally profiling memory T cells from a tuberculosis cohort identifies cell state associations with demographics, environment and disease. <i>Nature Immunology</i> , 2021, 22, 781-793.	14.5	52
7	CD1a selectively captures endogenous cellular lipids that broadly block T cell response. <i>Journal of Experimental Medicine</i> , 2021, 218, .	8.5	24
8	Benzofuran sulfonates and small self-lipid antigens activate type II NKT cells via CD1d. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, .	7.1	8
9	Rational design of a hydrolysis-resistant mycobacterial phosphoglycolipid antigen presented by CD1c to T cells. <i>Journal of Biological Chemistry</i> , 2021, 297, 101197.	3.4	5
10	Human T cell response to CD1a and contact dermatitis allergens in botanical extracts and commercial skin care products. <i>Science Immunology</i> , 2020, 5, .	11.9	42
11	Human $\alpha\beta$ T cells recognize CD1b by two distinct mechanisms. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020, 117, 22944-22952.	7.1	34
12	CD1b Tetramers Broadly Detect T Cells That Correlate With Mycobacterial Exposure but Not Tuberculosis Disease State. <i>Frontiers in Immunology</i> , 2020, 11, 199.	4.8	22
13	Total Synthesis of a Mycolic Acid from <i>Mycobacterium tuberculosis</i> . <i>Angewandte Chemie - International Edition</i> , 2020, 59, 7555-7560.	13.8	14
14	Total Synthesis of a Mycolic Acid from <i>Mycobacterium tuberculosis</i> . <i>Angewandte Chemie</i> , 2020, 132, 7625-7630.	2.0	1
15	Peripheral Blood Mucosal-Associated Invariant T Cells in Tuberculosis Patients and Healthy <i>Mycobacterium tuberculosis</i> -Exposed Controls. <i>Journal of Infectious Diseases</i> , 2020, 222, 995-1007.	4.0	19
16	Asymmetric Total Synthesis of Mycobacterial Diacyl Trehaloses Demonstrates a Role for Lipid Structure in Immunogenicity. <i>ACS Chemical Biology</i> , 2020, 15, 1835-1841.	3.4	10
17	RISK6, a 6-gene transcriptomic signature of TB disease risk, diagnosis and treatment response. <i>Scientific Reports</i> , 2020, 10, 8629.	3.3	90
18	Total Synthesis of an Immunogenic Trehalose Phospholipid from <i>Salmonella</i> Typhi and Elucidation of Its <i>sn</i> -Regiochemistry by Mass Spectrometry. <i>Organic Letters</i> , 2019, 21, 5126-5131.	4.6	7

#	ARTICLE	IF	CITATIONS
19	CD1b presents self and <i>Borrelia burgdorferi</i> diacylglycerols to human T cells. <i>European Journal of Immunology</i> , 2019, 49, 737-746.	2.9	10
20	Discovery of <i>Salmonella</i> trehalose phospholipids reveals functional convergence with mycobacteria. <i>Journal of Experimental Medicine</i> , 2019, 216, 757-771.	8.5	20
21	A TCR $\hat{2}$ -Chain Motif Biases toward Recognition of Human CD1 Proteins. <i>Journal of Immunology</i> , 2019, 203, 3395-3406.	0.8	10
22	A T-cell receptor escape channel allows broad T-cell response to CD1b and membrane phospholipids. <i>Nature Communications</i> , 2019, 10, 56.	12.8	31
23	CD1b Tetramers Identify T Cells that Recognize Natural and Synthetic Diacylated Sulfoglycolipids from <i>Mycobacterium tuberculosis</i> . <i>Cell Chemical Biology</i> , 2018, 25, 392-402.e14.	5.2	23
24	Molecular recognition of microbial lipid-based antigens by T cells. <i>Cellular and Molecular Life Sciences</i> , 2018, 75, 1623-1639.	5.4	10
25	T cell autoreactivity directed toward CD1c itself rather than toward carried self lipids. <i>Nature Immunology</i> , 2018, 19, 397-406.	14.5	52
26	Total Synthesis of <i>Mycobacterium tuberculosis</i> Dideoxymycobactin $\hat{8}$ 38 and Stereoisomers: Diverse CD1a-Restricted T Cells Display a Common Hierarchy of Lipopeptide Recognition. <i>Chemistry - A European Journal</i> , 2017, 23, 1694-1701.	3.3	13
27	A molecular basis of human T cell receptor autoreactivity toward self-phospholipids. <i>Science Immunology</i> , 2017, 2, .	11.9	39
28	CD1b-mycolic acid tetramers demonstrate T cell fine specificity for mycobacterial lipid tails. <i>European Journal of Immunology</i> , 2017, 47, 1525-1534.	2.9	49
29	CD1b-autoreactive T cells contribute to hyperlipidemia-induced skin inflammation in mice. <i>Journal of Clinical Investigation</i> , 2017, 127, 2339-2352.	8.2	59
30	Mammalian CD1 and MR1 genes. <i>Immunogenetics</i> , 2016, 68, 515-523.	2.4	26
31	T cell receptor recognition of CD1b presenting a mycobacterial glycolipid. <i>Nature Communications</i> , 2016, 7, 13257.	12.8	59
32	Human autoreactive T cells recognize CD1b and phospholipids. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2016, 113, 380-385.	7.1	85
33	CD1 and mycobacterial lipids activate human T cells. <i>Immunological Reviews</i> , 2015, 264, 138-153.	6.0	72
34	Lipid and small-molecule display by CD1 and MR1. <i>Nature Reviews Immunology</i> , 2015, 15, 643-654.	22.7	120
35	Donor Unrestricted T Cells: A Shared Human T Cell Response. <i>Journal of Immunology</i> , 2015, 195, 1927-1932.	0.8	77
36	Expression Patterns of Bovine CD1 In Vivo and Assessment of the Specificities of the Anti-Bovine CD1 Antibodies. <i>PLoS ONE</i> , 2015, 10, e0121923.	2.5	11

#	ARTICLE	IF	CITATIONS
37	TCR Bias and Affinity Define Two Compartments of the CD1bâ€“Glycolipid-Specific T Cell Repertoire. <i>Journal of Immunology</i> , 2014, 192, 4054-4060.	0.8	64
38	Discovery of Invariant T Cells by Next-Generation Sequencing of the Human TCR Î±-Chain Repertoire. <i>Journal of Immunology</i> , 2014, 193, 5338-5344.	0.8	23
39	CD1a-autoreactive T cells recognize natural skin oils that function as headless antigens. <i>Nature Immunology</i> , 2014, 15, 177-185.	14.5	141
40	Targeted Delivery of Mycobacterial Antigens to Human Dendritic Cells via Siglec-7 Induces Robust T Cell Activation. <i>Journal of Immunology</i> , 2014, 193, 1560-1566.	0.8	54
41	Cutting Edge: CD1a Tetramers and Dextramers Identify Human Lipopeptideâ€“Specific T Cells Ex Vivo. <i>Journal of Immunology</i> , 2013, 191, 4499-4503.	0.8	70
42	Lipoproteins Are Major Targets of the Polyclonal Human T Cell Response to <i>Mycobacterium tuberculosis</i> . <i>Journal of Immunology</i> , 2013, 190, 278-284.	0.8	22
43	CD1a, CD1b, and CD1c in Immunity Against Mycobacteria. <i>Advances in Experimental Medicine and Biology</i> , 2013, 783, 181-197.	1.6	46
44	A conserved human T cell population targets mycobacterial antigens presented by CD1b. <i>Nature Immunology</i> , 2013, 14, 706-713.	14.5	187
45	The molecular basis for Mucosal-Associated Invariant T cell recognition of MR1 proteins. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, E1771-8.	7.1	85
46	The bovine CD1D gene has an unusual gene structure and is expressed but cannot present Î±-galactosylceramide with a C26 fatty acid. <i>International Immunology</i> , 2013, 25, 91-98.	4.0	16
47	Î± T Cell Homing to Skin and Migration to Skin-Draining Lymph Nodes Is CCR7 Independent. <i>Journal of Immunology</i> , 2012, 188, 578-584.	0.8	38
48	CD1b tetramers bind Î± T cell receptors to identify a mycobacterial glycolipid-reactive T cell repertoire in humans. <i>Journal of Experimental Medicine</i> , 2011, 208, 1741-1747.	8.5	132
49	Immune response of cattle immunized with a conjugate of the glycolipid glucose monomycolate and protein. <i>Veterinary Immunology and Immunopathology</i> , 2011, 142, 265-270.	1.2	5
50	CD1a-autoreactive T cells are a normal component of the human Î± T cell repertoire. <i>Nature Immunology</i> , 2010, 11, 1102-1109.	14.5	221
51	Crystal Structure of Bovine CD1b3 with Endogenously Bound Ligands. <i>Journal of Immunology</i> , 2010, 185, 376-386.	0.8	15
52	Lion (<i>Panthera leo</i>) and cheetah (<i>Acinonyx jubatus</i>) IFN-Î³ sequences. <i>Veterinary Immunology and Immunopathology</i> , 2010, 134, 296-298.	1.2	7
53	Conservation of mucosal associated invariant T (MAIT) cells and the MR1 restriction element in ruminants, and abundance of MAIT cells in spleen. <i>Veterinary Research</i> , 2010, 41, 62.	3.0	45
54	CD1c bypasses lysosomes to present a lipopeptide antigen with 12 amino acids. <i>Journal of Experimental Medicine</i> , 2009, 206, 1409-1422.	8.5	47

#	ARTICLE	IF	CITATIONS
55	Low cross-reactivity of T cell responses against lipids from <i>Mycobacterium bovis</i> and <i>M. avium paratuberculosis</i> during natural infection. <i>European Journal of Immunology</i> , 2009, 39, 3031-3041.	2.9	29
56	The evolved functions of CD1 during infection. <i>Current Opinion in Immunology</i> , 2009, 21, 397-403.	5.5	43
57	The bovine T cell receptor alpha/delta locus contains over 400 V genes and encodes V genes without CDR2. <i>Immunogenetics</i> , 2009, 61, 541-549.	2.4	22
58	Functional CD1d and/or NKT cell invariant chain transcript in horse, pig, African elephant and guinea pig, but not in ruminants. <i>Molecular Immunology</i> , 2009, 46, 1424-1431.	2.2	51
59	Two canine CD1a proteins are differentially expressed in skin. <i>Immunogenetics</i> , 2008, 60, 315-324.	2.4	28
60	Bovine tuberculosis as a model for human tuberculosis: advantages over small animal models. <i>Microbes and Infection</i> , 2008, 10, 711-715.	1.9	59
61	Massive, sustained $\gamma\delta$ T cell migration from the bovine skin in vivo. <i>Journal of Leukocyte Biology</i> , 2007, 81, 968-973.	3.3	28
62	Highly diverse TCR γ chain repertoire in bovine tissues due to the use of up to four D segments per γ chain. <i>Molecular Immunology</i> , 2007, 44, 3155-3161.	2.2	21
63	Role of lipid trimming and CD1 groove size in cellular antigen presentation. <i>EMBO Journal</i> , 2006, 25, 2989-2999.	7.8	50
64	The Bovine CD1 Family Contains Group 1 CD1 Proteins, but No Functional CD1d. <i>Journal of Immunology</i> , 2006, 176, 4888-4893.	0.8	64
65	T-cell activation by lipopeptide antigens. <i>Current Opinion in Immunology</i> , 2005, 17, 222-229.	5.5	22
66	CD1d-restricted T cell activation by nonlipidic small molecules. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 13578-13583.	7.1	91
67	Expansion of human gammadelta T cells after in vitro stimulation with <i>Campylobacter jejuni</i> . <i>International Immunology</i> , 2003, 15, 373-382.	4.0	18
68	<i>Campylobacter</i> DNA Is Present in Circulating Myelomonocytic Cells of Healthy Persons and in Persons with Guillain-Barré Syndrome. <i>Journal of Infectious Diseases</i> , 2002, 185, 262-265.	4.0	9