

Birgit Filipiak-Pittroff

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3113088/publications.pdf>

Version: 2024-02-01

10
papers

1,221
citations

1163117

8
h-index

1474206

9
g-index

10
all docs

10
docs citations

10
times ranked

851
citing authors

#	ARTICLE	IF	CITATIONS
1	Long-term effects of hydrolyzed formulae on atopic diseases in the GINI study. <i>Allergy: European Journal of Allergy and Clinical Immunology</i> , 2021, 76, 1903-1907.	5.7	17
2	Reply to Wjst. <i>Pediatric Allergy and Immunology</i> , 2018, 29, 458-459.	2.6	0
3	Full breastfeeding and allergies from infancy until adolescence in the GINI plus cohort. <i>Pediatric Allergy and Immunology</i> , 2018, 29, 96-101.	2.6	8
4	Allergies in high-risk schoolchildren after early intervention with cow's milk protein hydrolysates: 10-year results from the German Infant Nutritional Intervention (GINI) study. <i>Journal of Allergy and Clinical Immunology</i> , 2013, 131, 1565-1573.e5.	2.9	141
5	Reply. <i>Journal of Allergy and Clinical Immunology</i> , 2013, 132, 770-771.	2.9	5
6	Preventive effect of hydrolyzed infant formulas persists until age 6 years: Long-term results from the German Infant Nutritional Intervention Study (GINI). <i>Journal of Allergy and Clinical Immunology</i> , 2008, 121, 1442-1447.	2.9	222
7	Certain hydrolyzed formulas reduce the incidence of atopic dermatitis but not that of asthma: Three-year results of the German Infant Nutritional Intervention Study. <i>Journal of Allergy and Clinical Immunology</i> , 2007, 119, 718-725.	2.9	163
8	Effect of breast-feeding on the development of atopic dermatitis during the first 3 years of life—results from the GINI-birth cohort study. <i>Journal of Pediatrics</i> , 2004, 144, 602-607.	1.8	149
9	The effect of hydrolyzed cow's milk formula for allergy prevention in the first year of life: The German Infant Nutritional Intervention Study, a randomized double-blind trial. <i>Journal of Allergy and Clinical Immunology</i> , 2003, 111, 533-540.	2.9	392
10	Effect of exclusive breast-feeding and early solid food avoidance on the incidence of atopic dermatitis in high-risk infants at 1-year of age. <i>Pediatric Allergy and Immunology</i> , 2002, 13, 234-242.	2.6	124