Yongsheng Tong

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3134722/publications.pdf

Version: 2024-02-01

1478505 1125743 22 200 13 6 citations h-index g-index papers 22 22 22 192 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Symptomatology in 1,112 women screened positive and negative using the Edinburgh postnatal depression scale (EPDS): longitudinal observations from the first trimester to 6 weeks postpartum of a Chinese cohort. Journal of Psychosomatic Obstetrics and Gynaecology, 2022, 43, 453-463.	2.1	2
2	Comparison of Psycho-Social Factors Associated With Suicidal Ideation and Suicide Attempts Among People Living With HIV in Central West China. Frontiers in Public Health, 2022, 10, 832624.	2.7	2
3	Trends in suicide rates and the case-fatality of pesticide self-poisoning in an agricultural county in china, 2009 to 2014. Journal of Affective Disorders, 2021, 283, 52-59.	4.1	9
4	Comparisons of Characteristics Between Psychological Support Hotline Callers With and Without COVID-19 Related Psychological Problems in China. Frontiers in Psychiatry, 2021, 12, 648974.	2.6	7
5	Factors related to retinal nerve fiber layer thickness in bipolar disorder patients and major depression patients. BMC Psychiatry, 2021, 21, 301.	2.6	8
6	Suicide attempt risks among hotline callers with and without the coronavirus disease 2019 related psychological distress: a case-control study. BMC Psychiatry, 2021, 21, 363.	2.6	6
7	Auditory Hallucinations, Depressive Symptoms, and CurrentÂSuicidal Ideation or Behavior Among Patients with Acute-episode Schizophrenia. Archives of Suicide Research, 2021, , 1-16.	2.3	2
8	Comparison of Characteristics of Suicide Attempts Registered in Urban and Rural Areas in China. Frontiers in Psychiatry, 2021, 12, 805324.	2.6	0
9	Prospective study of association of characteristics of hotline psychological intervention in 778 high-risk callers with subsequent suicidal act. Australian and New Zealand Journal of Psychiatry, 2020, 54, 1182-1191.	2.3	9
10	Suicidal ideation, suicide attempts, and neurocognitive dysfunctions among patients with firstâ€episode schizophrenia. Suicide and Life-Threatening Behavior, 2020, 50, 1181-1188.	1.9	10
11	Interrelationships Between BDNF, Superoxide Dismutase, and Cognitive Impairment in Drug-Naive First-Episode Patients With Schizophrenia. Schizophrenia Bulletin, 2020, 46, 1498-1510.	4.3	42
12	Relationship of the high proportion of suicidal acts involving ingestion of pesticides to the low male-to-female ratio of suicide rates in China. Epidemiology and Psychiatric Sciences, 2020, 29, e114.	3.9	15
13	Association between TNF-alpha polymorphism and the age of first suicide attempt in chronic patients with schizophrenia. Aging, 2020, 12, 1433-1445.	3.1	7
14	Association of retinal nerve fiber abnormalities with serum CNTF and cognitive functions in schizophrenia patients. Peerl, 2020, 8, e9279.	2.0	14
15	Differences between medically treated and untreated non-fatal self-harm reported by hotline callers in China. PeerJ, 2019, 7, e7868.	2.0	2
16	Prevalence of psychiatric diagnoses in individuals who die by suicide or attempt suicide in China based on independent structured diagnostic interviews with different informants. Journal of Psychiatric Research, 2018, 98, 30-38.	3.1	8
17	Authors' reply. British Journal of Psychiatry, 2017, 210, 167-168.	2.8	О
18	DSM-IV Axis II personality disorders and suicide and attempted suicide in China. British Journal of Psychiatry, 2016, 209, 319-326.	2.8	13

#	Article	IF	CITATIONS
19	Suicidal Behavior in Relatives or Associates Moderates the Strength of Common Risk Factors for Suicide. Suicide and Life-Threatening Behavior, 2015, 45, 505-517.	1.9	4
20	Cohort-specific risk of suicide for different mental disorders in China. British Journal of Psychiatry, 2010, 196, 467-473.	2.8	36
21	Relationship of the High Proportion of Suicidal Acts Involving Ingestion of Pesticides to the Low Male-to-Female Ratio of Suicide Rates in China. SSRN Electronic Journal, 0, , .	0.4	O
22	Mental health problems among hotline callers during the early stage of COVID-19 pandemic. PeerJ, 0, 10, e13419.	2.0	4