

Claudete A Mangolin

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3186777/publications.pdf>

Version: 2024-02-01

81
papers

987
citations

516710

16
h-index

552781

26
g-index

81
all docs

81
docs citations

81
times ranked

1041
citing authors

#	ARTICLE	IF	CITATIONS
1	Cell wall polysaccharides from <i>in vitro</i> propagated <i>Cereus hildmannianus</i> K. Schum. plants. <i>Natural Product Research</i> , 2022, 36, 2424-2428.	1.8	4
2	Biochemical composition of the pericarp cell wall of popcorn inbred lines with different popping expansion. <i>Current Research in Food Science</i> , 2022, 5, 102-106.	5.8	6
3	<i>Cereus hildmannianus</i> (K.) Schum. (Cactaceae): Ethnomedical uses, phytochemistry and biological activities. <i>Journal of Ethnopharmacology</i> , 2021, 264, 113339.	4.1	10
4	Prospecção biotecnológica de bactérias endofíticas associadas a cactáceas. <i>Revista Ibero-americana De Ciências Ambientais</i> , 2021, 12, 567-579.	0.1	0
5	ISSR markers to assess genetic diversity of cultivated populations from artificial selection of <i>Stevia rebaudiana</i> (Bert.) Bertonii. <i>Breeding Science</i> , 2020, 70, 508-514.	1.9	8
6	Selection based on meiotic behavior in <i>Urochloa decumbens</i> hybrids from non-shattered seed. <i>Tropical Grasslands - Forrajes Tropicales</i> , 2020, 8, 133-140.	0.5	5
7	Proteomic analysis of axillary buds of sugarcane at different cutting stages: evidence for alterations in axillary bud gene expression. <i>Crop and Pasture Science</i> , 2019, 70, 622.	1.5	3
8	Development and use of retrotransposons-based markers (IRAP/REMAP) to assess genetic divergence among table grape cultivars. <i>Plant Genetic Resources: Characterisation and Utilisation</i> , 2019, 17, 272-279.	0.8	5
9	High genetic differentiation of grapevine rootstock varieties determined by molecular markers and artificial neural networks. <i>Acta Scientiarum - Agronomy</i> , 2019, 42, e43475.	0.6	7
10	Clonal propagation of cv. Italy grapes and the generation of genetic divergence among vineyards. <i>Scientia Horticulturae</i> , 2019, 244, 263-269.	3.6	5
11	Extraction of Total Protein from Axillary Buds of Sugarcane (<i>Saccharum</i> spp.) for Proteomic Analysis. <i>Sugar Tech</i> , 2018, 20, 95-99.	1.8	1
12	Genetic variability of wild poinsettia populations in Brazilian agroecosystems. <i>Cogent Food and Agriculture</i> , 2018, 4, 1551749.	1.4	1
13	Differential genetic stability in vineyards of the cultivar 'Italy' (Vitis vinifera L.) cultivated in different regions of Southern and Southwestern Brazil. <i>Ciencia E Técnica Vitivinícola</i> , 2018, 33, 66-77.	0.9	1
14	Genome wide association study for gray leaf spot resistance in tropical maize core. <i>PLoS ONE</i> , 2018, 13, e0199539.	2.5	27
15	Genetic structure of phenotypic variants and somaclones of the genus <i>Cereus</i> (Cactaceae) as revealed by AFLP markers. <i>Journal of Horticultural Science and Biotechnology</i> , 2017, 92, 325-333.	1.9	5
16	Changes on microsatellites of expressed sequence tag of sugarcane (<i>Saccharum</i> spp) during vegetative propagation. <i>Genetics and Molecular Research</i> , 2017, 16, .	0.2	1
17	Genetic variability among elite popcorn lines based on molecular and morphoagronomic characteristics. <i>Genetics and Molecular Research</i> , 2017, 16, .	0.2	2
18	Genetic divergence and admixture of ancestral genome groups in the sugarcane variety 'RB867515' (<i>Saccharum</i> spp). <i>Genetics and Molecular Research</i> , 2016, 15, .	0.2	0

#	ARTICLE	IF	CITATIONS
19	Suplementos orgânicos para el cultivo in vitro del híbrido <i>Laeliocattleya</i> (Orchidaceae). <i>Idesia</i> , 2016, 34, 47-54.	0.3	14
20	Development of retrotransposon-based markers IRAP and REMAP for cassava (<i>Manihot esculenta</i>). <i>Genetics and Molecular Research</i> , 2016, 15, .	0.2	3
21	Genetic variability in the natural populations of <i>Lasioderma serricorne</i> (F.) (Coleoptera: Anobiidae), detected by RAPD markers and by esterase isozymes. <i>Bulletin of Entomological Research</i> , 2016, 106, 47-53.	1.0	5
22	Genetic structure and molecular divergence among samples of mandacaru (<i>Cereus</i> spp.; Cactaceae) as revealed by microsatellite markers. <i>Biochemical Systematics and Ecology</i> , 2016, 64, 38-45.	1.3	10
23	Genetic and chemical diversity in seeds of cactus mandacaru (<i>Cereus</i> sp.) from two edaphoclimatic regions contrasting. <i>Anais Da Academia Brasileira De Ciencias</i> , 2015, 87, 765-776.	0.8	9
24	Matrilineage differentiation of the genus <i>Tetragonisca</i> using mitochondrial DNA markers and the polymerase chain reaction-restriction fragment length polymorphism technique. <i>Genetics and Molecular Research</i> , 2015, 14, 12828-12840.	0.2	2
25	Genetic diversity of popcorn genotypes using molecular analysis. <i>Genetics and Molecular Research</i> , 2015, 14, 9829-9840.	0.2	4
26	In silico analysis of the 16S rRNA gene of endophytic bacteria, isolated from the aerial parts and seeds of important agricultural crops. <i>Genetics and Molecular Research</i> , 2015, 14, 9703-9721.	0.2	22
27	Genetic diversity assessed by microsatellite markers in sweet corn cultivars. <i>Scientia Agricola</i> , 2015, 72, 513-519.	1.2	13
28	Lipid profile and antiproliferative activity of callus cultures of <i>Cereus peruvianus</i> Mill. <i>Industrial Crops and Products</i> , 2015, 69, 408-414.	5.2	19
29	Evidence of High Gene Flow Between Samples of Horseweed (<i>Conyza canadensis</i>) and Hairy Fleabane (<i>Conyza bonariensis</i>) as Revealed by Isozyme Polymorphisms. <i>Weed Science</i> , 2015, 63, 604-612.	1.5	11
30	Population structure and genetic diversity of Brazilian popcorn germplasm inferred by microsatellite markers. <i>Electronic Journal of Biotechnology</i> , 2015, 18, 181-187.	2.2	19
31	Genetic diversity and structure of populations in <i>Pilosocereus gounellei</i> (F.A.C.Weber ex K.Schum.) (Cactaceae) in the Caatinga biome as revealed by heterologous microsatellite primers. <i>Biochemical Systematics and Ecology</i> , 2015, 58, 7-12.	1.3	11
32	High Polymorphism in Est-SSR Loci for Cellulose Synthase and Î ² -Amylase of Sugarcane Varieties (<i>Saccharum</i> spp.) Used by the Industrial Sector for Ethanol Production. <i>Applied Biochemistry and Biotechnology</i> , 2015, 175, 965-973.	2.9	3
33	Primary Characterization and Evaluation of Anti Ulcerogenic Activity of an Aqueous Extract from Callus Culture of <i>Cereus peruvianus</i> Mill. (Cactaceae). <i>Current Pharmaceutical Biotechnology</i> , 2015, 16, 462-467.	1.6	9
34	Genetic structure of <i>Pilosocereus gounellei</i> (Cactaceae) as revealed by AFLP marker to guide proposals for improvement and restoration of degraded areas in Caatinga biome. <i>Genetics and Molecular Research</i> , 2015, 14, 16966-16974.	0.2	6
35	Esterase polymorphism for genetic diversity analysis of some accessions of a native forage grass, <i>Mesosetum chaseae</i> Lucas, from the Brazilian Pantanal. <i>Tropical Grasslands - Forrajes Tropicales</i> , 2015, 3, 194.	0.5	1
36	Seleção de primers para análise inter simple sequence repeats na cultivar 'Itália' de <i>Vitis vinifera</i> L.. <i>Ciencia E Tecnica Vitivinicola</i> , 2014, 29, 81-87.	0.9	1

#	ARTICLE	IF	CITATIONS
37	Genetic divergence among sweet corn lines estimated by microsatellite markers. <i>Genetics and Molecular Research</i> , 2014, 13, 10415-10426.	0.2	6
38	Population genetics of the sugarcane borer <i>Diatraea saccharalis</i> (Fabr.) (Lepidoptera: Crambidae). <i>Acta Scientiarum - Agronomy</i> , 2014, 36, 189.	0.6	8
39	<i>Conyza sumatrensis</i>: A new weed species resistant to glyphosate in the Americas. <i>Weed Biology and Management</i> , 2014, 14, 106-114.	1.4	21
40	Caracterizaço gentica de amostras de Conyza sp. do estado do Paran. <i>Planta Daninha</i> , 2014, 32, 173-179.	0.5	5
41	Diallel analysis of maize inbred lines for grain yield, oil and protein content. <i>Crop Breeding and Applied Biotechnology</i> , 2014, 14, 23-28.	0.4	12
42	Use of differential levels of mean observed heterozygosity in microsatellite loci of commercial varieties of sugarcane (<i>Saccharum</i> spp). <i>Genetics and Molecular Research</i> , 2014, 13, 10130-10141.	0.2	5
43	A Collection of Popcorn as a Reservoir of Genes for the Generation of Lineages. <i>Molecular Biotechnology</i> , 2013, 53, 300-307.	2.4	9
44	Genetic relationships among accessions of mandacaru (<i>Cereus</i> spp.: Cactaceae) using amplified fragment length polymorphisms (AFLP). <i>Biochemical Systematics and Ecology</i> , 2013, 48, 12-19.	1.3	10
45	Genetic diversity analysis of <i>Capsicum</i> spp germplasm bank accessions based on \hat{I}^2 -esterase polymorphism. <i>Genetics and Molecular Research</i> , 2013, 12, 1155-1167.	0.2	1
46	Molecular characterization of endophytes isolated from <i>Saccharum</i> spp based on esterase and ribosomal DNA (ITS1-5.8S-ITS2) analyses. <i>Genetics and Molecular Research</i> , 2013, 12, 4095-4105.	0.2	7
47	Use of complex supplements and light-differential effects for micropropagation of <i>Hadrolaelia purpurata</i> (= <i>Laelia purpurata</i>) and <i>Encyclia randii</i> orchids. <i>Acta Scientiarum - Agronomy</i> , 2012, 34, .	0.6	4
48	Activated charcoal and graphite for the micropropagation of <i>Cattleya bicolor</i> Lindl. and a orchid double-hybrid "BLC Pastoral Innocence"™. <i>Acta Scientiarum - Agronomy</i> , 2012, 34, .	0.6	7
49	Selection of high heterozygosity popcorn varieties in Brazil based on SSR markers. <i>Genetics and Molecular Research</i> , 2012, 11, 1851-1860.	0.2	6
50	Application of microsatellite markers to evaluate the heterozygosity from the popcorn composite CMS43 (<i>Zea mays</i> L.) during eight cycles of selection. <i>Plant Breeding</i> , 2012, 131, 479-485.	1.9	4
51	Esterase polymorphisms for analysis of genetic diversity and structure of soybean (<i>Glycine max</i>) cultivars. <i>Genetics and Molecular Research</i> , 2011, 10, 2472-2481.	0.2	1
52	The use of esterase polymorphism for analysis of the genetic diversity and structure of stevia (<i>Stevia</i>) Tj ETQq0 0 0 13 BT /Overlock 10 Tf		
53	Esterase Polymorphism and the Analysis of Genetic Diversity and Structure in Cactus Populations Descended from <i>Cereus peruvianus</i> Plants Regenerated In Vitro. <i>Biochemical Genetics</i> , 2011, 49, 270-282.	1.7	12
54	Diversidade gentica de dourado utilizado em programas de repovoamento no rio Paranapanema. <i>Pesquisa Agropecuaria Brasileira</i> , 2011, 46, 167-173.	0.9	4

#	ARTICLE	IF	CITATIONS
55	Potential use of major royal jelly proteins (MRJPs) as molecular markers for royal jelly production in Africanized honeybee colonies. <i>Apidologie</i> , 2010, 41, 160-168.	2.0	17
56	Microsatellite analysis of the parental contribution of <i>Piaractus mesopotamicus</i> to the production of offspring in the semi-natural system of reproduction. <i>Brazilian Archives of Biology and Technology</i> , 2010, 53, 389-396.	0.5	10
57	Isoenzymatic polymorphism in the leaf-cutting ant <i>Atta capiguara</i> Gonsalves (hymenoptera:) Tj ETQq1 1 0.784314 rgBT /Overlock 10	1.2	3
58	Caracterizao gentica de <i>Brycon orbignyanus</i> utilizando o sistema seminatural. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2010, 62, 184-191.	0.4	11
59	Avaliao gentica de populaes naturais e de estoques de um programa de repovoamento de pacu (<i>Piaractus mesopotamicus</i>) utilizando marcadores microssatlite. <i>Arquivo Brasileiro De Medicina Veterinaria E Zootecnia</i> , 2010, 62, 954-963.	0.4	4
60	Genetic diversity of breeding popcorn lines determined by SSR markers. <i>Electronic Journal of Biotechnology</i> , 2010, 13, .	2.2	0
61	Efficiency of RAPD versus SSR markers for determining genetic diversity among popcorn lines. <i>Genetics and Molecular Research</i> , 2010, 9, 9-18.	0.2	56
62	Esterase Polymorphism for Analysis of Genetic Diversity and Structure of Wild Poinsettia (<i>Euphorbia</i>) Tj ETQq0 0 0 rgBT /Overlock 10 Tf	1.5	16
63	IN VITRO CELL AND TISSUE CULTURE OF THE <i>CEREUS PERUVIANUS</i> MILL. (CACTACEAE) FOR THE CONSERVATION AND MANAGEMENT OF BIODIVERSITY: CURRENT STATUS. <i>Acta Horticulturae</i> , 2009, , 249-254.	0.2	2
64	Genetic diversity in somatic mutants of grape (<i>Vitis vinifera</i>) cultivar Italia based on random amplified polymorphic DNA. <i>Genetics and Molecular Research</i> , 2009, 8, 28-38.	0.2	16
65	Soybean (<i>Glycine max</i>) Root Lignification Induced by Ferulic Acid. The Possible Mode of Action. <i>Journal of Chemical Ecology</i> , 2008, 34, 1230-1241.	1.8	102
66	Genetic diversity associated with in vitro and conventional bud propagation of <i>Saccharum</i> varieties using RAPD analysis. <i>Plant Breeding</i> , 2008, 127, 160-165.	1.9	16
67	Heterozygosity following half-sib recurrent selection in popcorn using isoenzyme markers. <i>Electronic Journal of Biotechnology</i> , 2008, 11, 0-0.	2.2	3
68	Seed germination of the <i>Cereus peruvianus</i> Mill. (Cactaceae) somaclones follows a relatively simple protocol. <i>Seed Science and Technology</i> , 2008, 36, 595-600.	1.4	8
69	Molecular mapping in tropical maize (<i>Zea mays</i> L.) using microsatellite markers. 1. Map construction and localization of loci showing distorted segregation. <i>Hereditas</i> , 2004, 139, 96-106.	1.4	37
70	Molecular mapping in tropical maize (<i>Zea mays</i> L.) using microsatellite markers. 2. Quantitative trait loci (QTL) for grain yield, plant height, ear height and grain moisture. <i>Hereditas</i> , 2004, 139, 107-115.	1.4	77
71	Mapping QTLs for kernel oil content in a tropical maize population. <i>Euphytica</i> , 2004, 137, 251-259.	1.2	43
72	RAPD markers to evaluate callus tissue of <i>Cereus peruvianus</i> Mill. (Cactaceae) maintained in different growth regulator combinations. <i>Biochemical Genetics</i> , 2002, 40, 351-358.	1.7	16

#	ARTICLE	IF	CITATIONS
73	Genetic diversity in tropical maize inbred lines: heterotic group assignment and hybrid performance determined by RFLP markers. <i>Plant Breeding</i> , 2000, 119, 491-496.	1.9	70
74	Two-dimensional electrophoresis of <i>Cereus peruvianus</i> (Cactaceae) callus tissue proteins. <i>Electrophoresis</i> , 1999, 20, 626-629.	2.4	11
75	Effect of 5-Aminouracil on Callus Tissue Growth and Mdh Genes Expression in <i>Cereus peruvianus</i> (Cactaceae). <i>Cytologia</i> , 1997, 62, 283-289.	0.6	2
76	Isozyme variability in plants regenerated from calli of <i>Cereus peruvianus</i> (Cactaceae). <i>Biochemical Genetics</i> , 1997, 35, 189-204.	1.7	18
77	Malate dehydrogenase isozymes (MDH; EC 1.1.1.37) in long-term callus culture of <i>Cereus peruvianus</i> (Cactaceae) exposed to sugar and temperature stress. <i>Biochemical Genetics</i> , 1997, 35, 155-164.	1.7	17
78	Isozyme extraction from shoot tissue of <i>Cereus peruvianus</i> (Cactaceae) for electrophoretic analysis. <i>Biochemical Genetics</i> , 1997, 35, 205-210.	1.7	2
79	Alcohol dehydrogenase (EC 1.1.1.1) isozymes as markers at 2,4-dichlorophenoxyacetic acid $1/2$ kinetin combinations in callus cultures of <i>Cereus peruvianus</i> (Cactaceae). <i>Biochemical Genetics</i> , 1994, 32, 191-200.	1.7	10
80	Isozyme patterns in callus cultures and in plants regenerated from calli of <i>Cereus peruvianus</i> (Cactaceae). <i>Biochemical Genetics</i> , 1994, 32, 237-247.	1.7	22
81	Malate dehydrogenase (MDH; EC 1.1.1.37) isozymes in tissues and callus cultures of <i>Cereus peruvianus</i> (cactaceae). <i>Biochemical Genetics</i> , 1993, 31, 167-172.	1.7	18