Georges Leclercq

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3205268/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Small-scale manufacturing of neoantigen-encoding messenger RNA for early-phase clinical trials. Cytotherapy, 2022, 24, 213-222.	0.7	8
2	A Novel Non-Coding Variant in DCLRE1C Results in Deregulated Splicing and Induces SCID Through the Generation of a Truncated ARTEMIS Protein That Fails to Support V(D)J Recombination and DNA Damage Repair. Frontiers in Immunology, 2021, 12, 674226.	4.8	2
3	<i>In vitro</i> OP9-DL1 co-culture and subsequent maturation in the presence of IL-21 generates tumor antigen-specific T cells with a favorable less-differentiated phenotype and enhanced functionality. Oncolmmunology, 2021, 10, 1954800.	4.6	3
4	T-BET and EOMES Accelerate and Enhance Functional Differentiation of Human Natural Killer Cells. Frontiers in Immunology, 2021, 12, 732511.	4.8	0
5	T-BET and EOMES Accelerate and Enhance Functional Differentiation of Human Natural Killer Cells. Frontiers in Immunology, 2021, 12, 732511.	4.8	24
6	Human Thymic CD10+ PD-1+ Intraepithelial Lymphocyte Precursors Acquire Interleukin-15 Responsiveness at the CD1a– CD95+ CD28– CCR7– Developmental Stage. International Journal of Molecular Sciences, 2020, 21, 8785.	4.1	7
7	Conventional and Computational Flow Cytometry Analyses Reveal Sustained Human Intrathymic T Cell Development From Birth Until Puberty. Frontiers in Immunology, 2020, 11, 1659.	4.8	3
8	Distinct and temporary-restricted epigenetic mechanisms regulate human αβ and γδT cell development. Nature Immunology, 2020, 21, 1280-1292.	14.5	43
9	HES1 and HES4 have non-redundant roles downstream of Notch during early human T-cell development. Haematologica, 2020, 106, 130-141.	3.5	20
10	Distinct Notch1 and <i>BCL11B</i> requirements mediate human γδ/αβ T cell development. EMBO Reports, 2020, 21, e49006.	4.5	31
11	The transcription factor ETS1 is an important regulator of human NK cell development and terminal differentiation. Blood, 2020, 136, 288-298.	1.4	33
12	T-cells with a single tumor antigen-specific T-cell receptor can be generated <i>in vitro</i> from clinically relevant stem cell sources. Oncolmmunology, 2020, 9, 1727078.	4.6	4
13	Rapid and Effective Generation of Nanobody Based CARs using PCR and Gibson Assembly. International Journal of Molecular Sciences, 2020, 21, 883.	4.1	24
14	Treatment of a patient with severe cytomegalovirus (CMV) infection after haploidentical stem cell transplantation with donor-derived CMV-specific T cells. Acta Clinica Belgica, 2020, 76, 1-5.	1.2	1
15	Integrated scRNA-Seq Identifies Human Postnatal Thymus Seeding Progenitors and Regulatory Dynamics of Differentiating Immature Thymocytes. Immunity, 2020, 52, 1088-1104.e6.	14.3	79
16	Regulatory Role for NK Cells in a Mouse Model of Systemic Juvenile Idiopathic Arthritis. Journal of Immunology, 2019, 203, 3339-3348.	0.8	14
17	Porcine NK cells display features associated with antigen-presenting cells. Journal of Leukocyte Biology, 2018, 103, 129-140.	3.3	22
18	Innate lymphoid cells in the upper airways: importance of CD117 and IL-1RI expression. European Respiratory Journal, 2018, 52, 1800742.	6.7	14

#	Article	IF	CITATIONS
19	Comparative analysis of the immune responses induced by native versus recombinant versions of the ASP-based vaccine against the bovine intestinal parasite Cooperia oncophora. International Journal for Parasitology, 2018, 48, 41-49.	3.1	9
20	Antigen receptor-redirected T cells derived from hematopoietic precursor cells lack expression of the endogenous TCR/CD3 receptor and exhibit specific antitumor capacities. Oncolmmunology, 2017, 6, e1283460.	4.6	22
21	A Murine Intestinal Intraepithelial NKp46-Negative Innate Lymphoid Cell Population Characterized by Group 1 Properties. Cell Reports, 2017, 19, 1431-1443.	6.4	24
22	The checkpoint for agonist selection precedes conventional selection in human thymus. Science Immunology, 2017, 2, .	11.9	40
23	Inflammatory Gene Expression Profile and Defective Interferonâ€î³ and Granzyme K in Natural Killer Cells From Systemic Juvenile Idiopathic Arthritis Patients. Arthritis and Rheumatology, 2017, 69, 213-224.	5.6	67
24	Differential Effects of CORM-2 and CORM-401 in Murine Intestinal Epithelial MODE-K Cells under Oxidative Stress. Frontiers in Pharmacology, 2017, 8, 31.	3.5	29
25	The Ly49E Receptor Inhibits the Immune Control of Acute Trypanosoma cruzi Infection. Frontiers in Immunology, 2016, 7, 472.	4.8	5
26	Activation by SLAM Family Receptors Contributes to NK Cell Mediated "Missing-Self―Recognition. PLoS ONE, 2016, 11, e0153236.	2.5	10
27	GATA3 induces human T-cell commitment by restraining Notch activity and repressing NK-cell fate. Nature Communications, 2016, 7, 11171.	12.8	57
28	NKT sublineage specification and survival requires the ubiquitin-modifying enzyme TNFAIP3/A20. Journal of Experimental Medicine, 2016, 213, 1973-1981.	8.5	31
29	Host protective ASP-based vaccine against the parasitic nematode Ostertagia ostertagi triggers NK cell activation and mixed IgG1-IgG2 response. Scientific Reports, 2016, 6, 29496.	3.3	20
30	Expression of the inhibitory Ly49E receptor is not critically involved in the immune response against cutaneous, pulmonary or liver tumours. Scientific Reports, 2016, 6, 30564.	3.3	7
31	The role of Ly49E receptor expression on murine intraepithelial lymphocytes in intestinal cancer development and progression. Cancer Immunology, Immunotherapy, 2016, 65, 1365-1375.	4.2	4
32	Difference in replication of low-passage MCMV HaNa1 in BALB/c, C57BL/6 and NOD mice and role of different branches of immunity in susceptibility. Virus Research, 2016, 221, 38-46.	2.2	2
33	ID: 195. Cytokine, 2015, 76, 99-100.	3.2	Ο
34	Mitochondria and NADPH oxidases are the major sources of TNF-α/cycloheximide-induced oxidative stress in murine intestinal epithelial MODE-K cells. Cellular Signalling, 2015, 27, 1141-1158.	3.6	22
35	In vitro human embryonic stem cell hematopoiesis mimics MYB-independent yolk sac hematopoiesis. Haematologica, 2015, 100, 157-166.	3.5	40
36	Antioxidant potential of CORM-A1 and resveratrol during TNF-α/cycloheximide-induced oxidative stress and apoptosis in murine intestinal epithelial MODE-K cells. Toxicology and Applied Pharmacology, 2015, 288, 161-178.	2.8	38

#	Article	IF	CITATIONS
37	Chimeric Antigen Receptor Transgenic, T Cell Receptor/CD3 Negative Monospecific T Cells Generated from Cord Blood CD34 Positive Cells. Blood, 2015, 126, 3087-3087.	1.4	1
38	Contribution of the Ly49E Natural Killer Receptor in the Immune Response to Plasmodium berghei Infection and Control of Hepatic Parasite Development. PLoS ONE, 2014, 9, e87463.	2.5	4
39	Ly49E Expression on CD8αα-Expressing Intestinal Intraepithelial Lymphocytes Plays No Detectable Role in the Development and Progression of Experimentally Induced Inflammatory Bowel Diseases. PLoS ONE, 2014, 9, e110015.	2.5	9
40	Systemic Juvenile Idiopathic Arthritis–like Syndrome in Mice Following Stimulation of the Immune System With Freund's Complete Adjuvant: Regulation by Interferonâ€Î³. Arthritis and Rheumatology, 2014, 66, 1340-1351.	5.6	71
41	Notch3 Activation Is Sufficient but Not Required for Inducing Human T-Lineage Specification. Journal of Immunology, 2014, 193, 5997-6004.	0.8	17
42	Spray-Dried Polyelectrolyte Microparticles in Oral Antigen Delivery: Stability, Biocompatibility, and Cellular Uptake. Biomacromolecules, 2014, 15, 2301-2309.	5.4	18
43	In vitro human embryonic stem cell hematopoiesis mimics MYB independent yolk sac hematopoiesis. Experimental Hematology, 2014, 42, S63.	0.4	1
44	β-Glucan microparticles are good candidates for mucosal antigen delivery in oral vaccination. Journal of Controlled Release, 2013, 172, 671-678.	9.9	123
45	cMYB expression during human in vitro hematopoiesis. Experimental Hematology, 2013, 41, S33.	0.4	Ο
46	Abundant stage-dependent Ly49E expression by liver NK cells is not essential for their differentiation and function. Journal of Leukocyte Biology, 2013, 93, 699-711.	3.3	18
47	Education of Murine NK Cells Requires Both <i>cis</i> and <i>trans</i> Recognition of MHC Class I Molecules. Journal of Immunology, 2013, 191, 5044-5051.	0.8	39
48	Specific Notch receptor–ligand interactions control human TCR-αβ/γÎ′ development by inducing differential Notch signal strength. Journal of Experimental Medicine, 2013, 210, 683-697.	8.5	95
49	Differential <i>Ly49e</i> Expression Pathways in Resting versus TCR-Activated Intraepithelial γδT Cells. Journal of Immunology, 2013, 190, 1982-1990.	0.8	12
50	Specific Notch receptor–ligand interactions control human TCR-ab/gd development by inducing differential Notch signal strength. Journal of Cell Biology, 2013, 201, i2-i2.	5.2	0
51	Notch induces human T-cell receptor Î ³ δ+ thymocytes to differentiate along a parallel, highly proliferative and bipotent CD4 CD8 double-positive pathway. Leukemia, 2012, 26, 127-138.	7.2	26
52	TNF-α/Cycloheximide-Induced Oxidative Stress and Apoptosis in Murine Intestinal Epithelial MODE-K Cells. Current Pharmaceutical Design, 2012, 18, 4414-4425.	1.9	24
53	RHAMM/HMMR (CD168) is not an ideal target antigen for immunotherapy of acute myeloid leukemia. Haematologica, 2012, 97, 1539-1547.	3.5	32
54	Matrix Metalloprotease 8-Dependent Extracellular Matrix Cleavage at the Blood-CSF Barrier Contributes to Lethality during Systemic Inflammatory Diseases. Journal of Neuroscience, 2012, 32, 9805-9816.	3.6	91

#	Article	IF	CITATIONS
55	In vitro generation of immune cells from pluripotent stem cells. Frontiers in Bioscience - Landmark, 2011, 16, 1488.	3.0	8
56	T-lymphoid differentiation potential measured in vitro is higher in CD34+CD38-/lo hematopoietic stem cells from umbilical cord blood than from bone marrow and is an intrinsic property of the cells. Haematologica, 2011, 96, 646-654.	3.5	33
57	Jagged2 acts as a Delta-like Notch ligand during early hematopoietic cell fate decisions. Blood, 2011, 117, 4449-4459.	1.4	89
58	Inhibitory receptors specific for MHC class I educate murine NK cells but not CD8αα intestinal intraepithelial T lymphocytes. Blood, 2011, 118, 339-347.	1.4	15
59	CD27â€deficient mice show normal NKâ€cell differentiation but impaired function upon stimulation. Immunology and Cell Biology, 2011, 89, 803-811.	2.3	26
60	Langerhans cells are not required for epidermal Vγ3 T cell homeostasis and function. Journal of Leukocyte Biology, 2011, 90, 61-68.	3.3	10
61	Continuous CD27 triggering <i>in vivo</i> strongly reduces NK cell numbers. European Journal of Immunology, 2010, 40, 1107-1117.	2.9	23
62	Human T Cell Differentiation: New Techniques, Old Challenges. , 2010, , 351-371.		0
63	Type I interferon drives tumor necrosis factor–induced lethal shock. Journal of Experimental Medicine, 2009, 206, 1873-1882.	8.5	74
64	Activated CD4 ⁺ CD25 ⁺ regulatory T cells inhibit osteoclastogenesis and collagen-induced arthritis. Annals of the Rheumatic Diseases, 2009, 68, 744-750.	0.9	133
65	Functionally Mature CD4 and CD8 TCRαβ Cells Are Generated in OP9-DL1 Cultures from Human CD34+ Hematopoietic Cells. Journal of Immunology, 2009, 183, 4859-4870.	0.8	46
66	Generation of T Cells from Human Embryonic Stem Cell-Derived Hematopoietic Zones. Journal of Immunology, 2009, 182, 6879-6888.	0.8	186
67	Differential expression of CD97 on human lymphocyte subsets and limited effect of CD97 antibodies on allogeneic T-cell stimulation. Immunology Letters, 2009, 123, 160-168.	2.5	18
68	A Role for cis Interaction between the Inhibitory Ly49A Receptor and MHC Class I for Natural Killer Cell Education. Immunity, 2009, 30, 337-347.	14.3	111
69	An early decrease in Notch activation is required for human TCR-αβ lineage differentiation at the expense of TCR-γδT cells. Blood, 2009, 113, 2988-2998.	1.4	97
70	Notch signaling is required for proliferation but not for differentiation at a well-defined β-selection checkpoint during human T-cell development. Blood, 2009, 113, 3254-3263.	1.4	70
71	CD4 and CD8 TCRαβ Cells Are selected On MHC Expressed On Thymocyte Precursors in OP9-DL1 Cultures Blood, 2009, 114, 3670-3670.	1.4	1
72	Human intrathymic development: a selective approach. Seminars in Immunopathology, 2008, 30, 411-423.	6.1	29

#	Article	IF	CITATIONS
73	CD27 Defines Phenotypically and Functionally Different Human NK Cell Subsets. Journal of Immunology, 2008, 180, 3739-3745.	0.8	173
74	Ly49E-dependent inhibition of natural killer cells by urokinase plasminogen activator. Blood, 2008, 112, 5046-5051.	1.4	20
75	Generation of T Cells from Human Embryonic Stem Cells Blood, 2008, 112, 1527-1527.	1.4	0
76	Mouse TCRαβ+CD8αα Intraepithelial Lymphocytes Express Genes That Down-Regulate Their Antigen Reactivity and Suppress Immune Responses. Journal of Immunology, 2007, 178, 4230-4239.	0.8	132
77	KLRG1 binds cadherins and preferentially associates with SHIP-1. International Immunology, 2007, 19, 391-400.	4.0	94
78	Interactions of Ly49 Family Receptors with MHC Class I Ligands in <i>trans</i> and <i>cis</i> . Journal of Immunology, 2007, 178, 1277-1284.	0.8	84
79	Notch signaling induces cytoplasmic CD3ϵ expression in human differentiating NK cells. Blood, 2007, 110, 2696-2703.	1.4	53
80	T-, B- and NK-lymphoid, but not myeloid cells arise from human CD34+CD38â^`CD7+ common lymphoid progenitors expressing lymphoid-specific genes. Leukemia, 2007, 21, 311-319.	7.2	77
81	Sa.122. Il-15 Augments CD4+ T-Cell Proliferation by Inhibiting the Suppressive Function of CD25highCD4+ Regulatory T-Cells. Clinical Immunology, 2006, 119, S148-S149.	3.2	Ο
82	Overexpression of HES-1 is not sufficient to impose T-cell differentiation on human hematopoietic stem cells. Blood, 2006, 107, 2879-2881.	1.4	15
83	A unique lymphotoxin Âbeta-dependent pathway regulates thymic emigration of VÂ14 invariant natural killer T cells. Proceedings of the National Academy of Sciences of the United States of America, 2006, 103, 9160-9165.	7.1	32
84	Negative thymocyte selection to HERV-K18 superantigens in humans. Blood, 2005, 105, 4377-4382.	1.4	34
85	Ly49 and CD94/NKG2 receptor acquisition by NK cells does not require lymphotoxin-β receptor expression. Blood, 2005, 106, 956-962.	1.4	10
86	Different thresholds of Notch signaling bias human precursor cells toward B-, NK-, monocytic/dendritic-, or T-cell lineage in thymus microenvironment. Blood, 2005, 106, 3498-3506.	1.4	84
87	Tangeretin inhibits extracellular-signal-regulated kinase (ERK) phosphorylation. FEBS Letters, 2005, 579, 1665-1669.	2.8	37
88	Defective CD4+CD25+ regulatory T cell functioning in collagen-induced arthritis: an important factor in pathogenesis, counter-regulated by endogenous IFN-gamma. Arthritis Research, 2005, 7, R402.	2.0	143
89	Role of natural killer cells in the rejection process of corneal allografts in rats. Transplantation, 2004, 77, 676-682.	1.0	27
90	Ly49E expression points toward overlapping, but distinct, natural killer (NK) cell differentiation kinetics and potential of fetal versus adult lymphoid progenitors. Journal of Leukocyte Biology, 2003, 73, 731-738.	3.3	20

#	Article	IF	CITATIONS
91	Active Form of Notch Imposes T Cell Fate in Human Progenitor Cells. Journal of Immunology, 2002, 169, 3021-3029.	0.8	100
92	Developmental and Functional Defects of Thymic and Epidermal Vγ3 Cells in IL-15-Deficient and IFN Regulatory Factor-1-Deficient Mice. Journal of Immunology, 2002, 168, 6486-6493.	0.8	76
93	Expression of Inhibitory Receptors Ly49E and CD94/NKG2 on Fetal Thymic and Adult Epidermal TCR Vγ3 Lymphocytes. Journal of Immunology, 2002, 168, 3295-3302.	0.8	54
94	HOX-A10 regulates hematopoietic lineage commitment: evidence for a monocyte-specific transcription factor. Blood, 2002, 99, 1197-1204.	1.4	64
95	NK cells developing in vitro from fetal mouse progenitors express at least one member of the Ly49 family that is acquired in a time-dependent and stochastic manner independently of CD94 and NKG2. European Journal of Immunology, 2002, 32, 868.	2.9	34
96	The Citrus Methoxyflavone Tangeretin Affects Human Cell-Cell Interactions. Advances in Experimental Medicine and Biology, 2002, 505, 135-139.	1.6	24
97	Expression of Ly49E and CD94/NKG2 on Fetal and Adult NK Cells. Journal of Immunology, 2001, 166, 4302-4311.	0.8	58
98	Enforced Expression of GATA-3 Severely Reduces Human Thymic Cellularity. Journal of Immunology, 2001, 167, 4468-4475.	0.8	37
99	Human B Cell Growth and Differentiation in the Spleen of Immunodeficient Mice. Journal of Immunology, 2001, 166, 2929-2936.	0.8	31
100	Langerhans Cells That Have Matured In Vivo in the Absence of T Cells Are Fully Capable of Inducing a Helper CD4 as Well as a Cytotoxic CD8 Response. Journal of Immunology, 2000, 165, 645-653.	0.8	19
101	Signals from the IL-9 Receptor Are Critical for the Early Stages of Human Intrathymic T Cell Development. Journal of Immunology, 2000, 164, 1761-1767.	0.8	31
102	Murine fetal natural killer cells are functionally and structurally distinct from adult natural killer cells. Journal of Leukocyte Biology, 1999, 66, 625-633.	3.3	17
103	Cytokine dependence of Vγ 3 thymocytes: mature but not immature Vγ3 cells require endogenous IL-2 and IL-7 to survive— evidence for cytokine redundancy. International Immunology, 1995, 7, 843-851.	4.0	20
104	Expression and Function of Fc Receptors in the Thymus. Critical Reviews in Immunology, 1995, 15, 215-233.	0.5	2
105	Human Fetal Liver Cells Differentiate Into Thymocytes in Chimeric Mouse Fetal Thymus Organ Culture. Advances in Experimental Medicine and Biology, 1994, 355, 27-31.	1.6	6
106	Presence of CD8α-CD8β-positive TcR γ/δ thymocytes in the fetal murine thymus and theirin vitro expansion with interleukin-7. European Journal of Immunology, 1992, 22, 2189-2193.	2.9	14
107	Interleukin 4 induces CD8α expression on T cell receptor Vγ5 thymocytes. European Journal of Immunology, 1991, 21, 1751-1754.	2.9	4
108	Enhanced oxygen metabolism of peritoneal macrophages in the presence of murine neuroblastoma cells is partly caused by enkephalins. Journal of Neuroimmunology, 1988, 19, 269-278.	2.3	3

#	Article	IF	CITATIONS
109	Immune response to murine neuroblastoma: Effects of in vitro culture of the tumor. Journal of Neuroimmunology, 1987, 15, 111-120.	2.3	3
110	Cellular immunity changes caused by LDH virus: Analogy with observations on neuroblastoma-bearing mice. International Journal of Cancer, 1987, 40, 669-675.	5.1	5