

Colin A B Jahoda

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3239185/publications.pdf>

Version: 2024-02-01

75
papers

5,663
citations

81900

39
h-index

76900

74
g-index

79
all docs

79
docs citations

79
times ranked

5750
citing authors

#	ARTICLE	IF	CITATIONS
1	Dominant effect of gap junction communication in wound-induced calcium wave, NFAT activation and wound closure in keratinocytes. <i>Journal of Cellular Physiology</i> , 2021, 236, 8171-8183.	4.1	6
2	Extracellular Vesicles from Activated Dermal Fibroblasts Stimulate Hair Follicle Growth Through Dermal Papilla-Secreted Norrin. <i>Stem Cells</i> , 2019, 37, 1166-1175.	3.2	44
3	Hair Follicle Dermal Cells Support Expansion of Murine and Human Embryonic and Induced Pluripotent Stem Cells and Promote Haematopoiesis in Mouse Cultures. <i>Stem Cells International</i> , 2018, 2018, 1-14.	2.5	3
4	Peptoid Efficacy against Polymicrobial Biofilms Determined by Using Propidium Monoazide-Modified Quantitative PCR. <i>ChemBioChem</i> , 2017, 18, 111-118.	2.6	31
5	Hierarchical patterning modes orchestrate hair follicle morphogenesis. <i>PLoS Biology</i> , 2017, 15, e2002117.	5.6	109
6	What Lies Beneath: Wnt/ β -Catenin Signaling and Cell Fate in the Lower Dermis. <i>Journal of Investigative Dermatology</i> , 2016, 136, 1084-1087.	0.7	6
7	Giant Panda (<i>Ailuropoda melanoleuca</i>) Buccal Mucosa Tissue as a Source of Multipotent Progenitor Cells. <i>PLoS ONE</i> , 2015, 10, e0138840.	2.5	6
8	Dermal white adipose tissue: a new component of the thermogenic response. <i>Journal of Lipid Research</i> , 2015, 56, 2061-2069.	4.2	104
9	Type VII collagen regulates tumour expression of organic anion transporting polypeptide OATP1B3, promotes front to rear polarity and increases structural organisation in 3D spheroid cultures of recessive dystrophic epidermolysis bullosa tumour keratinocytes. <i>Journal of Cell Science</i> , 2014, 127, 740-51.	2.0	22
10	Defining dermal adipose tissue. <i>Experimental Dermatology</i> , 2014, 23, 629-631.	2.9	218
11	Microenvironmental reprogramming by three-dimensional culture enables dermal papilla cells to induce de novo human hair-follicle growth. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 19679-19688.	7.1	309
12	Isolation and Establishment of Hair Follicle Dermal Papilla Cell Cultures. <i>Methods in Molecular Biology</i> , 2013, 989, 285-292.	0.9	34
13	Development of the Mouse Dermal Adipose Layer Occurs Independently of Subcutaneous Adipose Tissue and Is Marked by Restricted Early Expression of FABP4. <i>PLoS ONE</i> , 2013, 8, e59811.	2.5	66
14	Reprogramming of Human Hair Follicle Dermal Papilla Cells into Induced Pluripotent Stem Cells. <i>Journal of Investigative Dermatology</i> , 2012, 132, 1725-1727.	0.7	27
15	An <i>in vivo</i> mouse model of human skin replacement for wound healing and cell therapy studies. <i>Journal of Plastic, Reconstructive and Aesthetic Surgery</i> , 2012, 65, 1129-1131.	1.0	8
16	Human hair follicle dermal cells and skin fibroblasts show differential activation of NF- κ B in response to pro-inflammatory challenge. <i>Experimental Dermatology</i> , 2012, 21, 158-160.	2.9	12
17	Generation and Characterization of Multipotent Stem Cells from Established Dermal Cultures. <i>PLoS ONE</i> , 2012, 7, e50742.	2.5	42
18	Niche Crosstalk: Intercellular Signals at the Hair Follicle. <i>Cell</i> , 2011, 146, 678-681.	28.9	53

#	ARTICLE	IF	CITATIONS
19	Modulation in Proteolytic Activity Is Identified as a Hallmark of Exogen by Transcriptional Profiling of Hair Follicles. <i>Journal of Investigative Dermatology</i> , 2011, 131, 2349-2357.	0.7	15
20	Modelling the hair follicle dermal papilla using spheroid cell cultures. <i>Experimental Dermatology</i> , 2010, 19, 546-548.	2.9	105
21	Genome-wide association study in alopecia areata implicates both innate and adaptive immunity. <i>Nature</i> , 2010, 466, 113-117.	27.8	651
22	Effects of Physiological Electric Fields on Migration of Human Dermal Fibroblasts. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2320-2327.	0.7	153
23	KGF and EGF signalling block hair follicle induction and promote interfollicular epidermal fate in developing mouse skin. <i>Development (Cambridge)</i> , 2009, 136, 2153-2164.	2.5	78
24	Dynamic complexes of A-type lamins and emerin influence adipogenic capacity of the cell via nucleocytoplasmic distribution of β -catenin. <i>Journal of Cell Science</i> , 2009, 122, 401-413.	2.0	74
25	Dynamic expression of Syndecan-1 during hair follicle morphogenesis. <i>Gene Expression Patterns</i> , 2009, 9, 454-460.	0.8	20
26	From Telogen to Exogen: Mechanisms Underlying Formation and Subsequent Loss of the Hair Club Fiber. <i>Journal of Investigative Dermatology</i> , 2009, 129, 2100-2108.	0.7	81
27	Exogen involves gradual release of the hair club fibre in the vibrissa follicle model. <i>Experimental Dermatology</i> , 2009, 18, 793-795.	2.9	12
28	An improved method of human keratinocyte culture from skin explants: cell expansion is linked to markers of activated progenitor cells. <i>Experimental Dermatology</i> , 2009, 18, 720-726.	2.9	27
29	Multipotent skin-derived precursors: from biology to clinical translation. <i>Current Opinion in Biotechnology</i> , 2009, 20, 522-530.	6.6	61
30	Keratin 10 (K10) is expressed suprabasally throughout the limbus of embryonic and neonatal rat corneas, with interrupted expression in the adult limbus. <i>Experimental Eye Research</i> , 2009, 89, 435-438.	2.6	7
31	Id2, Id3, and Id4 proteins show dynamic changes in expression during vibrissae follicle development. <i>Developmental Dynamics</i> , 2008, 237, 1653-1661.	1.8	7
32	A Highly Enriched Niche of Precursor Cells with Neuronal and Glial Potential Within the Hair Follicle Dermal Papilla of Adult Skin. <i>Stem Cells</i> , 2008, 26, 163-172.	3.2	198
33	C/EBP β identifies differentiating preadipocytes around hair follicles in foetal and neonatal rat and mouse skin. <i>Experimental Dermatology</i> , 2008, 17, 675-680.	2.9	24
34	Dynamic expression of the zinc-finger transcription factor Trps1 during hair follicle morphogenesis and cycling. <i>Gene Expression Patterns</i> , 2008, 8, 51-57.	0.8	42
35	The MAGUK-family protein CASK is targeted to nuclei of the basal epidermis and controls keratinocyte proliferation. <i>Journal of Cell Science</i> , 2008, 121, 2705-2717.	2.0	21
36	Hair follicle stem cells. <i>Seminars in Cell and Developmental Biology</i> , 2007, 18, 245-254.	5.0	75

#	ARTICLE	IF	CITATIONS
37	The Wnt inhibitor, Dickkopf 4, is induced by canonical Wnt signaling during ectodermal appendage morphogenesis. <i>Developmental Biology</i> , 2007, 305, 498-507.	2.0	99
38	Transcriptional profiling of developing mouse epidermis reveals novel patterns of coordinated gene expression. <i>Developmental Dynamics</i> , 2007, 236, 961-970.	1.8	54
39	Cultured Cells from the Adult Human Hair Follicle Dermis can be Directed Toward Adipogenic and Osteogenic Differentiation. <i>Journal of Investigative Dermatology</i> , 2005, 124, 1090-1091.	0.7	32
40	Plasticity of Rodent and Human Hair Follicle Dermal Cells: Implications for Cell Therapy and Tissue Engineering. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2005, 10, 180-183.	0.8	59
41	The role of BMP signalling in the control of ID3 expression in the hair follicle. <i>Experimental Dermatology</i> , 2004, 13, 621-629.	2.9	20
42	The WNT Signalling Modulator, Wise, is Expressed in an Interaction-Dependent Manner During Hair-Follicle Cycling. <i>Journal of Investigative Dermatology</i> , 2004, 123, 613-621.	0.7	32
43	Transformation of amnion epithelium into skin and hair follicles. <i>Differentiation</i> , 2004, 72, 558-565.	1.9	35
44	Cultured human and rat tooth papilla cells induce hair follicle regeneration and fiber growth. <i>Differentiation</i> , 2004, 72, 566-575.	1.9	49
45	Total Deletion of <i>in Vivo</i> Telomere Elongation Capacity: An Ambitious but Possibly Ultimate Cure for All Age-Related Human Cancers. <i>Annals of the New York Academy of Sciences</i> , 2004, 1019, 147-170.	3.8	20
46	Genomic organization and analysis of the hairless gene in four hypotrichotic rat strains. <i>Mammalian Genome</i> , 2004, 15, 975-981.	2.2	6
47	The lanceolate hair rat phenotype results from a missense mutation in a calcium coordinating site of the desmoglein 4 gene. <i>Genomics</i> , 2004, 83, 747-756.	2.9	49
48	Hair Follicle Stem Cells. <i>Journal of Investigative Dermatology Symposium Proceedings</i> , 2003, 8, 28-38.	0.8	133
49	Cell Movement in the Hair Follicle Dermis – More Than a Two-Way Street?. <i>Journal of Investigative Dermatology</i> , 2003, 121, ix-xi.	0.7	45
50	Hair follicle dermal cells differentiate into adipogenic and osteogenic lineages. <i>Experimental Dermatology</i> , 2003, 12, 849-859.	2.9	246
51	Desmoglein 4 in Hair Follicle Differentiation and Epidermal Adhesion. <i>Cell</i> , 2003, 113, 249-260.	28.9	301
52	Hair follicle dermal cells repopulate the mouse haematopoietic system. <i>Journal of Cell Science</i> , 2002, 115, 3967-3974.	2.0	165
53	Genes that are differentially expressed in rat vibrissa follicle germinative epithelium in vivo show altered expression patterns after extended organ culture. <i>Experimental Dermatology</i> , 2002, 11, 542-555.	2.9	3
54	Defolliculated (Dfl): A Dominant Mouse Mutation Leading to Poor Sebaceous Gland Differentiation and Total Elimination of Pelage Follicles. <i>Journal of Investigative Dermatology</i> , 2002, 119, 32-37.	0.7	50

#	ARTICLE	IF	CITATIONS
55	A correlation between versican and neurofilament expression patterns during the development and adult cycling of rat vibrissa follicles. <i>Mechanisms of Development</i> , 2001, 101, 227-231.	1.7	5
56	Hair follicle dermal sheath cells: unsung participants in wound healing. <i>Lancet, The</i> , 2001, 358, 1445-1448.	13.7	224
57	Emulsion-Derived Foams (PolyHIPEs) Containing Poly(μ -caprolactone) as Matrixes for Tissue Engineering. <i>Biomacromolecules</i> , 2001, 2, 154-164.	5.4	199
58	In Vivo Induction of Hair Growth by Dermal Cells Isolated from Hair Follicles After Extended Organ Culture. <i>Journal of Investigative Dermatology</i> , 2001, 117, 596-604.	0.7	19
59	Hair follicle predetermination. <i>Journal of Cell Science</i> , 2001, 114, 3419-3431.	2.0	195
60	Localisation of members of the notch system and the differentiation of vibrissa hair follicles: Receptors, ligands, and fringe modulators. <i>Developmental Dynamics</i> , 2000, 218, 426-437.	1.8	55
61	Skin stem cells "a hairy issue.. <i>Nature Medicine</i> , 2000, 6, 1095-1097.	30.7	23
62	Trans-gender induction of hair follicles. <i>Nature</i> , 1999, 402, 33-34.	27.8	235
63	Hair Cycle Stage of the Mouse Vibrissa Follicle Determines Subsequent Fiber Growth and Follicle Behavior In Vitro. <i>Journal of Investigative Dermatology</i> , 1997, 108, 495-500.	0.7	29
64	DERMAL-EPIDERMAL INTERACTIONS. <i>Dermatologic Clinics</i> , 1996, 14, 573-583.	1.7	109
65	Human Hair Follicle Regeneration Following Amputation and Grafting into the Nude Mouse. <i>Journal of Investigative Dermatology</i> , 1996, 107, 804-807.	0.7	73
66	Hair Follicle and Fiber Reconstruction. <i>Journal of Investigative Dermatology</i> , 1995, 104, 39-40.	0.7	4
67	Hair follicle reconstruction in vitro. <i>Journal of Dermatological Science</i> , 1994, 7, S84-S97.	1.9	12
68	Hair Follicle Stem Cells: Characteristics and Possible Significance. <i>Skin Pharmacology and Physiology</i> , 1994, 7, 16-19.	2.5	11
69	A Quantitative Study of the Differential Expression of Alpha-Smooth Muscle Actin in Cell Populations of Follicular and Non-Follicular Origin. <i>Journal of Investigative Dermatology</i> , 1993, 101, 577-583.	0.7	29
70	Induction of Hair Growth in Ear Wounds by Cultured Dermal Papilla Cells. <i>Journal of Investigative Dermatology</i> , 1993, 101, 584-590.	0.7	107
71	Human Hair Follicle Germinative Epidermal Cell Culture. <i>Journal of Investigative Dermatology</i> , 1993, 101, 634-638.	0.7	29
72	The DEBR Rat Model for Alopecia Areata. <i>Journal of Investigative Dermatology</i> , 1991, 96, S97.	0.7	18

#	ARTICLE	IF	CITATIONS
73	Inductive Properties of Hair Follicle Cells. Annals of the New York Academy of Sciences, 1991, 642, 226-241.	3.8	68
74	Dermal-epidermal interactions. Clinics in Dermatology, 1988, 6, 74-82.	1.6	64
75	Histochemical localization of skin glycosaminoglycans during feather development in the chick embryo. Roux's Archives of Developmental Biology, 1987, 196, 303-315.	1.2	6