Cleusa P Ferri

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3264947/publications.pdf

Version: 2024-02-01

147 papers 38,002 citations

41344 49 h-index 145 g-index

156 all docs

156 docs citations

156 times ranked 57398 citing authors

#	Article	IF	CITATIONS
1	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2197-2223.	13.7	7,061
2	Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2163-2196.	13.7	6,376
3	Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 743-800.	13.7	4,951
4	Global prevalence of dementia: a Delphi consensus study. Lancet, The, 2005, 366, 2112-2117.	13.7	4,496
5	The global prevalence of dementia: A systematic review and metaanalysis. Alzheimer's and Dementia, 2013, 9, 63.	0.8	3,654
6	Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 2287-2323.	13.7	2,184
7	Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 1015-1035.	13.7	2,005
8	Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2129-2143.	13.7	1,013
9	Contribution of chronic diseases to disability in elderly people in countries with low and middle incomes: a 10/66 Dementia Research Group population-based survey. Lancet, The, 2009, 374, 1821-1830.	13.7	370
10	Prevalence of dementia in Latin America, India, and China: a population-based cross-sectional survey. Lancet, The, 2008, 372, 464-474.	13.7	358
11	The protocols for the 10/66 dementia research group population-based research programme. BMC Public Health, 2007, 7, 165.	2.9	298
12	Burden of disease in Brazil, 1990–2016: a systematic subnational analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 760-775.	13.7	267
13	Dementia incidence and mortality in middle-income countries, and associations with indicators of cognitive reserve: a 10/66 Dementia Research Group population-based cohort study. Lancet, The, 2012, 380, 50-58.	13.7	254
14	Prevalence, Distribution, and Impact of Mild Cognitive Impairment in Latin America, China, and India: A 10/66 Population-Based Study. PLoS Medicine, 2012, 9, e1001170.	8.4	143
15	Prevalence of stroke and related burden among older people living in Latin America, India and China. Journal of Neurology, Neurosurgery and Psychiatry, 2011, 82, 1074-1082.	1.9	137
16	The Impact of Epidemic Violence on the Prevalence of Psychiatric Disorders in Sao Paulo and Rio de Janeiro, Brazil. PLoS ONE, 2013, 8, e63545.	2.5	109
17	The 10/66 Dementia Research Group's fully operationalised DSM-IV dementia computerized diagnostic algorithm, compared with the 10/66 dementia algorithm and a clinician diagnosis: a population validation study. BMC Public Health, 2008, 8, 219.	2.9	108
18	Frailty and the prediction of dependence and mortality in low- and middle-income countries: a 10/66 population-based cohort study. BMC Medicine, 2015, 13, 138.	5.5	101

#	Article	IF	CITATIONS
19	Prevalence of anxiety and its correlates among older adults in Latin America, India and China: cross-cultural study. British Journal of Psychiatry, 2011, 199, 485-491.	2.8	97
20	A comparative cross-cultural study of the prevalence of late life depression in low and middle income countries. Journal of Affective Disorders, 2016, 190, 362-368.	4.1	96
21	"The contribution of chronic diseases to the prevalence of dependence among older people in Latin America, China and India: a 10/66 Dementia Research Group population-based survey". BMC Geriatrics, 2010, 10, 53.	2.7	94
22	Measuring disability across cultures — the psychometric properties of the WHODAS II in older people from seven lowâ€and middleâ€income countries. The 10/66 Dementia Research Group populationâ€based survey. International Journal of Methods in Psychiatric Research, 2010, 19, 1-17.	2.1	88
23	Behavioral and Psychological Symptoms of Dementia in developing countries. International Psychogeriatrics, 2004, 16, 441-459.	1.0	87
24	Determinants of cognitive performance and decline in 20 diverse ethno-regional groups: A COSMIC collaboration cohort study. PLoS Medicine, 2019, 16, e1002853.	8.4	86
25	Population normative data for the 10/66 Dementia Research Group cognitive test battery from Latin America, India and China: a cross-sectional survey. BMC Neurology, 2009, 9, 48.	1.8	85
26	Helping carers to careâ€"The 10/66 dementia research group's randomized control trial of a caregiver intervention in Russia. International Journal of Geriatric Psychiatry, 2009, 24, 347-354.	2.7	85
27	A brief dementia screener suitable for use by nonâ€specialists in resource poor settingsâ€"the crossâ€cultural derivation and validation of the brief Community Screening Instrument for Dementia. International Journal of Geriatric Psychiatry, 2011, 26, 899-907.	2.7	85
28	The impact of maternal experience of violence and common mental disorders on neonatal outcomes: a survey of adolescent mothers in Sao Paulo, Brazil. BMC Public Health, 2007, 7, 209.	2.9	84
29	Dietary fish and meat intake and dementia in Latin America, China, and India: a 10/66 Dementia Research Group population-based study. American Journal of Clinical Nutrition, 2009, 90, 392-400.	4.7	84
30	Hypertension prevalence, awareness, treatment and control among older people in Latin America, India and China. Journal of Hypertension, 2012, 30, 177-187.	0.5	83
31	Asthma and Suicide Mortality in Young People: A 12-Year Follow-Up Study. American Journal of Psychiatry, 2010, 167, 1092-1099.	7.2	82
32	Remembering the forgotten non-communicable diseases. BMC Medicine, 2014, 12, 200.	5.5	82
33	Chronic disease prevalence and care among the elderly in urban and rural Beijing, China - a 10/66 Dementia Research Group cross-sectional survey. BMC Public Health, 2009, 9, 394.	2.9	81
34	Psychometric properties of EURO-D, a geriatric depression scale: a cross-cultural validation study. BMC Psychiatry, 2015, 15, 12.	2.6	81
35	Dementia in low-income and middle-income countries: Different realities mandate tailored solutions. PLoS Medicine, 2017, 14, e1002271.	8.4	81
36	Equity in the delivery of community healthcare to older people: findings from 10/66 Dementia Research Group cross-sectional surveys in Latin America, China, India and Nigeria. BMC Health Services Research, 2011, 11, 153.	2.2	80

#	Article	IF	Citations
37	Strain and its correlates among carers of people with dementia in lowâ€income and middleâ€income countries. A 10/66 Dementia Research Group populationâ€based survey. International Journal of Geriatric Psychiatry, 2012, 27, 670-682.	2.7	80
38	Late-life depression in Peru, Mexico and Venezuela: the 10/66 population-based study. British Journal of Psychiatry, 2009, 195, 510-515.	2.8	79
39	Rate of cognitive decline in relation to sex after 60 yearsâ€ofâ€age: A systematic review. Geriatrics and Gerontology International, 2014, 14, 23-31.	1.5	75
40	Use of licit and illicit substances among adolescents in Brazil $\hat{a} \in$ "A national survey. Addictive Behaviors, 2012, 37, 1171-1175.	3.0	74
41	Prevalence and correlates for sleep complaints in older adults in low and middle income countries: A 10/66 Dementia Research Group study. Sleep Medicine, 2012, 13, 697-702.	1.6	73
42	Crack Cocaine: An Increase in Use among Patients Attending Clinics in Sa \hat{A} Paulo: 1990-1993. Substance Use and Misuse, 1996, 31, 519-527.	1.4	67
43	Unawareness of memory impairment in dementia: a population-based study. International Psychogeriatrics, 2012, 24, 931-939.	1.0	66
44	Ageing and dementia in low and middle income countries–Using research to engage with public and policy makers. International Review of Psychiatry, 2008, 20, 332-343.	2.8	62
45	Validity and reliability of the Severity of Dependence Scale (SDS) in a Brazilian sample of drug users. Drug and Alcohol Review, 2000, 19, 451-455.	2.1	59
46	Dementia underdiagnosis in Brazil. Lancet, The, 2015, 385, 418-419.	13.7	58
47	Co-occurrence of anxiety and depression amongst older adults in low- and middle-income countries: findings from the 10/66 study. Psychological Medicine, 2011, 41, 2047-2056.	4.5	53
48	Socioeconomic Factors and All Cause and Cause-Specific Mortality among Older People in Latin America, India, and China: A Population-Based Cohort Study. PLoS Medicine, 2012, 9, e1001179.	8.4	52
49	Helping carers to care: the 10/66 dementia research group's randomized control trial of a caregiver intervention in Peru. Revista Brasileira De Psiquiatria, 2011, 33, 47-54.	1.7	51
50	Route of cocaine administration. Addictive Behaviors, 1999, 24, 815-821.	3.0	50
51	The Epidemiology of Dependence in Older People in Nigeria: Prevalence, Determinants, Informal Care, and Health Service Utilization. A 10/66 Dementia Research Group Crossâ€Sectional Survey. Journal of the American Geriatrics Society, 2009, 57, 1620-1627.	2.6	50
52	Association between sleep duration and all-cause mortality in old age: 9-year follow-up of the BambuÃ-Cohort Study, Brazil. Journal of Sleep Research, 2011, 20, 303-310.	3.2	50
53	Reducing the Risk of Cognitive Decline and Dementia: WHO Recommendations. Frontiers in Neurology, 2021, 12, 765584.	2.4	42
54	Violence during pregnancy and newborn outcomes: a cohort study in a disadvantaged population in Brazil. European Journal of Public Health, 2011, 21, 92-97.	0.3	41

#	Article	IF	Citations
55	The epidemiology of dependency among urban-dwelling older people in the Dominican Republic; a cross-sectional survey. BMC Public Health, 2008, 8, 285.	2.9	40
56	Violence and post-traumatic stress disorder in Sao Paulo and Rio de Janeiro, Brazil: the protocol for an epidemiological and genetic survey. BMC Psychiatry, 2009, 9, 34.	2.6	38
57	Suicide attempts prior to starting methadone maintenance treatment in Taiwan. Drug and Alcohol Dependence, 2010, 109, 139-143.	3.2	38
58	Severity of heroin dependence in Taiwan: Reliability and validity of the Chinese version of the Severity of Dependence Scale (SDS[Ch]). Addictive Behaviors, 2008, 33, 1590-1593.	3.0	37
59	Alcohol consumption in late-life $\hat{a}\in$ " The first Brazilian National Alcohol Survey (BNAS). Addictive Behaviors, 2008, 33, 1598-1601.	3.0	37
60	The Prevalence, Correlates and Impact of Dementia in Cuba. Neuroepidemiology, 2008, 31, 243-251.	2.3	37
61	A Population-Based Study of the Association between <i>Trypanosoma cruzi</i> Infection and Cognitive Impairment in Old Age (The BambuÃ-Study). Neuroepidemiology, 2009, 32, 122-128.	2.3	37
62	Nutrition, mental health and violence: from pregnancy to postpartum Cohort of women attending primary care units in Southern Brazil - ECCAGE study. BMC Psychiatry, 2010, 10, 66.	2.6	34
63	The association between common physical impairments and dementia in low and middle income countries, and, among people with dementia, their association with cognitive function and disability. A 10/66 Dementia Research Group population-based study. International Journal of Geriatric Psychiatry. 2011. 26. 511-519.	2.7	33
64	Early life exposure to violence and substance misuse in adulthoodâ€"The first Brazilian national survey. Addictive Behaviors, 2011, 36, 251-255.	3.0	32
65	Midlife Hypertensive Status and Cognitive Function 20ÂYears Later: The <scp>S</scp> outhall and <scp>B</scp> rent Revisited Study. Journal of the American Geriatrics Society, 2013, 61, 1489-1498.	2.6	32
66	An investigation of associations between alcohol use disorder and polymorphisms on <i>ALDH2</i> , <i>BDNF</i> , <i>5â€HTTLPR</i> , and <i>MTHFR</i> genes in older Korean men. International Journal of Geriatric Psychiatry, 2010, 25, 441-448.	2.7	31
67	Resource utilisation for neuropsychiatric disorders in developing countries:. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 218-227.	3.1	30
68	Dimensions Underlying the Mini-Mental State Examination in a Sample With Low-Education Levels: The BambuÃ-Health and Aging Study. American Journal of Geriatric Psychiatry, 2009, 17, 863-872.	1.2	29
69	The Prevalence, Correlates, Detection and Control of Diabetes among Older People in Low and Middle Income Countries. A 10/66 Dementia Research Group Population-Based Survey. PLoS ONE, 2016, 11, e0149616.	2.5	27
70	Reducing the Number of People with Dementia Through Primary Prevention in Mozambique, Brazil, and Portugal: An Analysis of Population-Based Data. Journal of Alzheimer's Disease, 2019, 70, S283-S291.	2.6	27
71	Interactions between genetic admixture, ethnic identity, APOE genotype and dementia prevalence in an admixed Cuban sample; a cross-sectional population survey and nested case-control study. BMC Medical Genetics, 2011, 12, 43.	2.1	25
72	Epidemiological methods for research with drug misusers: review of methods for studying prevalence and morbidity. Revista De Saude Publica, 1999, 33, 206-215.	1.7	24

#	Article	IF	Citations
73	Nonprescribed use of tranquilizers or sedatives by adolescents: a Brazilian national survey. BMC Public Health, 2013, 13, 499.	2.9	24
74	The prevalence of dementia in a Portuguese community sample: a 10/66 Dementia Research Group study. BMC Geriatrics, 2017, 17, 261.	2.7	24
75	Suicide and Other-Cause Mortality after Early Exposure to Smoking and Second Hand Smoking: A 12-Year Population-Based Follow-Up Study. PLoS ONE, 2015, 10, e0130044.	2.5	23
76	Pragmatic randomized controlled trial of a web-based intervention for alcohol use among Brazilian college students: Motivation as a moderating effect. Drug and Alcohol Dependence, 2019, 199, 92-100.	3.2	23
77	No Association between Fish Intake and Depression in over 15,000 Older Adults from Seven Low and Middle Income Countries–The 10/66 Study. PLoS ONE, 2012, 7, e38879.	2.5	21
78	A systematic review and meta-analysis of dementia prevalence in seven developing countries: A STRiDE project. Global Public Health, 2020, 15, 1878-1893.	2.0	21
79	Factors associated with adverse reactions to cocaine among a sample of long-term, high-dose users in São Paulo, Brazil. Addictive Behaviors, 2004, 29, 365-374.	3.0	20
80	HIGH DOSE COCAINE USE IN SÃ f O PAULO: A COMPARISON OF TREATMENT AND COMMUNITY SAMPLES. Substance Use and Misuse, 2001, 36, 237-255.	1.4	19
81	Chronic diseases among older people and co-resident psychological morbidity: a 10/66 Dementia Research Group population-based survey. International Psychogeriatrics, 2011, 23, 1489-1501.	1.0	19
82	Tobacco smoking and risk for dementia: evidence from the 10/66 population-based longitudinal study. Aging and Mental Health, 2020, 24, 1796-1806.	2.8	18
83	Frailty Modifies the Association of Hypertension With Cognition in Older Adults: Evidence From the ELSI-Brazil. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 1134-1143.	3.6	18
84	Violence and depressive symptoms during pregnancy: a primary care study in Brazil. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 983-988.	3.1	17
85	The prevalence and social patterning of chronic diseases among older people in a population undergoing health transition. A 10/66 Group cross-sectional population-based survey in the Dominican Republic. BMC Public Health, 2010, 10, 344.	2.9	17
86	Tobacco use and dementia: evidence from the 1066 dementia populationâ€based surveys in Latin America, China and India. International Journal of Geriatric Psychiatry, 2011, 26, 1177-1185.	2.7	16
87	Contributions of parenting styles and parental drunkenness to adolescent drinking. Revista Brasileira De Psiquiatria, 2019, 41, 511-517.	1.7	16
88	Betel Chewing and Arecoline Affects Eotaxin-1, Asthma and Lung Function. PLoS ONE, 2014, 9, e91889.	2.5	16
89	Mixed methods implementation research of cognitive stimulation therapy (CST) for dementia in low and middle-income countries: study protocol for Brazil, India and Tanzania (CST-International). BMJ Open, 2019, 9, e030933.	1.9	15
90	Differences in factors associated with first treatment entry and treatment reâ€entry among cocaine users. Addiction, 2002, 97, 825-832.	3.3	14

#	Article	IF	Citations
91	Mental and personality disorders and abstinence from alcohol: results from a national household survey. Psychological Medicine, 2011, 41, 809-818.	4.5	14
92	Psychosocial factors associated with mother–child violence: a household survey. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 77-86.	3.1	14
93	Cognitive function among older adults. Revista De Saude Publica, 2018, 52, 4s.	1.7	13
94	Mortality Rates and Mortality Risk Factors in Older Adults with Dementia from Low- and Middle-Income Countries: The 10/66 Dementia Research Group Population-Based Cohort Study. Journal of Alzheimer's Disease, 2020, 75, 581-593.	2.6	13
95	Does multisite sampling improve patient heterogeneity in drug misuse research?. Drug and Alcohol Dependence, 2001, 63, 79-85.	3.2	12
96	The psychological impact of heavy drinking among the elderly on their co-residents: The 10/66 group population based survey in the Dominican Republic. Drug and Alcohol Dependence, 2011, 114, 82-86.	3.2	12
97	Leg length, skull circumference, and the prevalence of dementia in low and middle income countries: a 10/66 population-based cross sectional survey. International Psychogeriatrics, 2011, 23, 202-213.	1.0	12
98	Translating science into policy: mental health challenges during the COVID-19 pandemic. Revista Brasileira De Psiquiatria, 2021, 43, 638-649.	1.7	12
99	Delivering Cognitive Stimulation Therapy (CST) Virtually: Developing and Field-Testing a New Framework. Clinical Interventions in Aging, 2022, Volume 17, 97-116.	2.9	12
100	Letter to the Editor. Addiction, 1998, 93, 287-288.	3.3	11
101	Underdetection of Psychiatric Disorders During Prenatal Care: A Survey of Adolescents in Sao Paulo, Brazil. Journal of Adolescent Health, 2012, 50, 93-96.	2.5	11
102	Patterns of domestic violence and alcohol consumption among women and the effectiveness of a brief intervention in a household setting:Âa protocol study. BMC Women's Health, 2015, 15, 78.	2.0	11
103	Neuropsychological and Behavioral Disturbance Correlates of Unawareness of Memory Impairment in Dementia. Journal of Geriatric Psychiatry and Neurology, 2015, 28, 3-11.	2.3	11
104	Alcohol dependence criteria in DSM-III-R: presence of symptoms according to degree of severity. Addiction, 1994, 89, 1129-1134.	3.3	10
105	Population Ageing in Latin America: Dementia and Related Disorders. Revista Brasileira De Psiquiatria, 2012, 34, 371-374.	1.7	10
106	Abuse Against Women, Depression, and Infant Morbidity. American Journal of Preventive Medicine, 2012, 43, 188-195.	3.0	10
107	Beyond the project: Building a strategic theory of change to address dementia care, treatment and support gaps across seven middle-income countries. Dementia, 2022, 21, 114-135.	2.0	10
108	Depression increases in patients with Parkinson?s disease according to the increasing severity of the cognitive impairment. Arquivos De Neuro-Psiquiatria, 2014, 72, 426-429.	0.8	9

#	Article	IF	CITATIONS
109	Is there evidence for recommending specific intravenous immunoglobulin formulations? A systematic review of head-to-head randomized controlled trials. European Journal of Pharmacology, 2015, 747, 96-104.	3.5	9
110	Endovascular thrombectomy for the treatment of acute ischemic stroke. Arquivos De Neuro-Psiquiatria, 2016, 74, 67-74.	0.8	9
111	Tobacco use among the elderly: The first Brazilian National Survey (BNAS). Aging and Mental Health, 2010, 14, 720-724.	2.8	8
112	Incidence of dementia and association with APOE genotype in older Cubans. Dementia E Neuropsychologia, 2014, 8, 356-363.	0.8	8
113	Intermittent preventive treatment of malaria during Pathways to homelessness among older people in Rio de Janeiro, Brazil. Bulletin of the World Health Organization, 2007, 85, 888-889.	3.3	7
114	Correlated Factors and Prevalence of Alcohol Treatment in Brazil. Addictive Disorders and Their Treatment, 2015, 14, 40-46.	0.5	7
115	Alcohol and tobacco consumption concordance and its correlates in older couples in Latin America. Geriatrics and Gerontology International, 2017, 17, 1849-1857.	1.5	7
116	Alcohol Consumption by Older People in Brazil: A Systematic Review of Population-based Studies. Addictive Disorders and Their Treatment, 2019, 18, 229-237.	0.5	7
117	Validation of the 10/66 Dementia Research Group's 10/66 Dementia diagnosis in Iran. International Psychogeriatrics, 2009, 21, 604.	1.0	6
118	10/66 Dementia Research Group: recently published survey data for seven Latin America sites. International Psychogeriatrics, 2010, 22, 158-159.	1.0	6
119	Rational use of diagnostic tests for clinical decision making. Revista Da Associação Médica Brasileira, 2019, 65, 452-459.	0.7	6
120	Genetics of cognitive trajectory in Brazilians: 15 years of follow-up from the BambuÃ-Epigen Cohort Study of Aging. Scientific Reports, 2019, 9, 18085.	3.3	6
121	Experiences of stigma and discrimination among people living with dementia and family carers in Brazil: qualitative study. Ageing and Society, 0, , 1-22.	1.7	6
122	Harmonization of epidemiological studies on dementia in Latin America Why does it matter?. Dementia E Neuropsychologia, 2019, 13, 363-366.	0.8	6
123	The prevalence, correlates and impact of anaemia among older people in Cuba, Dominican Republic, Mexico, Puerto Rico and Venezuela. British Journal of Haematology, 2013, 160, 387-398.	2.5	5
124	Description of the crossâ€cultural process adopted in the STRiDE (STrengthening Responses to dementia) Tj ETQo Assessment and Disease Monitoring, 2022, 14, e12293.	q0 0 0 rgB 2.4	T /Overlock 5
125	Prevalence, correlates, and risk perception of HIV infection among heroin users in Central Taiwan. Kaohsiung Journal of Medical Sciences, 2013, 29, 673-679.	1.9	4
126	Nonprescribed use of tranquilizers and use of other drugs among Brazilian students. Revista Brasileira De Psiquiatria, 2014, 36, 16-23.	1.7	4

#	Article	IF	CITATIONS
127	Implementação em Portugal de um estudo de prevalência da demência e da depressão geriátrica: a metodologia do 10/66 Dementia Research Group. Revista Portuguesa De Saude Publica, 2016, 34, 134-143.	0.3	4
128	Incidence of hospitalâ€acquired pressure injury: A cohort study of adults admitted to public and private hospitals in Sao Paulo, Brazil. Wound Repair and Regeneration, 2021, 29, 79-86.	3.0	4
129	Alcohol consumption among older adults: Findings from the ELSIâ€Brazil study. International Journal of Geriatric Psychiatry, 2021, , .	2.7	4
130	Development and acceptability of a brief, evidence-based Dementia Awareness for Caregivers course in low- and middle-income countries. Dementia, 2022, 21, 598-617.	2.0	4
131	P2-171 Pain among older people and its impact on disability: a 10/66 cross-sectional population-based surveys in Latin America, India and China. Journal of Epidemiology and Community Health, 2011, 65, A268-A268.	3.7	2
132	P2-37 Association between sleep duration and all-cause mortality in old age: 9-year follow-up of the Bambui Cohort Study, Brazil. Journal of Epidemiology and Community Health, 2011, 65, A229-A230.	3.7	2
133	Genomic African and Native American Ancestry and 15‥ear Cognitive Trajectory: Bambui Study, Brazil. Journal of the American Geriatrics Society, 2018, 66, 1956-1962.	2.6	2
134	Alcohol and ageing: rapid changes in populations present new challenges for an old problem. Substance Use and Misuse, 2019, 54, 1580-1581.	1.4	2
135	Violence between Different Members in the Family System: A Household Survey in a Brazilian City. Journal of Family Issues, 2020, 41, 1237-1257.	1.6	2
136	Brief interventions for older adults (BIO) delivered by non-specialist community health workers to reduce at-risk drinking in primary care: a study protocol for a randomised controlled trial. BMJ Open, 2021, 11, e043918.	1.9	2
137	Burden of dementia in Brazil. Arquivos De Neuro-Psiquiatria, 2020, 78, 755-756.	0.8	2
138	Depression in older people is associated with increased risk of developing dementia. Evidence-Based Mental Health, 2010, 13, 112-112.	4.5	1
139	P2-536 Co-occurrence of anxiety and depression among older adults in low and middle-income countries: findings from the $10/66$ study. Journal of Epidemiology and Community Health, $2011,65$, A369-A369.	3.7	1
140	Potential for dementia prevention in Latin America and Africa based on population-attributable fraction estimates. The Lancet Global Health, 2019, 7, e1324.	6.3	1
141	Attenuated effect of the apolipoprotein E É>4 allele on neuropsychiatric symptoms among African older adults. International Psychogeriatrics, 2021, 33, 213-215.	1.0	1
142	Innovation and achievement for primary care in Brazil: new challenges. BJGP Open, 2017, 1, bjgpopen17X100857.	1.8	1
143	Measures of effect for cross-sectional studies. Revista Brasileira De Psiquiatria, 2008, 30, 399-399.	1.7	1
144	P3â€080: APOE Genotype and the Prevalence and Incidence of Dementia in Admixed Cuban Population: A 10/66 Populationâ€Based Cohort Study. Alzheimer's and Dementia, 2016, 12, P849.	0.8	0

#	Article	IF	CITATIONS
145	Substance Use Among Older Adults: Epidemiological Aspects, Associated Variables, and Organic Risks., 2021,, 71-82.		O
146	Chronic Disease Prevalence and Care Among the Elderly in Urban and Rural Beijing,China â€" A 10/66 Dementia Research Group Cross-Sectional Survey. , 2011, , 13-32.		0
147	Late-life drinking and smoking in primary care users in Brazil. Aging and Mental Health, 2023, 27, 797-802.	2.8	O