
## Cleusa P Ferri

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3264947/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Late-life drinking and smoking in primary care users in Brazil. Aging and Mental Health, 2023, 27, 797-802.	2.8	0
2	Beyond the project: Building a strategic theory of change to address dementia care, treatment and support gaps across seven middle-income countries. Dementia, 2022, 21, 114-135.	2.0	10
3	Delivering Cognitive Stimulation Therapy (CST) Virtually: Developing and Field-Testing a New Framework. Clinical Interventions in Aging, 2022, Volume 17, 97-116.	2.9	12
4	Description of the cross ultural process adopted in the STRiDE (STrengthening Responses to dementia) Tj ETQ Assessment and Disease Monitoring, 2022, 14, e12293.	q0 0 0 rgE 2.4	3T /Overlock 5
5	Development and acceptability of a brief, evidence-based Dementia Awareness for Caregivers course in low- and middle-income countries. Dementia, 2022, 21, 598-617.	2.0	4
6	Incidence of hospitalâ€acquired pressure injury: A cohort study of adults admitted to public and private hospitals in Sao Paulo, Brazil. Wound Repair and Regeneration, 2021, 29, 79-86.	3.0	4
7	Substance Use Among Older Adults: Epidemiological Aspects, Associated Variables, and Organic Risks. , 2021, , 71-82.		0
8	Translating science into policy: mental health challenges during the COVID-19 pandemic. Revista Brasileira De Psiquiatria, 2021, 43, 638-649.	1.7	12
9	Attenuated effect of the apolipoprotein E ɛ4 allele on neuropsychiatric symptoms among African older adults. International Psychogeriatrics, 2021, 33, 213-215.	1.0	1
10	Brief interventions for older adults (BIO) delivered by non-specialist community health workers to reduce at-risk drinking in primary care: a study protocol for a randomised controlled trial. BMJ Open, 2021, 11, e043918.	1.9	2
11	Frailty Modifies the Association of Hypertension With Cognition in Older Adults: Evidence From the ELSI-Brazil. Journals of Gerontology - Series A Biological Sciences and Medical Sciences, 2021, 76, 1134-1143.	3.6	18
12	Alcohol consumption among older adults: Findings from the ELSIâ€Brazil study. International Journal of Geriatric Psychiatry, 2021, , .	2.7	4
13	Reducing the Risk of Cognitive Decline and Dementia: WHO Recommendations. Frontiers in Neurology, 2021, 12, 765584.	2.4	42
14	Tobacco smoking and risk for dementia: evidence from the 10/66 population-based longitudinal study. Aging and Mental Health, 2020, 24, 1796-1806.	2.8	18
15	Violence between Different Members in the Family System: A Household Survey in a Brazilian City. Journal of Family Issues, 2020, 41, 1237-1257.	1.6	2
16	A systematic review and meta-analysis of dementia prevalence in seven developing countries: A STRiDE project. Global Public Health, 2020, 15, 1878-1893.	2.0	21
17	Mortality Rates and Mortality Risk Factors in Older Adults with Dementia from Low- and Middle-Income Countries: The 10/66 Dementia Research Group Population-Based Cohort Study. Journal of Alzheimer's Disease, 2020, 75, 581-593.	2.6	13
18	Burden of dementia in Brazil. Arquivos De Neuro-Psiquiatria, 2020, 78, 755-756.	0.8	2

#	Article	IF	CITATIONS
19	Determinants of cognitive performance and decline in 20 diverse ethno-regional groups: A COSMIC collaboration cohort study. PLoS Medicine, 2019, 16, e1002853.	8.4	86
20	Potential for dementia prevention in Latin America and Africa based on population-attributable fraction estimates. The Lancet Global Health, 2019, 7, e1324.	6.3	1
21	Reducing the Number of People with Dementia Through Primary Prevention in Mozambique, Brazil, and Portugal: An Analysis of Population-Based Data. Journal of Alzheimer's Disease, 2019, 70, S283-S291.	2.6	27
22	Alcohol and ageing: rapid changes in populations present new challenges for an old problem. Substance Use and Misuse, 2019, 54, 1580-1581.	1.4	2
23	Pragmatic randomized controlled trial of a web-based intervention for alcohol use among Brazilian college students: Motivation as a moderating effect. Drug and Alcohol Dependence, 2019, 199, 92-100.	3.2	23
24	Rational use of diagnostic tests for clinical decision making. Revista Da Associação Médica Brasileira, 2019, 65, 452-459.	0.7	6
25	Contributions of parenting styles and parental drunkenness to adolescent drinking. Revista Brasileira De Psiquiatria, 2019, 41, 511-517.	1.7	16
26	Mixed methods implementation research of cognitive stimulation therapy (CST) for dementia in low and middle-income countries: study protocol for Brazil, India and Tanzania (CST-International). BMJ Open, 2019, 9, e030933.	1.9	15
27	Genetics of cognitive trajectory in Brazilians: 15 years of follow-up from the BambuÃ-Epigen Cohort Study of Aging. Scientific Reports, 2019, 9, 18085.	3.3	6
28	Alcohol Consumption by Older People in Brazil: A Systematic Review of Population-based Studies. Addictive Disorders and Their Treatment, 2019, 18, 229-237.	0.5	7
29	Harmonization of epidemiological studies on dementia in Latin America Why does it matter?. Dementia E Neuropsychologia, 2019, 13, 363-366.	0.8	6
30	Cognitive function among older adults. Revista De Saude Publica, 2018, 52, 4s.	1.7	13
31	Genomic African and Native American Ancestry and 15â€Year Cognitive Trajectory: Bambui Study, Brazil. Journal of the American Geriatrics Society, 2018, 66, 1956-1962.	2.6	2
32	Burden of disease in Brazil, 1990–2016: a systematic subnational analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 760-775.	13.7	267
33	Alcohol use and burden for 195 countries and territories, 1990–2016: a systematic analysis for the Global Burden of Disease Study 2016. Lancet, The, 2018, 392, 1015-1035.	13.7	2,005
34	Alcohol and tobacco consumption concordance and its correlates in older couples in Latin America. Geriatrics and Gerontology International, 2017, 17, 1849-1857.	1.5	7
35	Psychosocial factors associated with mother–child violence: a household survey. Social Psychiatry and Psychiatric Epidemiology, 2017, 52, 77-86.	3.1	14
36	The prevalence of dementia in a Portuguese community sample: a 10/66 Dementia Research Group study. BMC Geriatrics, 2017, 17, 261.	2.7	24

#	Article	IF	CITATIONS
37	Dementia in low-income and middle-income countries: Different realities mandate tailored solutions. PLoS Medicine, 2017, 14, e1002271.	8.4	81
38	Innovation and achievement for primary care in Brazil: new challenges. BJGP Open, 2017, 1, bjgpopen17X100857.	1.8	1
39	Endovascular thrombectomy for the treatment of acute ischemic stroke. Arquivos De Neuro-Psiquiatria, 2016, 74, 67-74.	0.8	9
40	The Prevalence, Correlates, Detection and Control of Diabetes among Older People in Low and Middle Income Countries. A 10/66 Dementia Research Group Population-Based Survey. PLoS ONE, 2016, 11, e0149616.	2.5	27
41	P3â€080: APOE Genotype and the Prevalence and Incidence of Dementia in Admixed Cuban Population: A 10/66 Populationâ€Based Cohort Study. Alzheimer's and Dementia, 2016, 12, P849.	0.8	Ο
42	Implementação em Portugal de um estudo de prevalência da demência e da depressão geriátrica: a metodologia do 10/66 Dementia Research Group. Revista Portuguesa De Saude Publica, 2016, 34, 134-143.	0.3	4
43	A comparative cross-cultural study of the prevalence of late life depression in low and middle income countries. Journal of Affective Disorders, 2016, 190, 362-368.	4.1	96
44	Patterns of domestic violence and alcohol consumption among women and the effectiveness of a brief intervention in a household setting:Âa protocol study. BMC Women's Health, 2015, 15, 78.	2.0	11
45	Suicide and Other-Cause Mortality after Early Exposure to Smoking and Second Hand Smoking: A 12-Year Population-Based Follow-Up Study. PLoS ONE, 2015, 10, e0130044.	2.5	23
46	Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 743-800.	13.7	4,951
47	Dementia underdiagnosis in Brazil. Lancet, The, 2015, 385, 418-419.	13.7	58
48	Psychometric properties of EURO-D, a geriatric depression scale: a cross-cultural validation study. BMC Psychiatry, 2015, 15, 12.	2.6	81
49	Correlated Factors and Prevalence of Alcohol Treatment in Brazil. Addictive Disorders and Their Treatment, 2015, 14, 40-46.	0.5	7
50	Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: a systematic analysis for the Global Burden of Disease Study 2013. Lancet, The, 2015, 386, 2287-2323.	13.7	2,184
51	Frailty and the prediction of dependence and mortality in low- and middle-income countries: a 10/66 population-based cohort study. BMC Medicine, 2015, 13, 138.	5.5	101
52	ls there evidence for recommending specific intravenous immunoglobulin formulations? A systematic review of head-to-head randomized controlled trials. European Journal of Pharmacology, 2015, 747, 96-104.	3.5	9
53	Neuropsychological and Behavioral Disturbance Correlates of Unawareness of Memory Impairment in Dementia. Journal of Geriatric Psychiatry and Neurology, 2015, 28, 3-11.	2.3	11
54	Depression increases in patients with Parkinson?s disease according to the increasing severity of the cognitive impairment. Arquivos De Neuro-Psiquiatria, 2014, 72, 426-429.	0.8	9

#	Article	IF	CITATIONS
55	Incidence of dementia and association with APOE genotype in older Cubans. Dementia E Neuropsychologia, 2014, 8, 356-363.	0.8	8
56	Rate of cognitive decline in relation to sex after 60 yearsâ€ofâ€age: A systematic review. Geriatrics and Gerontology International, 2014, 14, 23-31.	1.5	75
57	Remembering the forgotten non-communicable diseases. BMC Medicine, 2014, 12, 200.	5.5	82
58	Nonprescribed use of tranquilizers and use of other drugs among Brazilian students. Revista Brasileira De Psiquiatria, 2014, 36, 16-23.	1.7	4
59	Betel Chewing and Arecoline Affects Eotaxin-1, Asthma and Lung Function. PLoS ONE, 2014, 9, e91889.	2.5	16
60	Nonprescribed use of tranquilizers or sedatives by adolescents: a Brazilian national survey. BMC Public Health, 2013, 13, 499.	2.9	24
61	Prevalence, correlates, and risk perception of HIV infection among heroin users in Central Taiwan. Kaohsiung Journal of Medical Sciences, 2013, 29, 673-679.	1.9	4
62	The prevalence, correlates and impact of anaemia among older people in Cuba, Dominican Republic, Mexico, Puerto Rico and Venezuela. British Journal of Haematology, 2013, 160, 387-398.	2.5	5
63	The global prevalence of dementia: A systematic review and metaanalysis. Alzheimer's and Dementia, 2013, 9, 63.	0.8	3,654
64	Midlife Hypertensive Status and Cognitive Function 20ÂYears Later: The <scp>S</scp> outhall and <scp>B</scp> rent Revisited Study. Journal of the American Geriatrics Society, 2013, 61, 1489-1498.	2.6	32
65	The Impact of Epidemic Violence on the Prevalence of Psychiatric Disorders in Sao Paulo and Rio de Janeiro, Brazil. PLoS ONE, 2013, 8, e63545.	2.5	109
66	Hypertension prevalence, awareness, treatment and control among older people in Latin America, India and China. Journal of Hypertension, 2012, 30, 177-187.	0.5	83
67	Unawareness of memory impairment in dementia: a population-based study. International Psychogeriatrics, 2012, 24, 931-939.	1.0	66
68	Dementia incidence and mortality in middle-income countries, and associations with indicators of cognitive reserve: a 10/66 Dementia Research Group population-based cohort study. Lancet, The, 2012, 380, 50-58.	13.7	254
69	Common values in assessing health outcomes from disease and injury: disability weights measurement study for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2129-2143.	13.7	1,013
70	Disability-adjusted life years (DALYs) for 291 diseases and injuries in 21 regions, 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2197-2223.	13.7	7,061
71	Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet, The, 2012, 380, 2163-2196.	13.7	6,376
72	Population Ageing in Latin America: Dementia and Related Disorders. Revista Brasileira De Psiquiatria, 2012, 34, 371-374.	1.7	10

#	Article	IF	CITATIONS
73	Prevalence and correlates for sleep complaints in older adults in low and middle income countries: A 10/66 Dementia Research Group study. Sleep Medicine, 2012, 13, 697-702.	1.6	73
74	Use of licit and illicit substances among adolescents in Brazil — A national survey. Addictive Behaviors, 2012, 37, 1171-1175.	3.0	74
75	Underdetection of Psychiatric Disorders During Prenatal Care: A Survey of Adolescents in Sao Paulo, Brazil. Journal of Adolescent Health, 2012, 50, 93-96.	2.5	11
76	Abuse Against Women, Depression, and Infant Morbidity. American Journal of Preventive Medicine, 2012, 43, 188-195.	3.0	10
77	Socioeconomic Factors and All Cause and Cause-Specific Mortality among Older People in Latin America, India, and China: A Population-Based Cohort Study. PLoS Medicine, 2012, 9, e1001179.	8.4	52
78	No Association between Fish Intake and Depression in over 15,000 Older Adults from Seven Low and Middle Income Countries–The 10/66 Study. PLoS ONE, 2012, 7, e38879.	2.5	21
79	Strain and its correlates among carers of people with dementia in lowâ€income and middleâ€income countries. A 10/66 Dementia Research Group populationâ€based survey. International Journal of Geriatric Psychiatry, 2012, 27, 670-682.	2.7	80
80	Prevalence, Distribution, and Impact of Mild Cognitive Impairment in Latin America, China, and India: A 10/66 Population-Based Study. PLoS Medicine, 2012, 9, e1001170.	8.4	143
81	Early life exposure to violence and substance misuse in adulthood—The first Brazilian national survey. Addictive Behaviors, 2011, 36, 251-255.	3.0	32
82	The psychological impact of heavy drinking among the elderly on their co-residents: The 10/66 group population based survey in the Dominican Republic. Drug and Alcohol Dependence, 2011, 114, 82-86.	3.2	12
83	Helping carers to care: the 10/66 dementia research group's randomized control trial of a caregiver intervention in Peru. Revista Brasileira De Psiquiatria, 2011, 33, 47-54.	1.7	51
84	P2-171 Pain among older people and its impact on disability: a 10/66 cross-sectional population-based surveys in Latin America, India and China. Journal of Epidemiology and Community Health, 2011, 65, A268-A268.	3.7	2
85	P2-37 Association between sleep duration and all-cause mortality in old age: 9-year follow-up of the Bambui Cohort Study, Brazil. Journal of Epidemiology and Community Health, 2011, 65, A229-A230.	3.7	2
86	P2-536 Co-occurrence of anxiety and depression among older adults in low and middle-income countries: findings from the 10/66 study. Journal of Epidemiology and Community Health, 2011, 65, A369-A369.	3.7	1
87	Leg length, skull circumference, and the prevalence of dementia in low and middle income countries: a 10/66 population-based cross sectional survey. International Psychogeriatrics, 2011, 23, 202-213.	1.0	12
88	Chronic diseases among older people and co-resident psychological morbidity: a 10/66 Dementia Research Group population-based survey. International Psychogeriatrics, 2011, 23, 1489-1501.	1.0	19
89	Association between sleep duration and all-cause mortality in old age: 9-year follow-up of the BambuÃ- Cohort Study, Brazil. Journal of Sleep Research, 2011, 20, 303-310.	3.2	50
90	Interactions between genetic admixture, ethnic identity, APOE genotype and dementia prevalence in an admixed Cuban sample; a cross-sectional population survey and nested case-control study. BMC Medical Genetics, 2011, 12, 43.	2.1	25

#	Article	IF	CITATIONS
91	Equity in the delivery of community healthcare to older people: findings from 10/66 Dementia Research Group cross-sectional surveys in Latin America, China, India and Nigeria. BMC Health Services Research, 2011, 11, 153.	2.2	80
92	The association between common physical impairments and dementia in low and middle income countries, and, among people with dementia, their association with cognitive function and disability. A 10/66 Dementia Research Group population-based study. International Journal of Geriatric Psychiatry, 2011, 26, 511-519.	2.7	33
93	A brief dementia screener suitable for use by nonâ€specialists in resource poor settings—the crossâ€cultural derivation and validation of the brief Community Screening Instrument for Dementia. International Journal of Geriatric Psychiatry, 2011, 26, 899-907.	2.7	85
94	Tobacco use and dementia: evidence from the 1066 dementia populationâ€based surveys in Latin America, China and India. International Journal of Geriatric Psychiatry, 2011, 26, 1177-1185.	2.7	16
95	Violence during pregnancy and newborn outcomes: a cohort study in a disadvantaged population in Brazil. European Journal of Public Health, 2011, 21, 92-97.	0.3	41
96	Prevalence of stroke and related burden among older people living in Latin America, India and China. Journal of Neurology, Neurosurgery and Psychiatry, 2011, 82, 1074-1082.	1.9	137
97	Co-occurrence of anxiety and depression amongst older adults in low- and middle-income countries: findings from the 10/66 study. Psychological Medicine, 2011, 41, 2047-2056.	4.5	53
98	Mental and personality disorders and abstinence from alcohol: results from a national household survey. Psychological Medicine, 2011, 41, 809-818.	4.5	14
99	Prevalence of anxiety and its correlates among older adults in Latin America, India and China: cross-cultural study. British Journal of Psychiatry, 2011, 199, 485-491.	2.8	97
100	Chronic Disease Prevalence and Care Among the Elderly in Urban and Rural Beijing,China — A 10/66 Dementia Research Group Cross-Sectional Survey. , 2011, , 13-32.		0
101	10/66 Dementia Research Group: recently published survey data for seven Latin America sites. International Psychogeriatrics, 2010, 22, 158-159.	1.0	6
102	Violence and depressive symptoms during pregnancy: a primary care study in Brazil. Social Psychiatry and Psychiatric Epidemiology, 2010, 45, 983-988.	3.1	17
103	The prevalence and social patterning of chronic diseases among older people in a population undergoing health transition. A 10/66 Group cross-sectional population-based survey in the Dominican Republic. BMC Public Health, 2010, 10, 344.	2.9	17
104	An investigation of associations between alcohol use disorder and polymorphisms on <i>ALDH2</i> , <i>BDNF</i> , <i>5â€HTTLPR</i> , and <i>MTHFR</i> genes in older Korean men. International Journal of Geriatric Psychiatry, 2010, 25, 441-448.	2.7	31
105	Measuring disability across cultures — the psychometric properties of the WHODAS II in older people from seven low―and middleâ€income countries. The 10/66 Dementia Research Group populationâ€based survey. International Journal of Methods in Psychiatric Research, 2010, 19, 1-17.	2.1	88
106	"The contribution of chronic diseases to the prevalence of dependence among older people in Latin America, China and India: a 10/66 Dementia Research Group population-based survey". BMC Geriatrics, 2010, 10, 53.	2.7	94
107	Nutrition, mental health and violence: from pregnancy to postpartum Cohort of women attending primary care units in Southern Brazil - ECCAGE study. BMC Psychiatry, 2010, 10, 66.	2.6	34
108	Depression in older people is associated with increased risk of developing dementia. Evidence-Based Mental Health, 2010, 13, 112-112.	4.5	1

#	Article	IF	CITATIONS
109	Asthma and Suicide Mortality in Young People: A 12-Year Follow-Up Study. American Journal of Psychiatry, 2010, 167, 1092-1099.	7.2	82
110	Tobacco use among the elderly: The first Brazilian National Survey (BNAS). Aging and Mental Health, 2010, 14, 720-724.	2.8	8
111	Suicide attempts prior to starting methadone maintenance treatment in Taiwan. Drug and Alcohol Dependence, 2010, 109, 139-143.	3.2	38
112	Late-life depression in Peru, Mexico and Venezuela: the 10/66 population-based study. British Journal of Psychiatry, 2009, 195, 510-515.	2.8	79
113	A Population-Based Study of the Association between <i>Trypanosoma cruzi</i> Infection and Cognitive Impairment in Old Age (The BambuÃ-Study). Neuroepidemiology, 2009, 32, 122-128.	2.3	37
114	Dietary fish and meat intake and dementia in Latin America, China, and India: a 10/66 Dementia Research Group population-based study. American Journal of Clinical Nutrition, 2009, 90, 392-400.	4.7	84
115	Population normative data for the 10/66 Dementia Research Group cognitive test battery from Latin America, India and China: a cross-sectional survey. BMC Neurology, 2009, 9, 48.	1.8	85
116	Chronic disease prevalence and care among the elderly in urban and rural Beijing, China - a 10/66 Dementia Research Group cross-sectional survey. BMC Public Health, 2009, 9, 394.	2.9	81
117	Helping carers to care—The 10/66 dementia research group's randomized control trial of a caregiver intervention in Russia. International Journal of Geriatric Psychiatry, 2009, 24, 347-354.	2.7	85
118	Violence and post-traumatic stress disorder in Sao Paulo and Rio de Janeiro, Brazil: the protocol for an epidemiological and genetic survey. BMC Psychiatry, 2009, 9, 34.	2.6	38
119	The Epidemiology of Dependence in Older People in Nigeria: Prevalence, Determinants, Informal Care, and Health Service Utilization. A 10/66 Dementia Research Group Cross‧ectional Survey. Journal of the American Geriatrics Society, 2009, 57, 1620-1627.	2.6	50
120	Contribution of chronic diseases to disability in elderly people in countries with low and middle incomes: a 10/66 Dementia Research Group population-based survey. Lancet, The, 2009, 374, 1821-1830.	13.7	370
121	Dimensions Underlying the Mini-Mental State Examination in a Sample With Low-Education Levels: The BambuÃ-Health and Aging Study. American Journal of Geriatric Psychiatry, 2009, 17, 863-872.	1.2	29
122	Validation of the 10/66 Dementia Research Group's 10/66 Dementia diagnosis in Iran. International Psychogeriatrics, 2009, 21, 604.	1.0	6
123	The 10/66 Dementia Research Group's fully operationalised DSM-IV dementia computerized diagnostic algorithm, compared with the 10/66 dementia algorithm and a clinician diagnosis: a population validation study. BMC Public Health, 2008, 8, 219.	2.9	108
124	The epidemiology of dependency among urban-dwelling older people in the Dominican Republic; a cross-sectional survey. BMC Public Health, 2008, 8, 285.	2.9	40
125	Severity of heroin dependence in Taiwan: Reliability and validity of the Chinese version of the Severity of Dependence Scale (SDS[Ch]). Addictive Behaviors, 2008, 33, 1590-1593.	3.0	37
126	Alcohol consumption in late-life — The first Brazilian National Alcohol Survey (BNAS). Addictive Behaviors, 2008, 33, 1598-1601.	3.0	37

#	Article	IF	CITATIONS
127	Prevalence of dementia in Latin America, India, and China: a population-based cross-sectional survey. Lancet, The, 2008, 372, 464-474.	13.7	358
128	Ageing and dementia in low and middle income countries–Using research to engage with public and policy makers. International Review of Psychiatry, 2008, 20, 332-343.	2.8	62
129	The Prevalence, Correlates and Impact of Dementia in Cuba. Neuroepidemiology, 2008, 31, 243-251.	2.3	37
130	Measures of effect for cross-sectional studies. Revista Brasileira De Psiquiatria, 2008, 30, 399-399.	1.7	1
131	Intermittent preventive treatment of malaria during Pathways to homelessness among older people in Rio de Janeiro, Brazil. Bulletin of the World Health Organization, 2007, 85, 888-889.	3.3	7
132	The protocols for the 10/66 dementia research group population-based research programme. BMC Public Health, 2007, 7, 165.	2.9	298
133	The impact of maternal experience of violence and common mental disorders on neonatal outcomes: a survey of adolescent mothers in Sao Paulo, Brazil. BMC Public Health, 2007, 7, 209.	2.9	84
134	Global prevalence of dementia: a Delphi consensus study. Lancet, The, 2005, 366, 2112-2117.	13.7	4,496
135	Behavioral and Psychological Symptoms of Dementia in developing countries. International Psychogeriatrics, 2004, 16, 441-459.	1.0	87
136	Resource utilisation for neuropsychiatric disorders in developing countries:. Social Psychiatry and Psychiatric Epidemiology, 2004, 39, 218-227.	3.1	30
137	Factors associated with adverse reactions to cocaine among a sample of long-term, high-dose users in São Paulo, Brazil. Addictive Behaviors, 2004, 29, 365-374.	3.0	20
138	Differences in factors associated with first treatment entry and treatment reâ€entry among cocaine users. Addiction, 2002, 97, 825-832.	3.3	14
139	Does multisite sampling improve patient heterogeneity in drug misuse research?. Drug and Alcohol Dependence, 2001, 63, 79-85.	3.2	12
140	HIGH DOSE COCAINE USE IN SÃ $f$ O PAULO: A COMPARISON OF TREATMENT AND COMMUNITY SAMPLES. Substance Use and Misuse, 2001, 36, 237-255.	1.4	19
141	Validity and reliability of the Severity of Dependence Scale (SDS) in a Brazilian sample of drug users. Drug and Alcohol Review, 2000, 19, 451-455.	2.1	59
142	Epidemiological methods for research with drug misusers: review of methods for studying prevalence and morbidity. Revista De Saude Publica, 1999, 33, 206-215.	1.7	24
143	Route of cocaine administration. Addictive Behaviors, 1999, 24, 815-821.	3.0	50
144	Letter to the Editor. Addiction, 1998, 93, 287-288.	3.3	11

#	Article	IF	CITATIONS
145	Crack Cocaine: An Increase in Use among Patients Attending Clinics in SaÂ <sup>-</sup> o Paulo: 1990-1993. Substance Use and Misuse, 1996, 31, 519-527.	1.4	67
146	Alcohol dependence criteria in DSM-III-R: presence of symptoms according to degree of severity. Addiction, 1994, 89, 1129-1134.	3.3	10
147	Experiences of stigma and discrimination among people living with dementia and family carers in Brazil: qualitative study. Ageing and Society, 0, , 1-22.	1.7	6