

Yoshinori Asakawa

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3280072/publications.pdf>

Version: 2024-02-01

345
papers

9,745
citations

47006
47
h-index

102487
66
g-index

359
all docs

359
docs citations

359
times ranked

4029
citing authors

#	ARTICLE	IF	CITATIONS
1	Chemosystematics of the Hepaticae. <i>Phytochemistry</i> , 2004, 65, 623-669.	2.9	206
2	Phytochemical and biological studies of bryophytes. <i>Phytochemistry</i> , 2013, 91, 52-80.	2.9	199
3	Biologically active compounds from bryophytes. <i>Pure and Applied Chemistry</i> , 2007, 79, 557-580.	1.9	175
4	Recent advances in phytochemistry of bryophytes-acetogenins, terpenoids and bis(bibenzyl)s from selected Japanese, Taiwanese, New Zealand, Argentinean and European liverworts. <i>Phytochemistry</i> , 2001, 56, 297-312.	2.9	159
5	Chemical Constituents of Bryophytes: Structures and Biological Activity. <i>Journal of Natural Products</i> , 2018, 81, 641-660.	3.0	141
6	Novel neurotrophic isocuparane-type sesquiterpene dimers, mastigophorenes A, B, C and D, isolated from the liverwort Mastigophora diciados. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 2737.	0.9	103
7	Cyclic bis(bibenzyls) and related compounds from the liverworts <i>Marchantia polymorpha</i> and <i>Marchantia palmata</i> . <i>Phytochemistry</i> , 1987, 26, 1811-1816.	2.9	91
8	New Bibenzyl Cannabinoid from the New Zealand Liverwort <i>Radula marginata</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2002, 50, 1390-1392.	1.3	90
9	Prenyl bibenzyls from the liverworts <i>Radula perrottetii</i> and <i>Radula complanata</i> . <i>Phytochemistry</i> , 1991, 30, 235-251.	2.9	89
10	Chemical Constituents of the Ascomycete <i>Daldinia concentrica</i> . <i>Journal of Natural Products</i> , 2002, 65, 1869-1874.	3.0	88
11	Chemical structures of macrocyclic bis(bibenzyls) isolated from liverworts (Hepaticae). <i>Spectroscopy</i> , 2000, 14, 149-175.	0.8	86
12	Introduction. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, 95, 1-16.	1.1	77
13	Riccardin C, a novel cyclic bibenzyl derivative from <i>Reboulia hemisphaerica</i> . <i>Phytochemistry</i> , 1982, 21, 2143-2144.	2.9	74
14	Inedible mushrooms: a good source of biologically active substances. <i>Chemical Record</i> , 2006, 6, 79-99.	5.8	74
15	Anti-Influenza Activity of Marchantins, Macrocylic Bisbibenzyls Contained in Liverworts. <i>PLoS ONE</i> , 2011, 6, e19825.	2.5	73
16	Bryophytes: Bio- and Chemical Diversity, Bioactivity and Chemosystematics. <i>Heterocycles</i> , 2009, 77, 99.	0.7	70
17	Phenolic constituents of the liverwort: Four novel cyclic bisbibenzyl dimers from <i>Blasia pusilla</i> L. <i>Tetrahedron</i> , 1996, 52, 14487-14500.	1.9	69
18	Cohaeirins A and B, azaphilones from the fungus <i>Hypoxyylon cohaerens</i> , and comparison of HPLC-based metabolite profiles in <i>Hypoxyylon</i> sect. <i>Annulata</i> . <i>Phytochemistry</i> , 2005, 66, 797-809.	2.9	67

#	ARTICLE	IF	CITATIONS
19	Riccardin C: A natural product that functions as a liver X receptor (LXR) α agonist and an LXR β antagonist. <i>FEBS Letters</i> , 2005, 579, 5299-5304.	2.8	66
20	A potent cytotoxic warburganal and related drimane-type sesquiterpenoids from <i>Polygonum hydropiper</i> . <i>Phytochemistry</i> , 1980, 21, 2895-2898.	2.9	65
21	Neogrinolin derivatives possessing anti-oxidative activity from the mushroom <i>Albatrellus ovinus</i> . <i>Phytochemistry</i> , 2002, 59, 731-737.	2.9	65
22	Riccardin A and riccardin B, two novel cyclic bis(bibenzyls) possessing cytotoxicity from the liverwort <i>Riccardia multifida</i> (L.) S. Gray. <i>Journal of Organic Chemistry</i> , 1983, 48, 2164-2167.	3.2	64
23	Biosynthesis of cyclic bis(bibenzyls) in <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1999, 50, 589-598.	2.9	64
24	Biologically Active Substances of Japanese Inedible Mushrooms. <i>Heterocycles</i> , 1998, 47, 1067.	0.7	62
25	Grifolin derivatives from <i>Albatrellus caeruleoporus</i> , new inhibitors of nitric oxide production in RAW 264.7 cells. <i>Bioorganic and Medicinal Chemistry</i> , 2006, 14, 164-168.	3.0	62
26	Perrottetins E, F, and G from (liverwort)-isolation, structure determination, and synthesis of perrottetin e. <i>Tetrahedron Letters</i> , 1985, 26, 6097-6100.	1.4	58
27	New Chlorinated Cyclic Bis(bibenzyls) from the Liverworts <i>Herbertus sakuraii</i> and <i>Mastigophora diclados</i> . <i>Tetrahedron</i> , 2000, 56, 3153-3159.	1.9	57
28	In vitro antitrypanosomal activity of plant terpenes against <i>Trypanosoma brucei</i> . <i>Phytochemistry</i> , 2011, 72, 2024-2030.	2.9	57
29	Ent-sesquiterpenoids and cyclic bis(bibenzyls) from the german liverwort <i>Marchantia Polymorpha</i> . <i>Phytochemistry</i> , 1990, 29, 1577-1584.	2.9	55
30	(+)-Cavicularin: A novel optically active cyclic bibenzyl-dihydrophenanthrene derivative from the liverwort <i>Cavicularia densa</i> Steph. <i>Tetrahedron Letters</i> , 1996, 37, 4745-4748.	1.4	54
31	Changes in secondary metabolism during stromatal ontogeny of <i>Hypoxylon fragiforme</i> . <i>Mycological Research</i> , 2006, 110, 811-820.	2.5	54
32	Sesquiterpenes from <i>Porella</i> species. <i>Phytochemistry</i> , 1978, 17, 457-460.	2.9	53
33	Distribution of novel cyclic bisbibenzyls in <i>Marchantia</i> and <i>Riccardia</i> species. <i>Phytochemistry</i> , 1983, 22, 1413-1415.	2.9	53
34	Structures of a Novel Binaphthyl and Three Novel Benzophenone Derivatives with Plant-Growth Inhibitory Activity from the Fungus <i>Daldinia concentrica</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1528-1530.	1.3	53
35	(+)-Osmundalactone, β -lactones and spiromentins from the fungus <i>Paxillus atrotomentosus</i> . <i>Phytochemistry</i> , 1995, 40, 1251-1257.	2.9	53
36	Terpenoids of the liverwort <i>Frullanoides densifolia</i> and <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 1993, 32, 335-348.	2.9	52

#	ARTICLE		IF	CITATIONS
37	Bryophytes as a source of bioactive volatile terpenoids – A review. <i>Food and Chemical Toxicology</i> , 2019, 132, 110649.		3.6	52
38	Seven new bibenzyls and a dihydrochalcone from <i>Radula variabilis</i> . <i>Phytochemistry</i> , 1978, 17, 2005-2010.		2.9	51
39	Total assignment of ¹ H and ¹³ C NMR spectra of marchantins isolated from liverworts and its application to structure determination of two new macrocyclic bis(bibenzyls) from and. <i>Tetrahedron Letters</i> , 1985, 26, 4735-4738.		1.4	51
40	Cryptoporic acids A-G, drimane-type sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . <i>Phytochemistry</i> , 1992, 31, 579-592.		2.9	51
41	Terpenoids and aromatic compounds from six liverworts. <i>Phytochemistry</i> , 1996, 41, 207-211.		2.9	50
42	Pinguisane and dimeric pinguisane-type sesquiterpenoids from the Japanese liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . <i>Phytochemistry</i> , 2000, 53, 593-604.		2.9	50
43	Cyclic azaphilones daldinins E and F from the ascomycete fungus <i>Hypoxylon fuscum</i> (Xylariaceae). <i>Phytochemistry</i> , 2004, 65, 469-473.		2.9	50
44	Novel cytotoxic kaurane-type diterpenoids from the New Zealand Liverwort <i>Jungermannia</i> species. <i>Tetrahedron</i> , 2005, 61, 4531-4544.		1.9	50
45	Inhibition of Nitric Oxide Production in RAW 264.7 Cells by Azaphilones from Xylariaceous Fungi. <i>Biological and Pharmaceutical Bulletin</i> , 2006, 29, 34-37.		1.4	50
46	Chemical Constituents of Bryophytes. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , .		1.1	50
47	Sesquiterpenes from Japanese liverworts. <i>Phytochemistry</i> , 1981, 20, 2359-2366.		2.9	49
48	A comparative study on three chemo-types of the liverwort <i>Conocephalum conicum</i> using volatile constituents. <i>Phytochemistry</i> , 1997, 44, 1265-1270.		2.9	48
49	Cytochromes P-450 catalyze the formation of marchantins A and C in <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1999, 52, 1195-1202.		2.9	48
50	Antioxidant Activity of Curtisiains I - L from the Inedible Mushroom <i>Paxillus curtisii</i> . <i>Planta Medica</i> , 2003, 69, 1063-1066.		1.3	48
51	Prenyl bibenzyls from the liverwort <i>Radula kojana</i> . <i>Phytochemistry</i> , 1991, 30, 219-234.		2.9	47
52	Sesquiterpenoids from the liverworts <i>Bazzania trilobata</i> and <i>Porella canariensis</i> . <i>Phytochemistry</i> , 1996, 42, 1361-1366.		2.9	47
53	Molecular chemotaxonomy of Daldinia and other Xylariaceae. <i>Mycological Research</i> , 2001, 105, 1191-1205.		2.5	47
54	Cytotoxic and Apoptosis-Inducing <i>< i>ent</i>-Kaurane-Type Diterpenoids from the Japanese Liverwort < i>Jungermannia truncata</i> N< small>EES</small></i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2002, 50, 808-813.		1.3	47

#	ARTICLE	IF	CITATIONS
55	New Azaphilones from the Inedible Mushroom <i>Hypoxylon rubiginosum</i> . <i>Journal of Natural Products</i> , 2004, 67, 1152-1155.	3.0	47
56	Novel bibenzyl derivatives and ent-cuparene-type sesquiterpenoids from <i>Radula</i> species. <i>Phytochemistry</i> , 1982, 21, 2481-2490.	2.9	46
57	Fusicoccane-, dolabellane- and rearranged labdane-type diterpenoids from the liverwort <i>Pleurozia gigantea</i> . <i>Phytochemistry</i> , 1990, 29, 2597-2603.	2.9	46
58	Structures of Daldinins A-C, Three Novel Azaphilone Derivatives from Ascomycetous Fungus <i>Daldinia concentrica</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2397-2399.	1.3	46
59	Herbertane-type sesquiterpenoids from the liverwort <i>Herbertus sakuraii</i> . <i>Phytochemistry</i> , 2000, 55, 247-253.	2.9	46
60	Bis(bibenzyls) from Liverworts Inhibit Lipopolysaccharide-Induced Inducible NOS in RAW 264.7 Cells: A Study of Structure-Activity Relationships and Molecular Mechanism. <i>Journal of Natural Products</i> , 2005, 68, 1779-1781.	3.0	46
61	Isoprenyl Phenyl Ethers from Liverworts of the Genus <i>Trichocolea</i> : Cytotoxic Activity, Structural Corrections, and Synthesis. <i>Journal of Natural Products</i> , 1996, 59, 729-733.	3.0	45
62	Isopladiochins C and D, New Type of Macroyclic Bis(bibenzyls), Having Two Biphenyl Linkages from the Liverwort <i>Plagiochila fruticosa</i> . <i>Chemistry Letters</i> , 1996, 25, 741-742.	1.3	45
63	Cytochalasins from a <i>Daldinia</i> sp. of fungus. <i>Phytochemistry</i> , 1996, 41, 821-828.	2.9	45
64	Iridoid glucosides from roots of Vietnamese <i>Paederia scandens</i> . <i>Phytochemistry</i> , 2002, 60, 505-514.	2.9	45
65	Sassafrins A-D, new antimicrobial azaphilones from the fungus <i>Creosphaeria sassafras</i> . <i>Tetrahedron</i> , 2005, 61, 1743-1748.	1.9	45
66	Terpenoids and bibenzyls of 25 liverwort <i>Frullania</i> species. <i>Phytochemistry</i> , 1981, 20, 2187-2194.	2.9	42
67	Application of Chromatographic and Spectroscopic Methods towards the Quality Assessment of Ginger (<i>Zingiber officinale</i>) Rhizomes from Ecological Plantations. <i>International Journal of Molecular Sciences</i> , 2017, 18, 452.	4.1	42
68	New ent-Kaurene-Type Diterpenoids Possessing Cytotoxicity from the New Zealand Liverwort <i>Jungermannia</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1189-1192.	1.3	41
69	Sesquiterpenes of six <i>Porella</i> species (hepaticae). <i>Phytochemistry</i> , 1976, 15, 1929-1931.	2.9	40
70	Isotachin A and isotachin B, two sulphur-containing acrylates from the liverwort <i>isotachis japonica</i> . <i>Phytochemistry</i> , 1985, 24, 1505-1508.	2.9	40
71	Cytotoxic, radical scavenging and antimicrobial activities of sesquiterpenoids from the Tahitian liverwort <i>Mastigophora diclados</i> (Brid.) Nees (Mastigophoraceae). <i>Journal of Natural Medicines</i> , 2010, 64, 417-422.	2.3	40
72	Homomono- and sesquiterpenoids from the liverwort <i>Lophocolea heterophylla</i> . <i>Phytochemistry</i> , 1990, 29, 2334-2337.	2.9	39

#	ARTICLE	IF	CITATIONS
73	Zierane sesquiterpene lactone, cembrane and fusicoccane diterpenoids, from the Tahitian liverwort <i>Chandonanthus hirtellus</i> . <i>Phytochemistry</i> , 2010, 71, 1387-1394.	2.9	39
74	Marchantin A Trimethyl Ether: Its Molecular Structure and Tubocurarine-like Skeletal Muscle Relaxation Activity.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 52-56.	1.3	38
75	Novel skeletal diterpenoids from the Japanese liverwort <i>Pallavicinia subciliata</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 178-180.	1.3	38
76	Volatile components of selected species of the liverwort genera <i>Frullania</i> and <i>Schusterella</i> (Frullaniaceae) from New Zealand, Australia and South America: a chemosystematic approach. <i>Phytochemistry</i> , 2003, 62, 439-452.	2.9	38
77	Apoptosis-Inducing Properties ofent-Kaurene-Type Diterpenoids from the Liverwort <i>Jungermannia truncata</i> . <i>Planta Medica</i> , 2003, 69, 377-379.	1.3	38
78	Kaurene Diterpene Induces Apoptosis in Human Leukemia Cells Partly through a Caspase-8-Dependent Pathway. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2004, 311, 115-122.	2.5	38
79	Antimicrobial Azaphilones from the Fungus <i>Hypoxylon multiforme</i> . <i>Planta Medica</i> , 2005, 71, 1058-1062.	1.3	38
80	Plagiochins A, B, C, and D, new type of macrocyclic bis(bibenzyls) having a biphenyl linkage between the ortho positions to the benzyl methylenes, from the liverwort subsp.. <i>Tetrahedron Letters</i> , 1987, 28, 6295-6298.	1.4	37
81	Sesquiterpene and other constituents of the liverwort <i>Dumortiera hirsuta</i> . <i>Phytochemistry</i> , 1997, 44, 293-298.	2.9	37
82	Chemical Constituents of Malagasy Liverworts, Part II: Mastigophoric Acid Methyl Ester of Biogenetic Interest from <i>Mastigophora diclados</i> (Lepicoleaceae Subf. Mastigophoroideae). <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 1382-1384.	1.3	37
83	Antimitotic activity of two macrocyclic bis(bibenzyls), isoplagiochins A and B from the Liverwort <i>Plagiochila fruticosa</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 493-496.	2.2	37
84	Bibenzyl derivatives from the Australian liverwort <i>Frullania falciloba</i> . <i>Phytochemistry</i> , 1987, 26, 1023-1025.	2.9	36
85	Five 10-phenyl-[11]-cytochalasans from a <i>Daldinia</i> fungal species. <i>Phytochemistry</i> , 1995, 40, 135-140.	2.9	36
86	Terpenoids from the Japanese liverworts <i>Jackiella javanica</i> and <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 1997, 46, 1203-1208.	2.9	36
87	Sesqui- and diterpenoids from two Japanese and three European liverworts. <i>Phytochemistry</i> , 2001, 56, 347-352.	2.9	36
88	Thelephantins A, B and C: three benzoyl p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . <i>Phytochemistry</i> , 2003, 62, 109-113.	2.9	36
89	Thelephantins D-H: five p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . <i>Phytochemistry</i> , 2003, 63, 919-924.	2.9	36
90	Chemical Constituents of Malagasy Liverworts, Part V: Prenyl Bibenzyls and Clerodane Diterpenoids with Nitric Oxide Inhibitory Activity from <i>Radula appressa</i> and <i>Thysananthus spathulistipus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 1046-1049.	1.3	36

#	ARTICLE	IF	CITATIONS
91	Synthesis of riccardin C and its seven analogues. Part 1: The role of their phenolic hydroxy groups as LXR \pm agonists. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 738-741.	2.2	36
92	Fatty acids and cyclic bis(bibenzyls) from the New Zealand liverwort <i>Monoclea forsteri</i> . <i>Phytochemistry</i> , 1988, 27, 2603-2608.	2.9	35
93	Terpenoids and aromatic compounds from selected Ecuadorian liverworts*. <i>Phytochemistry</i> , 1991, 30, 215-217.	2.9	35
94	Cytotoxic 2,3-Secoiridoid-Type Sesquiterpenoids from the Liverwort <i>Plagiochila ovalifolia</i> . <i>Planta Medica</i> , 1998, 64, 462-464.	1.3	35
95	Eremophilanolides and other constituents from the Argentine liverwort <i>Frullania brasiliensis</i> . <i>Phytochemistry</i> , 2002, 59, 205-213.	2.9	35
96	Anent-Kaurene Diterpene Enhances Apoptosis Induced by Tumor Necrosis Factor in Human Leukemia Cells. <i>Planta Medica</i> , 2004, 70, 723-727.	1.3	35
97	Efficient synthesis of isoplagiochin D, a macrocyclic bis(bibenzyls), by utilizing an intramolecular Suzuki-Miyaura reaction. <i>Tetrahedron Letters</i> , 2004, 45, 6941-6945.	1.4	35
98	Africane-Type Sesquiterpenoids from the Argentine Liverwort <i>Porella swartziana</i> and Their Antibacterial Activity. <i>Journal of Natural Products</i> , 2004, 67, 31-36.	3.0	35
99	ent-Verticillane-type diterpenoids from the Japanese liverwort <i>Jackiella javanica</i> . <i>Phytochemistry</i> , 2005, 66, 1662-1670.	2.9	35
100	New bibenzyls from <i>Radula complanata</i> . <i>Phytochemistry</i> , 1978, 17, 2115-2117.	2.9	34
101	A Highly efficient preparation of lunularic acid and some biological activities of stilbene and dihydrostilbene derivatives. <i>Phytochemistry</i> , 1988, 27, 109-113.	2.9	34
102	Highlights in phytochemistry of hepaticae-biologically active terpenoids and aromatic compounds. <i>Pure and Applied Chemistry</i> , 1994, 66, 2193-2196.	1.9	34
103	A Novel Class of Plant Type III Polyketide Synthase Involved in Orsellinic Acid Biosynthesis from <i>Rhododendron dauricum</i> . <i>Frontiers in Plant Science</i> , 2016, 7, 1452.	3.6	34
104	Sesquiterpenoids from <i>Chiloscyphus</i> , <i>Clasmatocolea</i> and <i>Frullania</i> species. <i>Phytochemistry</i> , 1983, 22, 961-964.	2.9	33
105	Sesquiterpenoids from the liverwort <i>Porella acutifolia</i> subsp. <i>Tosana</i> . <i>Phytochemistry</i> , 1991, 30, 567-573.	2.9	33
106	Terpenoids from six lophoziaeae liverworts. <i>Phytochemistry</i> , 1993, 34, 181-190.	2.9	33
107	Epi-neoverrucosane- and ent-clerodane-type diterpenoids and ent-2,3-secoiridoid- and calamenene-type sesquiterpenoids from the liverwort <i>heteroscyphus planus</i> . <i>Phytochemistry</i> , 1995, 38, 119-127.	2.9	33
108	Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> . <i>Journal of Natural Products</i> , 2001, 64, 1309-1317.	3.0	33

#	ARTICLE	IF	CITATIONS
109	The Novel Compounds That Activate Farnesoid X Receptor: the Diversity of Their Effects on Gene Expression. <i>Journal of Pharmacological Sciences</i> , 2008, 107, 285-294.	2.5	33
110	Distribution of Bibenzyls, Prenyl Bibenzyls, Bis-bibenzyls, and Terpenoids in the Liverwort Genus <i>< i>Radula</i></i> . <i>Journal of Natural Products</i> , 2020, 83, 756-769.	3.0	33
111	Concentriols B, C and D, three squalene-type triterpenoids from the ascomycete <i>Daldinia concentrica</i> . <i>Phytochemistry</i> , 2002, 61, 345-353.	2.9	32
112	Composition of the essential oil of the liverwort <i>Radula perrottetii</i> of Japanese origin. <i>Phytochemistry</i> , 2005, 66, 941-949.	2.9	32
113	Terrestrins A-G: p-Terphenyl derivatives from the inedible mushroom <i>Thelephora terrestris</i> . <i>Phytochemistry</i> , 2005, 66, 1052-1059.	2.9	32
114	Distribution of Cyclic and Acyclic Bis-bibenzyls in the Marchantiophyta (Liverworts), Ferns and Higher Plants and Their Biological Activities, Biosynthesis, and Total Synthesis. <i>Heterocycles</i> , 2012, 86, 891.	0.7	32
115	The superoxide release inhibitors, cryptoporic acids C, D, and E; dimeric drimane sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1989, , 258.	2.0	31
116	Cadinane-type sesquiterpenoids from the liverwort <i>Scapania undulata</i> . <i>Phytochemistry</i> , 1994, 37, 1323-1325.	2.9	31
117	Chenopodene, marchantin P and riccardin G from the liverwort <i>Marchantia chenopoda</i> . <i>Phytochemistry</i> , 1994, 36, 73-76.	2.9	31
118	Chapter Five: Distribution of Terpenoids and Aromatic Compounds in Selected Southern Hemispheric Liverworts. <i>Fieldiana Botany</i> , 2008, 47, 37.	0.3	31
119	Structures of four novel macrocyclic bis(bibenzyl) dimers, pusilatins A-D from the liverwort <i>Blasia pusilla</i> . <i>Tetrahedron Letters</i> , 1994, 35, 909-910.	1.4	30
120	Sesqui- and diterpenoids from the japanese liverwort <i>Jungermannia hattoriiana</i> . <i>Phytochemistry</i> , 1997, 45, 353-363.	2.9	30
121	Sesquiterpenoids from the three Japanese liverworts <i>Lejeunea aquatica</i> , <i>L. flava</i> and <i>L. japonica</i> . <i>Phytochemistry</i> , 1997, 46, 145-150.	2.9	30
122	Butenolides from <i>Marchantia paleacea</i> subspecies <i>diptera</i> . <i>Phytochemistry</i> , 1997, 46, 293-296.	2.9	30
123	Sesquiterpenoids, hopanoids and bis(bibenzyls) from the Argentine liverwort <i>Plagiochasma rupestre</i> . <i>Phytochemistry</i> , 1999, 52, 1323-1329.	2.9	30
124	Isolation, Synthesis and Biological Activity of Grifolic Acid Derivatives from the Inedible Mushroom <i>Albatrellus dispansus</i> . <i>Heterocycles</i> , 2005, 65, 2431.	0.7	30
125	Cryptoporic acids A and B, novel bitter drimane sesquiterpenoid ethers of isocitric acid, from the fungus. <i>Tetrahedron Letters</i> , 1987, 28, 6303-6304.	1.4	29
126	First syntheses of 1,13- and 1,15-dihydroxyherbertenes, and herbertenolide by applying intramolecular Heck reaction for the construction of adjacent quaternary centers. <i>Tetrahedron</i> , 2001, 57, 9299-9307.	1.9	29

#	ARTICLE	IF	CITATIONS
127	New Humulane-Type Sesquiterpenes from the Liverworts <i>Tylimanthus tenellus</i> and <i>Marchantia emarginata</i> subsp. <i>tosana</i> . Chemical and Pharmaceutical Bulletin, 2004, 52, 481-484.	1.3	29
128	Cinnamacrins A ¹ -C, Cinnafragrin D, and Cytostatic Metabolites with β -Glucosidase Inhibitory Activity from <i>Cinnamosma macrocarpa</i> . Journal of Natural Products, 2007, 70, 277-282.	3.0	29
129	Marchantiophyta (Liverworts): Rich Sources of Macroyclic Bis(bibenzyls). Heterocycles, 2008, 76, 99.	0.7	29
130	Cyclopropanochroman derivatives from the liverwort <i>Radula javanica</i> . Phytochemistry, 1991, 30, 325-328.	2.9	28
131	Terpenoids from some German and Russian liverworts. Phytochemistry, 1993, 33, 1445-1448.	2.9	28
132	A 10-phenyl-[11]-cytochalasan from a species of <i>Daldinia</i> . Phytochemistry, 1996, 42, 173-176.	2.9	28
133	Scapaundulins A and B, two novel dimeric labdane diterpenoids, and related compounds from the Japanese liverwort <i>Scapania undulata</i> (L.) Dum.. Tetrahedron Letters, 1997, 38, 1975-1978.	1.4	28
134	Volatile components of the liverworts <i>Archilejeunea olivacea</i> , <i>cheilolejeunea imbricata</i> and <i>Leptolejeunea elliptica</i> . Phytochemistry, 1997, 44, 1261-1264.	2.9	28
135	Ptychantols A-C, macrocyclic bis(bibenzyls), possessing a trans-stilbene structure from the liverwort <i>Ptychanthus striatus</i> . Phytochemistry, 1999, 52, 501-509.	2.9	28
136	Occurrence of Polygodial and 1-(2,4,6-Trimethoxyphenyl)-but-2-en-1-one from Some Ferns and Liverworts: Role of Pungent Components in Bryophytes and Pteridophytes Evolution.. Chemical and Pharmaceutical Bulletin, 2001, 49, 1380-1381.	1.3	28
137	Terpenoids and Aromatic Compounds from the New Zealand Liverworts <i>Plagiochila</i> , <i>Schistochila</i> , and <i>Heteroscyphus</i> Species. Chemical and Pharmaceutical Bulletin, 2004, 52, 556-560.	1.3	28
138	Insect Antifeedant Sesquiterpene Acetals from the Liverwort < i>Lepidolaena clavigera</i>. 2. Structures, Artifacts, and Activity. Journal of Natural Products, 2008, 71, 258-261.	3.0	28
139	Sesqui- and diterpenoids from <i>Plagiochila</i> species. Phytochemistry, 1994, 36, 1425-1430.	2.9	27
140	The absolute structures of new 1.BETA.-hydroxysacculatane-type diterpenoids with piscidal activity from the liverwort <i>Pellia endiviifolia</i> .. Chemical and Pharmaceutical Bulletin, 1995, 43, 2030-2032.	1.3	27
141	Terpenoid constituents of the liverwort <i>Heteroscyphus coalitus</i> . Phytochemistry, 1996, 41, 575-580.	2.9	27
142	Identification of cryptic species within liverwort <i>Conocephalum conicum</i> based on the volatile components. Phytochemistry, 2013, 95, 234-241.	2.9	27
143	Dietary intake of metals by the young adult population of Eastern Poland: Results from a market basket study. Journal of Trace Elements in Medicine and Biology, 2016, 35, 36-42.	3.0	27
144	Bibenzyl derivatives from <i>Frullania</i> Species. Phytochemistry, 1987, 26, 1117-1122.	2.9	26

#	ARTICLE	IF	CITATIONS
145	Labdane type diterpenoids from the liverwort <i>Frullania hamachiloba</i> . <i>Phytochemistry</i> , 1988, 27, 1789-1793.	2.9	26
146	Curtisians M-Q: Five Novel p-Terphenyl Derivatives from the Mushroom <i>Paxillus curtisii</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1064-1067.	1.3	26
147	A Comparison of Apoptosis and Necrosis Induced by ent-Kaurene-Type Diterpenoids in HL-60 Cells. <i>Planta Medica</i> , 2004, 70, 401-406.	1.3	26
148	In vitro antitrypanosomal activity of bis(bibenzyls)s and bibenzyls from liverworts against <i>Trypanosoma brucei</i> . <i>Journal of Natural Medicines</i> , 2012, 66, 377-382.	2.3	26
149	Occurrence of a potent piscicidal diterpenedial in the liverwort <i>Riccardia lobata</i> var. <i>Yakushimensis</i> . <i>Phytochemistry</i> , 1985, 24, 261-262.	2.9	25
150	Isotachin c and balantiolide two aromatic compounds from the new zealand liverwort <i>balantiopsis rosea</i> . <i>Phytochemistry</i> , 1986, 25, 2543-2546.	2.9	25
151	Novel ent-vibsane- and dolabellane-type diterpenoids from the liverwort <i>Odontoschisma denudatum</i> . <i>Tetrahedron Letters</i> , 1998, 39, 579-582.	1.4	25
152	Phytochemistry of Bryophytes., 1999, , 319-342.		25
153	Occurrence of a Bis-bibenzyl Derivative in the Japanese Fern <i>Hymenophyllum barbatum</i> : First Isolation and Identification of Perrottetin H from the Pteridophytes.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 297-298.	1.3	25
154	Identification and Characterization of Daurichromenic Acid Synthase Active in Anti-HIV Biosynthesis. <i>Plant Physiology</i> , 2017, 174, 2213-2230.	4.8	25
155	Bibenzyls from <i>Radula tokiensis</i> and <i>R. Japonica</i> . <i>Phytochemistry</i> , 1981, 20, 858-859.	2.9	24
156	Cuparane- and isocuparane-type sesquiterpenoids in liverworts of the genus <i>Herbertus</i> . <i>Phytochemistry</i> , 1982, 21, 2471-2473.	2.9	24
157	Distribution of cyclic bis(bibenzyls) in the South African liverwort <i>Marchantia polymorpha</i> †. <i>Phytochemistry</i> , 1988, 27, 161-163.	2.9	24
158	Mastigophorenes: novel dimeric isocuparane-type sesquiterpenoids from the liverwort <i>Mastigophora diclados</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1988, , 1341.	2.0	24
159	Two novel macrocyclic bis(Bibenzyls), Isoplagiochins A and B from the liverwort <i>Plagiochila Fruticosa</i> . <i>Tetrahedron Letters</i> , 1994, 35, 911-912.	1.4	24
160	Dumortenols, Novel Skeletal Sesquiterpenoids from the Argentinian Liverwort <i>Dumortiera hirsuta</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1997, 45, 2119-2121.	1.3	24
161	Eudesmane-type sesquiterpene lactones from the Japanese liverwort <i>Frullania densiloba</i> . <i>Phytochemistry</i> , 1997, 45, 555-558.	2.9	24
162	A 7-nordumortenone and other dumortane derivatives from the Argentine liverwort <i>Dumortiera hirsuta</i> . <i>Phytochemistry</i> , 1999, 51, 281-287.	2.9	24

#	ARTICLE	IF	CITATIONS
163	ent-Isopimarane-type diterpenoids from the New Zealand liverwort <i>Trichocolea mollissima</i> . <i>Phytochemistry</i> , 2003, 64, 1319-1325.	2.9	24
164	New Sesquiterpenoids from the New Zealand Liverwort <i>Chiloscyphus subporosus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 949-952.	1.3	24
165	Synthetic Transformation of Ptychanthin into Forskolin and 1,9-Dideoxyforskolin. <i>Journal of Organic Chemistry</i> , 2006, 71, 4619-4624.	3.2	24
166	Heterocyclic Compounds Found in Bryophytes. <i>Heterocycles</i> , 1997, 46, 795.	0.7	24
167	Studies on the liverwort sesquiterpene alcohol tamariscol. Synthesis and absolute configuration. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 435.	0.9	23
168	Neurotrophic secoaromadendrane-type sesquiterpenes from the liverwort <i>Plagiochila fruticosa</i> . <i>Phytochemistry</i> , 1991, 30, 4061-4065.	2.9	23
169	Two novel Diels-Alder reaction-type dimeric pinguisane sesquiterpenoids and related compounds from the liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . <i>Tetrahedron Letters</i> , 1998, 39, 2977-2980.	1.4	23
170	Triterpenoid constituents of the moss <i>Floribundaria aurea</i> subsp. <i>Nipponica</i> . <i>Phytochemistry</i> , 1998, 48, 297-299.	2.9	23
171	Curtisiains E-H: four p-terphenyl derivatives from the inedible mushroom <i>Paxillus curtisi</i> . <i>Phytochemistry</i> , 2003, 64, 649-654.	2.9	23
172	Thelephantins I-N p-terphenyl derivatives from the inedible mushroom <i>Hydnellum caeruleum</i> . <i>Phytochemistry</i> , 2004, 65, 1179-1184.	2.9	23
173	Bioactive compounds from liverworts: Inhibition of lipopolysaccharide-induced inducible NOS mRNA in RAW 264.7 cells by Herbertenoids and Cuparenoids. <i>Phytomedicine</i> , 2007, 14, 486-491.	5.3	23
174	An Aromatic Farnesyltransferase Functions in Biosynthesis of the Anti-HIV Meroterpenoid Daurichromenic Acid. <i>Plant Physiology</i> , 2018, 178, 535-551.	4.8	23
175	Sesquiterpene Lactones and Acetogenin Lactones from the Hepaticae and Chemosystematics of the Liverworts <i>Frullania</i> , <i>Plagiochila</i> and <i>Porella</i> . <i>Heterocycles</i> , 2001, 54, 1057.	0.7	23
176	Terpenoids and bibenzyls from some New Zealand liverworts. <i>Phytochemistry</i> , 1982, 21, 2663-2667.	2.9	22
177	Terpenoids from the french liverwort <i>targionia hypophylla</i> . <i>Phytochemistry</i> , 1986, 25, 2555-2556.	2.9	22
178	Sesqui- and diterpenoids from the panamanian liverwort <i>Bryopteris filicina</i> . <i>Phytochemistry</i> , 1994, 37, 433-439.	2.9	22
179	New acorane- and cuparane-type sesqui- and new labdane- and seco-labdanne-type diterpenoids from the Japanese liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. <i>Tetrahedron</i> , 1999, 55, 9117-9132.	1.9	22
180	Dimeric azaphilones from the xylariaceous ascomycete <i>Hypoxylon rutilum</i> . <i>Tetrahedron</i> , 2005, 61, 8451-8455.	1.9	22

#	ARTICLE	IF	CITATIONS
181	Diterpenoids and Aromatic Compounds from the Three New Zealand Liverworts <i>Jamesoniella kirkii</i> , <i>Balantiopsis rosea</i> , and <i>Radula</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 902-906.	1.3	22
182	Fingerprinting of Secondary Metabolites of Liverworts: Chemosystematic Approach. <i>Journal of AOAC INTERNATIONAL</i> , 2014, 97, 1234-1243.	1.5	22
183	Cytotoxic Activity of Riccardin and Perrottetin Derivatives from the Liverwort <i>Lunularia cruciata</i> . <i>Journal of Natural Products</i> , 2019, 82, 694-701.	3.0	22
184	Inhibitory Effect of Cryptoporic Acid E, a Product from Fungus <i>Cryptoporus volvatus</i> , on Colon Carcinogenesis Induced with N-Methyl-N-Nitrosourea in Rats and with 1,2-Dimethylhydrazine in Mice. <i>Japanese Journal of Cancer Research</i> , 1992, 83, 830-834.	1.7	21
185	Structures of five new highly oxygenated labdane-type diterpenoids, ptychantins A-E, closely related to forskolin from the liverwort <i>Ptýchanthus striatus</i> . <i>Tetrahedron Letters</i> , 1994, 35, 5457-5460.	1.4	21
186	Total Synthesis and Absolute Configuration of Riccardiphenols A and B, Isolated from the Liverwort <i>Riccardia crassa</i> . <i>Journal of Organic Chemistry</i> , 1996, 61, 5362-5370.	3.2	21
187	Induction of Apoptosis by Newent-Kaurene-Type Diterpenoids Isolated from the New Zealand Liverwort <i>Jungermannia</i> Species. <i>Planta Medica</i> , 2005, 71, 1005-1009.	1.3	21
188	Chemical constituents of the Vietnamese inedible mushroom <i>Xylaria intracolorata</i> . <i>Natural Product Research</i> , 2006, 20, 317-321.	1.8	21
189	Secondary Metabolites of <i>Cinnamosma madagascariensis</i> and Their β -Glucosidase Inhibitory Properties. <i>Journal of Natural Products</i> , 2008, 71, 123-126.	3.0	21
190	MAO-A Inhibitory Potential of Terpene Constituents from Ginger Rhizomes—A Bioactivity Guided Fractionation. <i>Molecules</i> , 2018, 23, 1301.	3.8	21
191	Bis-benzyls, Bibenzyls, and Terpenoids in 33 Genera of the Marchantiophyta (Liverworts): Structures, Synthesis, and Bioactivity. <i>Journal of Natural Products</i> , 2022, 85, 729-762.	3.0	21
192	Drimane-type sesquiterpenoids from <i>Cryptoporus volvatus</i> infected by <i>Paecilomyces varioti</i> . <i>Phytochemistry</i> , 1993, 33, 1055-1059.	2.9	20
193	Phytol esters and phaeophytins from the hornwort <i>Megaceros flagellaris</i> . <i>Phytochemistry</i> , 1996, 41, 1373-1376.	2.9	20
194	Chemical Conversion of Labdane-Type Diterpenoid Isolated from the Liverwort <i>Porella perrottetiana</i> into (-)-Ambrox. <i>Heterocycles</i> , 1998, 49, 315.	0.7	20
195	Biotransformation of Monoterpenoids. , 2010, , 669-801.		20
196	Vasorelaxant effects of macrocyclic bis(benzyls) from liverworts. <i>Bioorganic and Medicinal Chemistry</i> , 2011, 19, 4051-4056.	3.0	20
197	Total synthesis of riccardin C and ($\Delta\pm$)-cavicularin via Pd-catalyzed Ar-Ar cross couplings. <i>Tetrahedron</i> , 2013, 69, 6959-6968.	1.9	20
198	Clerodane- and halimane-type diterpenoids from the liverwort <i>Jungermannia hyalina</i> . <i>Phytochemistry</i> , 1995, 40, 209-212.	2.9	19

#	ARTICLE	IF	CITATIONS
199	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia exsertifolia</i> ssp. <i>cordifolia</i> . <i>Phytochemistry</i> , 1996, 41, 1129-1141.	2.9	19
200	Isolation, Structure Elucidation, and Chemical Derivatization of a New Cyclic Bisbibenzyl Dimer, Pusilatin E, from the Liverwort <i>Riccardia multifida</i> subsp. <i>decrescens</i> . <i>Journal of Natural Products</i> , 1997, 60, 145-147.	3.0	19
201	Diterpenoids from the Japanese liverwort <i>jungermannia</i> INFUSCA fn1 fn1 This paper is dedicated, with best wishes, to Professor G. H. Neil Towers on the occasion of his 75th birthday. In honour of professor G. H. Neil Towers 75TH birthday. <i>Phytochemistry</i> , 1998, 49, 601-608.	2.9	19
202	Occurrence of ent-Sesquiterpene in the Japanese Moss- <i>Plagiomnium acutum</i> : First Isolation and Identification of the ent-Sesqui- and Dolabellane-type Diterpenoids from the Musci.. <i>Chemical and Pharmaceutical Bulletin</i> , 1998, 46, 1488-1489.	1.3	19
203	Highly oxygenated sesquiterpenes from the liverwort <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 2000, 53, 271-276.	2.9	19
204	Chemical Constituents of Malagasy Liverworts, Part III: Sesquiterpenoids from <i>Bazzania decrescens</i> and <i>Bazzania madagassa</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 515-518.	1.3	19
205	Lanostane triterpenoids from the inedible mushroom <i>Fomitopsis spraguei</i> . <i>Phytochemistry</i> , 2005, 66, 1656-1661.	2.9	19
206	Bitter kaurane-type diterpene glucosides from the liverwort <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 1990, 29, 1619-1623.	2.9	18
207	Stereostructure of Plagiochiline A and Conversion of Plagiochiline A and Stearylvelutinal into Hot-Tasting Compounds by Human Saliva.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1542-1544.	1.3	18
208	Terpenoids from the liverworts <i>Symphyogyna brasiliensis</i> and unidentified <i>Frullania</i> species. <i>Phytochemistry</i> , 1995, 39, 99-103.	2.9	18
209	Novel sesquiterpenoids from the Colombian liverwort <i>Porella swartziana</i> . <i>Tetrahedron</i> , 1996, 52, 6339-6354.	1.9	18
210	Eudesmane-type sesquiterpenoids from the liverwort <i>Lepidozia vitrea</i> . <i>Phytochemistry</i> , 1996, 41, 833-836.	2.9	18
211	Terpenoid constituents of the new Zealand liverwort <i>Jamesoniella tasmanica</i> . <i>Phytochemistry</i> , 1996, 43, 1057-1064.	2.9	18
212	Sesquiterpene constituents from the liverwort <i>Bazzania japonica</i> . <i>Phytochemistry</i> , 2003, 63, 581-587.	2.9	18
213	Chemotaxonomic value of essential oil components in liverwort species. A review. <i>Flavour and Fragrance Journal</i> , 2015, 30, 189-196.	2.6	18
214	Geographical distribution of tamariscol, a mossy odorous sesquiterpene alcohol, in the liverwort <i>Frullania tamarisci</i> and related species. <i>Phytochemistry</i> , 1991, 30, 2295-2300.	2.9	17
215	Differential effects of cryptoporic acids D and E, inhibitors of superoxide anion radical release, on tumor promotion of okadaic acid in mouse skin. <i>Carcinogenesis</i> , 1991, 12, 1129-1131.	2.8	17
216	Volatile constituent of the liverwort <i>chiloscyphus pallidus</i> (mitt.) engel & schuster. <i>Flavour and Fragrance Journal</i> , 1994, 9, 237-240.	2.6	17

#	ARTICLE	IF	CITATIONS
217	Eudesmane-Type Sesquiterpenoids from Japanese Liverwort <i>Frullania tamarisci</i> subsp. <i>obscura</i> .. Chemical and Pharmaceutical Bulletin, 1998, 46, 542-544.	1.3	17
218	Chemical constituents of Malagasy liverworts: Cyclomyltaylanoids from <i>Bazzania madagassa</i> . Phytochemistry, 2006, 67, 2616-2622.	2.9	17
219	Volatile Components from Selected Mexican, Ecuadorian, Greek, German and Japanese Liverworts. Natural Product Communications, 2008, 3, 1934578X0800300.	0.5	17
220	Bryophytes: Liverworts, Mosses, and Hornworts: Extraction and Isolation Procedures. Methods in Molecular Biology, 2013, 1055, 1-20.	0.9	17
221	Trifarienols A and B, Isolated from the liverwort <i>Cheilolejeunea trifaria</i> . Sesquiterpenes having a new carbon skeleton, trifarane. Tetrahedron Letters, 1994, 35, 4787-4788.	1.4	16
222	Sesquiterpenes and a phenolic compound from the liverwort <i>Omphalanthus filiformis</i> . Phytochemistry, 1995, 38, 651-653.	2.9	16
223	Sesqui- and di-terpenoids from the liverwort <i>Jungermannia vulcanicola</i> . Phytochemistry, 1996, 42, 93-96.	2.9	16
224	New Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. Chemical and Pharmaceutical Bulletin, 1998, 46, 1184-1185.	1.3	16
225	Sesqui- and diterpenoids from <i>Ptilidium ciliare</i> and <i>Barbilophozia</i> species (liverworts). Phytochemistry, 1999, 51, 563-566.	2.9	16
226	A revision of the positive sign of the optical rotation and its maximum value of \pm -eudesmol. Phytochemistry, 1999, 52, 689-694.	2.9	16
227	Tyromycic Acids B ⁺ E, New Lanostane Triterpenoids from the Mushroom <i>Tyromyces fissilis</i> . Journal of Natural Products, 2004, 67, 148-151.	3.0	16
228	Penicillium sclerotiorum Catalyzes the Conversion of Herbertenediol into Its Dimers: Mastigophorenes A and B. Chemical and Pharmaceutical Bulletin, 2005, 53, 256-257.	1.3	16
229	Chemical constituents of the Vietnamese liverwort <i>Porella densifolia</i> . F ⁺ -toterap ⁻ A ⁻ c, 2010, 81, 659-661.	2.2	16
230	Studies on the Genus < i>Thysananthus</i>(Marchantiophyta, Lejeuneaceae) 3. Terpenoid Chemistry and Chemotaxonomy of Selected Species of < i>Thysananthus</i>and < i>Dendrolejeunea fruticosa</i>. Cryptogamie, Bryologie, 2011, 32, 199-209.	0.2	16
231	Phytochemical investigations and bioactivity evaluation of liverworts as a function of anti-inflammatory and antinociceptive properties in animal models. Pharmaceutical Biology, 2013, 51, 1008-1013.	2.9	16
232	Sesquiterpene derivatives and a norsesquiterpenoid from the liverworts <i>Riccardia crassa</i> and <i>Porella caespitans</i> var. <i>Setigera</i> . Phytochemistry, 1992, 32, 137-140.	2.9	15
233	Rearranged ent-eudesmane- and ent-eremophilane-type sesquiterpenoids from the liverwort <i>Frullania dilatata</i> . Phytochemistry, 1994, 37, 1317-1321.	2.9	15
234	Sesquiterpenoids from the liverwort <i>dicranolejeunea yoshinaga</i> (HATT.) MIZUT.. Chemical and Pharmaceutical Bulletin, 1995, 43, 714-716.	1.3	15

#	ARTICLE	IF	CITATIONS
235	Structure of conocephalenol, a sesquiterpenoid alcohol from the European liverwort <i>Conocephalum conicum</i> : determination of the absolute configuration by total synthesis. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1995, , 593.	0.9	15
236	Isolation, structure, and synthesis of chenopodanol and the absolute configuration of chenopodene and chenopodanol. <i>Canadian Journal of Chemistry</i> , 1997, 75, 634-640.	1.1	15
237	First synthesis of 1,9-dideoxyforskolin from ptychantin A. <i>Tetrahedron Letters</i> , 2003, 44, 2305-2306.	1.4	15
238	Tyromycic Acids F and G: Two New Triterpenoids from the Mushroom <i>Tyromyces fissionis</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1441-1443.	1.3	15
239	Constituents of Isotachis aubertii (Isotachidaceae) collected in Madagascar. <i>Biochemical Systematics and Ecology</i> , 2004, 32, 1073-1078.	1.3	15
240	Inhibitory activity of nitric oxide production in RAW 264.7 cells of daldinal A-C from the fungus <i>Daldinia childiae</i> and other metabolites isolated from inedible mushrooms. <i>Journal of Natural Medicines</i> , 2006, 60, 303-307.	2.3	15
241	Insecticidal Constituents from the Argentine Liverwort <i>Plagiochila bursata</i> . <i>Chemistry and Biodiversity</i> , 2010, 7, 1855-1861.	2.1	15
242	Terpenoids and Bibenzyls from Three Argentine Liverworts. <i>Molecules</i> , 2011, 16, 10471-10478.	3.8	15
243	Terpenoids and Aromatic Compounds from Bryophytes and their Central Nervous System Activity. <i>Current Organic Chemistry</i> , 2020, 24, 113-128.	1.6	15
244	Bis(bibenzyl) ethers from <i>Pellia endiviifolia</i> . <i>Phytochemistry</i> , 1991, 30, 1523-1530.	2.9	14
245	Trifarienols A-E, trifarane-type sesquiterpenoids from the Malaysian liverwort <i>Cheilolejeunea trifaria</i> . <i>Phytochemistry</i> , 1995, 40, 171-176.	2.9	14
246	Sesquiterpenoids from a Cell Suspension Culture of the Liverwort <i>Porella vernicosa</i> Lindb. , 1996, 11, 53-56.		14
247	A phenethyl glycoside from <i>Conocephalum conicum</i> . <i>Phytochemistry</i> , 1996, 43, 1087-1088.	2.9	14
248	A Novel Skeletal Diterpenoid from the German Liverwort <i>Barbilophozia hatcheri</i> (Evans) Loeske.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 138-139.	1.3	14
249	New Chroman Derivatives from the Japanese Liverworts <i>Metacalypogeia cordifolia</i> and <i>Cephalozia otaruensis</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2001, 49, 924-926.	1.3	14
250	Occurrence of a High Concentration of Spider Pheromones in the Ascomycete Fungus <i>Hypoxylon truncatum</i> . <i>Journal of Natural Products</i> , 2003, 66, 1613-1614.	3.0	14
251	Cytotoxic bibenzyls, and germacrane- and pinguisane-type sesquiterpenoids from Indonesian, Tahitian and Japanese liverworts. <i>Natural Product Communications</i> , 2011, 6, 303-9.	0.5	14
252	Chemosystematics of <i>Porella</i> (Marchantiophyta, Porellaceae). <i>Natural Product Communications</i> , 2011, 6, 315-21.	0.5	14

#	ARTICLE	IF	CITATIONS
253	Fusicorragatol from the Venezuelan liverwort <i>Plagiochila corrugata</i> .. Chemical and Pharmaceutical Bulletin, 1994, 42, 2650-2652.	1.3	13
254	Africane- and monocyclofarnesane-type sesquiterpenoids from the liverwort <i>Porella subobtusa</i> . Phytochemistry, 1996, 43, 1285-1291.	2.9	13
255	Two novel skeletal diterpenoids, neodenudatenones A and B, from the liverwort <i>Odontoschisma denudatum</i> . Tetrahedron Letters, 1998, 39, 3791-3794.	1.4	13
256	Synthesis of cryptoporic acid A methyl ester. Tetrahedron Letters, 2000, 41, 3099-3102.	1.4	13
257	Activation of p38 Mitogen-Activated Protein Kinase duringent-11 β -Hydroxy-16-kauren-15-one-Induced Apoptosis in Human Leukemia HL-60 Cells. Planta Medica, 2005, 71, 275-277.	1.3	13
258	Chemical Constituents of Malagasy Liverworts. 6. A Myltaylane Caffeate with Nitric Oxide Inhibitory Activity from <i>Bazzania nitida</i> . Journal of Natural Products, 2007, 70, 856-858.	3.0	13
259	Volatile components from selected Tahitian liverworts. Natural Product Communications, 2009, 4, 1387-92.	0.5	13
260	Distribution of drimane sesquiterpenoids and tocopherols in liverworts, ferns and higher plants: Polygonaceae, Canellaceae and Winteraceae species. Natural Product Communications, 2012, 7, 685-92.	0.5	13
261	Bisbibenzyl and sesquiterpenoids from the liver wort <i>Jubula japonica</i> . Phytochemistry, 1993, 34, 1135-1137.	2.9	12
262	Sesquiterpene esters from the liverwort <i>Plagiochila poreloides</i> . Phytochemistry, 1994, 37, 1091-1093.	2.9	12
263	Synthesis and the absolute configuration of the sesquiterpene aldehyde tridensenol from the Taiwanese liverwort <i>Bazzania tridens</i> . Journal of the Chemical Society Perkin Transactions 1, 1995, , 1513.	0.9	12
264	Nudenoic acid: A novel tricyclic sesquiterpenoid from the Taiwanese liverwort <i>Mylia nuda</i> . Tetrahedron Letters, 1996, 37, 9307-9308.	1.4	12
265	Bazzanane Sesquiterpenoids from the New Zealand Liverwort <i>Frullania falciloba</i> . Chemical and Pharmaceutical Bulletin, 2006, 54, 1347-1349.	1.3	12
266	New ent-Verticillane Diterpenoids from the Japanese Liverwort <i>Jackiella javanica</i> . Chemical and Pharmaceutical Bulletin, 2008, 56, 1184-1188.	1.3	12
267	Evaluation of anti-melanoma and tyrosinase inhibitory properties of marchantin A, a natural macrocyclic bisbibenzyl isolated from <i>Marchantia</i> species. Phytochemistry Letters, 2019, 31, 192-195.	1.2	12
268	Setiformenol, isolated from the liverwort <i>tetralophozia setiformis</i> , the first example of cembrane-type diterpene from bryophytes. Tetrahedron Letters, 1993, 34, 643-644.	1.4	11
269	Naphthalene and isocoumarin derivatives from the liverwort <i>Wettsteinia schusterana</i> . Phytochemistry, 1994, 37, 233-235.	2.9	11
270	A New Monocyclic Diterpene from the Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. Chemical and Pharmaceutical Bulletin, 1996, 44, 1628-1630.	1.3	11

#	ARTICLE	IF	CITATIONS
271	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia rotundata</i> . <i>Phytochemistry</i> , 1997, 44, 653-657.	2.9	11
272	Isolation and Structure of Striatenic Acid from Liverwort <i>Cheilolejeunea serpentina</i> and the Absolute Configuration by Synthesis. <i>Tetrahedron</i> , 2000, 56, 1655-1659.	1.9	11
273	Hydnellins A and B, Nitrogen-Containing Terphenyls from the Mushrooms <i>Hydnellum suaveolens</i> and <i>Hydnellum geogericum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 912-914.	1.3	11
274	Volatile Components from Selected Tahitian Liverworts. <i>Natural Product Communications</i> , 2009, 4, 1934578X0900401.	0.5	11
275	Volatile Components of the Stressed Liverwort <i>Conocephalum Conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	11
276	Chemotypes and Biomarkers of Seven Species of New Caledonian Liverworts from the Bazzanioideae Subfamily. <i>Molecules</i> , 2018, 23, 1353.	3.8	11
277	Phytochemicals from bryophytes: Structures and biological activity. <i>Journal of the Serbian Chemical Society</i> , 2021, 86, 1139-1175.	0.8	11
278	Sesquiterpenoids from the liverwort <i>Marsupella aquatica</i> . <i>Phytochemistry</i> , 1994, 37, 777-779.	2.9	10
279	Ent-longipinane-type sesquiterpenoid from the liverwort <i>Marsupella emarginata</i> . <i>Phytochemistry</i> , 1994, 37, 1767-1768.	2.9	10
280	Clerodane-type Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. <i>Chemical and Pharmaceutical Bulletin</i> , 2000, 48, 1818-1821.	1.3	10
281	Sesquiterpene hydrocarbons with trifarane backbone in the liverwort <i>Trocholejeunea sandvicensis</i> . <i>Phytochemistry</i> , 2001, 57, 499-506.	2.9	10
282	Expedient Synthetic Transformation of Ptychantins into Forskolin. <i>Synlett</i> , 2008, 2008, 929-931.	1.8	10
283	Chemical Constituents of Selected Japanese and New Zealand Liverworts. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.5	10
284	Liverworts-Potential Source of Medicinal Compounds. , 2012, 01, .		10
285	Identification of sesquiterpene lactones in the Bryophyta (mosses) <i>Takakia</i> : <i>Takakia</i> species are closely related chemically to the Marchantiophyta (liverworts). <i>Natural Product Communications</i> , 2015, 10, 5-8.	0.5	10
286	Volatile Components of the Stressed Liverwort <i>Conocephalum conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 103-4.	0.5	10
287	New Highly Oxygenated 6,7-Seco-kaurane- and Bis-kaurane-Type Diterpenoids from The Liverwort <i>Jungermannia exsertifolia</i> STEPH. ssp. <i>cordifolia</i> (DUM.) VANA.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2656-2658.	1.3	9
288	Structures of Four Novel Highly Oxygenated Labdane-Type Diterpenoids, Ptychantins F-I, from the Liverwort <i>Ptychanthus striatus</i> . <i>Chemistry Letters</i> , 1995, 24, 481-482.	1.3	9

#	ARTICLE	IF	CITATIONS
289	secocuparane-type sesquiterpenoid from the Japanese liverwort <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 2001, 56, 807-810.	2.9	9
290	A Novel Dimeric Lactone Bis-osmundalactone from the Japanese Inedible Mushroom <i>Paxillus atromentosus</i> var. <i>bambusinus</i> . <i>Heterocycles</i> , 2002, 56, 581.	0.7	9
291	Sesqui- and diterpenoids from three New Zealand liverworts: <i>Bazzania novae-zelandiae</i> , <i>Gackstroemiasp.</i> and <i>Dendromastigophorasp.</i> <i>Natural Product Research</i> , 2010, 24, 68-75.	1.8	9
292	Diversity of Secondary Metabolites in the Liverwort <i>< i>Syzygiella rubricaulis</i></i> (<i>< sc>Nees</sc></i> <i>< sc>Stephani</sc></i>) (<i>Jamesoniellaceae, Marchantiophyta</i>) from Neotropical High Mountains. <i>Chemistry and Biodiversity</i> , 2018, 15, e1800239.	2.1	9
293	Daurichromenic acid-producing oxidocyclase in the young leaves of <i>Rhododendron dauricum</i> . <i>Natural Product Communications</i> , 2014, 9, 1329-32.	0.5	9
294	Malagasy Liverworts, Source of New and Biologically Active Compounds. <i>Natural Product Communications</i> , 2007, 2, 1934578X0700200.	0.5	8
295	Pungent Aromatic Compound from New Zealand Liverwort <i>Hymenophyton flabellatum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2009, 57, 1015-1018.	1.3	8
296	Cytotoxic Bibenzyls, and Germacrane- and Pinguisane-type Sesquiterpenoids from Indonesian, Tahitian and Japanese Liverworts. <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.5	8
297	Chemosystematics of selected liverworts collected in Borneo . <i>Bryophyte Diversity and Evolution</i> , 2015, 31, 33.	1.1	8
298	Bryophytes: chemical diversity, synthesis and biotechnology. A review.. <i>Flavour and Fragrance Journal</i> , 2011, 26, n/a-n/a.	2.6	7
299	Pungent and Bitter, Cytotoxic and Antiviral Terpenoids from Some Bryophytes and Inedible Fungi. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.5	7
300	Phytochemistry of three selected liverworts: <i>Conocephalum conicum</i> , <i>Plagiochila barteri</i> and <i>P. terebrans</i> . <i>Arkivoc</i> , 2007, 2007, 22-29.	0.5	7
301	Recent Advances of Biologically Active Substances from the Marchantiophyta. <i>Natural Product Communications</i> , 2008, 3, 1934578X0800300.	0.5	6
302	Biotransformation of Sesquiterpenoids. , 2010, , 803-892.		6
303	Chemosystematics of <i>< i>Porella</i></i> (Marchantiophyta, Porellaceae). <i>Natural Product Communications</i> , 2011, 6, 1934578X1100600.	0.5	6
304	Constituents of the Argentine Liverwort <i>< i>Plagiochila diversifolia</i></i> and Their Insecticidal Activities. <i>Chemistry and Biodiversity</i> , 2017, 14, e1700229.	2.1	6
305	Transcriptome Analysis of Marchantin Biosynthesis from the Liverwort <i>< i>Marchantia polymorpha</i></i> . <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	6
306	Characteristic Scent from the Tahitian Liverwort, <i>Cyathodium foetidissimum</i>. <i>Journal of Oleo Science</i> , 2018, 67, 1265-1269.	1.4	6

#	ARTICLE	IF	CITATIONS
307	Syntheses of isocitric acid derivatives and biological evaluation. <i>Tetrahedron Letters</i> , 2000, 41, 3095-3098.	1.4	5
308	Pregnane-Type Steroids from the Inedible Mushroom <i>Thelephora terrestris</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 309-312.	1.3	5
309	Volatile Components of Selected Liverworts, and Cytotoxic, Radical Scavenging and Antimicrobial Activities of Their Crude Extracts. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.5	5
310	Distribution of Drimane Sesquiterpenoids and Tocopherols in Liverworts, Ferns and Higher Plants: Polygonaceae, Canellaceae and Winteraceae Species. <i>Natural Product Communications</i> , 2012, 7, 1934578X1200700.	0.5	5
311	Chemical variability of the Tahitian <i>Marchantia hexaptera</i> Reich.. <i>Phytochemistry Letters</i> , 2014, 10, xcix-ciii.	1.2	5
312	Biotransformation of Perrottetin F by <i>Aspergillus niger</i> : New Bioactive Secondary Metabolites. <i>Records of Natural Products</i> , 2021, 15, 281-292.	1.3	5
313	Volatile Components Emitted from the Liverwort <i>Marchantia paleacea</i> subsp. <i>diptera</i> . <i>Natural Product Communications</i> , 2016, 11, 263-4.	0.5	5
314	Chemical Relationships between Liverworts of the Family Lejeuneaceae (Porellales,) Tj ETQq0 0 0 rgBT /Overlock 10 _{0.5} Tf 50 462 ₄ Td (Jungener)		
315	Volatile Components Emitted from the Liverwort <i>Marchantia Paleacea</i> Subsp. <i>Diptera</i>. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	4
316	Management of Diabetic Bacterial Foot Infections with Organic Extracts of Liverwort <i>Marchantia debilis</i> from Cameroon. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	4
317	Chemotaxonomy and cytotoxicity of the liverwort <i>Porella Viridissima</i>. <i>Natural Product Research</i> , 2021, 35, 2099-2102.	1.8	4
318	Biotransformation of Sesquiterpenoids from Liverworts by Fungi and Mammals. <i>Natural Product Communications</i> , 2010, 5, 1934578X1000500.	0.5	3
319	Chemical Constituents of Marchantiophyta. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 25-561.	1.1	3
320	Influence of Thermal Processing and <i>in vitro</i> Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	3
321	Microbial Transformation of Some Natural and Synthetic Aromatic Compounds by Fungi: <i>Aspergillus</i> and <i>Neurospora</i> Strains. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	3
322	Chemo- and biocatalytic esterification of marchantin A and cytotoxic activity of ester derivatives. FÄ-toterapÄ-Ä¢, 2020, 142, 104520.	2.2	3
323	Chemical Diversity of Liverworts From<i>Frullania</i> Genus. <i>Natural Product Communications</i> , 2021, 16, 1934578X2199538.	0.5	3
324	Hunting for bis-bibenzyls in <i>Primula veris</i> subsp. <i>macrocalyx</i> (Bunge) LÄ¼di: Organ-specific accumulation and cytotoxic activity. <i>Phytochemistry Letters</i> , 2021, 44, 90-97.	1.2	3

#	ARTICLE	IF	CITATIONS
325	Modification of valencene by bio- and chemical transformation. <i>Natural Product Communications</i> , 2013, 8, 859-62.	0.5	3
326	Chemical Relationships Between Algae, Bryophytes, and Pteridophytes. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 705-726.	1.1	2
327	GC/MS Fingerprinting of Solvent Extracts and Essential Oils Obtained from Liverwort Species. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	2
328	Biotransformation of sesquiterpenoids from liverworts by fungi and mammals. <i>Natural Product Communications</i> , 2010, 5, 695-707.	0.5	2
329	Influence of Thermal Processing and in vitro Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. <i>Natural Product Communications</i> , 2016, 11, 1153-1156.	0.5	2
330	Global phytochemistry: research in Japan. <i>Phytochemistry</i> , 2003, 64, 909-912.	2.9	1
331	Chemosystematics of the Hepaticae. <i>ChemInform</i> , 2004, 35, no.	0.0	1
332	Guy Ourisson 1926â€“2006. <i>Phytochemistry</i> , 2007, 68, 1350-1351.	2.9	1
333	Biologically Active Compounds of the Marchantiophyta and Bryophyta. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 619-638.	1.1	1
334	Chemical Diversity of Bryophytes. <i>Progress in the Chemistry of Organic Natural Products</i> , 2013, , 21-24.	1.1	1
335	Chemosystematics of the Thai Liverwort <i>Cheilolejeunea</i> (Marchantiophyta, Lejeuneaceae). <i>Natural Product Communications</i> , 2013, 8, 1934578X1300800.	0.5	1
336	Chemical Evidence for the Liverwort Complex, Chiloscyphus concavus and C. horizontalis. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.5	1
337	Bis-benzyls from the Cameroon Liverwort <i>Marchantia debilis</i>. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	1
338	Terpenoids, Flavonoids and Acetogenins from Some Malagasy Plants. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	1
339	Dimeric and esterified sesquiterpenes from the liverwort Chiastocaulon caledonicum. <i>Phytochemistry</i> , 2020, 179, 112495.	2.9	1
340	Biotransformation of Sesquiterpenoids, Ionones, Damascones, Adamantanes, and Aromatic Compounds by Green Algae, Fungi, and Mammals., 2009, , 737-841.		1
341	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 412.	1.5	0
342	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 410-411.	1.5	0

#	ARTICLE	IF	CITATIONS
343	Chemosystematics of Marchantiophyta. Progress in the Chemistry of Organic Natural Products, 2013, , 639-704.	1.1	0
344	Sesqui- and Diterpenoids from Tahitian and Japanese Liverworts <i>Jungermannia</i> species. Natural Product Communications, 2017, 12, 1934578X1701200.	0.5	0
345	Heterocyclic Stilbene and Bibenzyl Derivatives in Liverworts: Distribution, Structures, Total Synthesis and Biological Activity. Heterocycles, 2022, 105, 179.	0.7	0