

# Yoshinori Asakawa

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3280072/publications.pdf>

Version: 2024-02-01

345  
papers

9,745  
citations

47006  
47  
h-index

102487  
66  
g-index

359  
all docs

359  
docs citations

359  
times ranked

4029  
citing authors

#	ARTICLE	IF	CITATIONS
1	Bis-bibenzyls, Bibenzyls, and Terpenoids in 33 Genera of the Marchantiophyta (Liverworts): Structures, Synthesis, and Bioactivity. <i>Journal of Natural Products</i> , 2022, 85, 729-762.	3.0	21
2	Heterocyclic Stilbene and Bibenzyl Derivatives in Liverworts: Distribution, Structures, Total Synthesis and Biological Activity. <i>Heterocycles</i> , 2022, 105, 179.	0.7	0
3	Chemotaxonomy and cytotoxicity of the liverwort <i>&lt; i&gt;Porella Viridissima&lt;/i&gt;</i> . <i>Natural Product Research</i> , 2021, 35, 2099-2102.	1.8	4
4	Chemical Diversity of Liverworts From <i>&lt; i&gt;Frullania&lt;/i&gt;</i> Genus. <i>Natural Product Communications</i> , 2021, 16, 1934578X2199538.	0.5	3
5	Biotransformation of Perrottetin F by <i>Aspergillus niger</i> : New Bioactive Secondary Metabolites. <i>Records of Natural Products</i> , 2021, 15, 281-292.	1.3	5
6	Hunting for bis-bibenzyls in <i>Primula veris</i> subsp. <i>macrocalyx</i> (Bunge) LÄ¼di: Organ-specific accumulation and cytotoxic activity. <i>Phytochemistry Letters</i> , 2021, 44, 90-97.	1.2	3
7	Phytochemicals from bryophytes: Structures and biological activity. <i>Journal of the Serbian Chemical Society</i> , 2021, 86, 1139-1175.	0.8	11
8	Dimeric and esterified sesquiterpenes from the liverwort <i>Chiastocalon caledonicum</i> . <i>Phytochemistry</i> , 2020, 179, 112495.	2.9	1
9	Terpenoids and Aromatic Compounds from Bryophytes and their Central Nervous System Activity. <i>Current Organic Chemistry</i> , 2020, 24, 113-128.	1.6	15
10	Distribution of Bibenzyls, Prenyl Bibenzyls, Bis-bibenzyls, and Terpenoids in the Liverwort Genus <i>&lt; i&gt;Radula&lt;/i&gt;</i> . <i>Journal of Natural Products</i> , 2020, 83, 756-769.	3.0	33
11	Chemo- and biocatalytic esterification of marchantin A and cytotoxic activity of ester derivatives. <i>FÄ-toterapÄ-Äc</i> , 2020, 142, 104520.	2.2	3
12	Bryophytes as a source of bioactive volatile terpenoids – A review. <i>Food and Chemical Toxicology</i> , 2019, 132, 110649.	3.6	52
13	Cytotoxic Activity of Riccardin and Perrottetin Derivatives from the Liverwort <i>&lt; i&gt;Lunularia cruciata&lt;/i&gt;</i> . <i>Journal of Natural Products</i> , 2019, 82, 694-701.	3.0	22
14	Evaluation of anti-melanoma and tyrosinase inhibitory properties of marchantin A, a natural macrocyclic bisbibenzyl isolated from <i>Marchantia</i> species. <i>Phytochemistry Letters</i> , 2019, 31, 192-195.	1.2	12
15	Chemical Constituents of Bryophytes: Structures and Biological Activity. <i>Journal of Natural Products</i> , 2018, 81, 641-660.	3.0	141
16	Characteristic Scent from the Tahitian Liverwort, <i>&amp;lt;i&amp;gt;Cyathodium foetidissimum&amp;lt;/i&amp;gt;</i> . <i>Journal of Oleo Science</i> , 2018, 67, 1265-1269.	1.4	6
17	Diversity of Secondary Metabolites in the Liverwort <i>&lt; i&gt;Syzygiella rubricaulis&lt;/i&gt;</i> ( <i>&lt; sc&gt;Nees&lt;/sc&gt;</i> ) <i>&lt; sc&gt;Stephani&lt;/sc&gt;</i> (Jamesoniellaceae, Marchantiophyta) from Neotropical High Mountains. <i>Chemistry and Biodiversity</i> , 2018, 15, e1800239.	2.1	9
18	Chemotypes and Biomarkers of Seven Species of New Caledonian Liverworts from the Bazzanioideae Subfamily. <i>Molecules</i> , 2018, 23, 1353.	3.8	11

#	ARTICLE	IF	CITATIONS
19	MAO-A Inhibitory Potential of Terpene Constituents from Ginger Rhizomes—A Bioactivity Guided Fractionation. <i>Molecules</i> , 2018, 23, 1301.	3.8	21
20	An Aromatic Farnesyltransferase Functions in Biosynthesis of the Anti-HIV Meroterpenoid Daurichromenic Acid. <i>Plant Physiology</i> , 2018, 178, 535-551.	4.8	23
21	Constituents of the Argentine Liverwort <i>&lt; i&gt;Plagiochila diversifolia&lt;/i&gt;</i> and Their Insecticidal Activities. <i>Chemistry and Biodiversity</i> , 2017, 14, e1700229.	2.1	6
22	Identification and Characterization of Daurichromenic Acid Synthase Active in Anti-HIV Biosynthesis. <i>Plant Physiology</i> , 2017, 174, 2213-2230.	4.8	25
23	Sesqui- and Diterpenoids from Tahitian and Japanese Liverworts <i>&lt; i&gt;Jungermannia&lt;/i&gt;</i> species. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	0
24	Microbial Transformation of Some Natural and Synthetic Aromatic Compounds by Fungi: <i>Aspergillus</i> and <i>Neurospora</i> Strains. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	3
25	Transcriptome Analysis of Marchantin Biosynthesis from the Liverwort <i>&lt; i&gt;Marchantia polymorpha&lt;/i&gt;</i> . <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	6
26	GC/MS Fingerprinting of Solvent Extracts and Essential Oils Obtained from Liverwort Species. <i>Natural Product Communications</i> , 2017, 12, 1934578X1701200.	0.5	2
27	Application of Chromatographic and Spectroscopic Methods towards the Quality Assessment of Ginger ( <i>Zingiber officinale</i> ) Rhizomes from Ecological Plantations. <i>International Journal of Molecular Sciences</i> , 2017, 18, 452.	4.1	42
28	Volatile Components Emitted from the Liverwort <i>&lt; i&gt;Marchantia Paleacea&lt;/i&gt;</i> Subsp. <i>&lt; i&gt;Diptera&lt;/i&gt;</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	4
29	Bis-benzyls from the Cameroon Liverwort <i>&lt; i&gt;Marchantia debilis&lt;/i&gt;</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	1
30	Terpenoids, Flavonoids and Acetogenins from Some Malagasy Plants. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	1
31	A Novel Class of Plant Type III Polyketide Synthase Involved in Orsellinic Acid Biosynthesis from <i>Rhododendron dauricum</i> . <i>Frontiers in Plant Science</i> , 2016, 7, 1452.	3.6	34
32	Volatile Components of the Stressed Liverwort <i>Conocephalum Conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	11
33	Management of Diabetic Bacterial Foot Infections with Organic Extracts of Liverwort <i>&lt; i&gt;Marchantia debilis&lt;/i&gt;</i> from Cameroon. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	4
34	Influence of Thermal Processing and <i>&lt; i&gt;in vitro&lt;/i&gt;</i> Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. <i>Natural Product Communications</i> , 2016, 11, 1934578X1601100.	0.5	3
35	Dietary intake of metals by the young adult population of Eastern Poland: Results from a market basket study. <i>Journal of Trace Elements in Medicine and Biology</i> , 2016, 35, 36-42.	3.0	27
36	Volatile Components of the Stressed Liverwort <i>Conocephalum conicum</i> . <i>Natural Product Communications</i> , 2016, 11, 103-4.	0.5	10

#	ARTICLE	IF	CITATIONS
37	Volatile Components Emitted from the Liverwort <i>Marchantia paleacea</i> subsp. <i>diptera</i> . Natural Product Communications, 2016, 11, 263-4.	0.5	5
38	Influence of Thermal Processing and in vitro Digestion on the Antioxidant Potential of Ginger and Ginger Containing Products. Natural Product Communications, 2016, 11, 1153-1156.	0.5	2
39	Chemotaxonomic value of essential oil components in liverwort species. A review. Flavour and Fragrance Journal, 2015, 30, 189-196.	2.6	18
40	<strong>Chemosystematics of selected liverworts collected in Borneo</strong>. Bryophyte Diversity and Evolution, 2015, 31, 33.	1.1	8
41	Identification of sesquiterpene lactones in the Bryophyta (mosses) Takakia: Takakia species are closely related chemically to the Marchantiophyta (liverworts). Natural Product Communications, 2015, 10, 5-8.	0.5	10
42	Pungent and Bitter, Cytotoxic and Antiviral Terpenoids from Some Bryophytes and Inedible Fungi. Natural Product Communications, 2014, 9, 1934578X1400900.	0.5	7
43	Chemical Evidence for the Liverwort Complex, <i>Chiloscyphus concavus</i> and <i>C. horizontalis</i> . Natural Product Communications, 2014, 9, 1934578X1400900.	0.5	1
44	Fingerprinting of Secondary Metabolites of Liverworts: Chemosystematic Approach. Journal of AOAC INTERNATIONAL, 2014, 97, 1234-1243.	1.5	22
45	Chemical variability of the Tahitian <i>Marchantia hexaptera</i> Reich.. Phytochemistry Letters, 2014, 10, xcix-ciii.	1.2	5
46	Daurichromenic acid-producing oxidocyclase in the young leaves of <i>Rhododendron dauricum</i> . Natural Product Communications, 2014, 9, 1329-32.	0.5	9
47	Total synthesis of riccardin C and ( $\Delta\pm$ )-cavicularin via Pd-catalyzed Ar–Ar cross couplings. Tetrahedron, 2013, 69, 6959-6968.	1.9	20
48	Bryophytes: Liverworts, Mosses, and Hornworts: Extraction and Isolation Procedures. Methods in Molecular Biology, 2013, 1055, 1-20.	0.9	17
49	Identification of cryptic species within liverwort <i>Conocephalum conicum</i> based on the volatile components. Phytochemistry, 2013, 95, 234-241.	2.9	27
50	Biologically Active Compounds of the Marchantiophyta and Bryophyta. Progress in the Chemistry of Organic Natural Products, 2013, , 619-638.	1.1	1
51	Introduction. Progress in the Chemistry of Organic Natural Products, 2013, 95, 1-16.	1.1	77
52	Chemical Diversity of Bryophytes. Progress in the Chemistry of Organic Natural Products, 2013, , 21-24.	1.1	1
53	Chemical Constituents of Marchantiophyta. Progress in the Chemistry of Organic Natural Products, 2013, , 25-561.	1.1	3
54	Chemosystematics of Marchantiophyta. Progress in the Chemistry of Organic Natural Products, 2013, , 639-704.	1.1	0

#	ARTICLE	IF	CITATIONS
55	Chemical Relationships Between Algae, Bryophytes, and Pteridophytes. Progress in the Chemistry of Organic Natural Products, 2013, , 705-726.	1.1	2
56	Chemical Constituents of Bryophytes. Progress in the Chemistry of Organic Natural Products, 2013, , .	1.1	50
57	Phytochemical and biological studies of bryophytes. Phytochemistry, 2013, 91, 52-80.	2.9	199
58	Phytochemical investigations and bioactivity evaluation of liverworts as a function of anti-inflammatory and antinociceptive properties in animal models. Pharmaceutical Biology, 2013, 51, 1008-1013.	2.9	16
59	Chemosystematics of the Thai Liverwort <i>Cheilolejeunea</i> (Marchantiophyta, Lejeuneaceae). Natural Product Communications, 2013, 8, 1934578X1300800.	0.5	1
60	Chemical Relationships between Liverworts of the Family Lejeuneaceae (Porellales,) Tj ETQq0 0 0 rgBT /Overlock 10 <sub>0.5</sub> Tf 50 542 <sub>4</sub> Td (Junges)		
61	Modification of valencene by bio- and chemical transformation. Natural Product Communications, 2013, 8, 859-62.	0.5	3
62	Distribution of Cyclic and Acyclic Bis-bibenzyls in the Marchantiophyta (Liverworts), Ferns and Higher Plants and Their Biological Activities, Biosynthesis, and Total Synthesis. Heterocycles, 2012, 86, 891.	0.7	32
63	Distribution of Drimane Sesquiterpenoids and Tocopherols in Liverworts, Ferns and Higher Plants: Polygonaceae, Canellaceae and Winteraceae Species. Natural Product Communications, 2012, 7, 1934578X1200700.	0.5	5
64	Liverworts-Potential Source of Medicinal Compounds. , 2012, 01, .		10
65	In vitro antitrypanosomal activity of bis(bibenzyls)s and bibenzyls from liverworts against Trypanosoma brucei. Journal of Natural Medicines, 2012, 66, 377-382.	2.3	26
66	Distribution of drimane sesquiterpenoids and tocopherols in liverworts, ferns and higher plants: Polygonaceae, Canellaceae and Winteraceae species. Natural Product Communications, 2012, 7, 685-92.	0.5	13
67	Studies on the Genus<i>Thysananthus</i> (Marchantiophyta, Lejeuneaceae) 3. Terpenoid Chemistry and Chemotaxonomy of Selected Species of<i>Thysananthus</i> and<i>Dendrolejeunea fruticosa</i>. Cryptogamie, Bryologie, 2011, 32, 199-209.	0.2	16
68	Cytotoxic Bibenzyls, and Germacrane- and Pinguisane-type Sesquiterpenoids from Indonesian, Tahitian and Japanese Liverworts. Natural Product Communications, 2011, 6, 1934578X1100600.	0.5	8
69	Chemosystematics of <i>Porella</i> (Marchantiophyta, Porellaceae). Natural Product Communications, 2011, 6, 1934578X1100600.	0.5	6
70	Terpenoids and Bibenzyls from Three Argentine Liverworts. Molecules, 2011, 16, 10471-10478.	3.8	15
71	In vitro antitrypanosomal activity of plant terpenes against Trypanosoma brucei. Phytochemistry, 2011, 72, 2024-2030.	2.9	57
72	Vasorelaxant effects of macrocyclic bis(bibenzyls) from liverworts. Bioorganic and Medicinal Chemistry, 2011, 19, 4051-4056.	3.0	20

#	ARTICLE	IF	CITATIONS
73	Bryophytes: chemical diversity, synthesis and biotechnology. A review.. Flavour and Fragrance Journal, 2011, 26, n/a-n/a.	2.6	7
74	Anti-Influenza Activity of Marchantins, Macroyclic Bisbibenzyls Contained in Liverworts. PLoS ONE, 2011, 6, e19825.	2.5	73
75	Cytotoxic bibenzyls, and germacrane- and pinguisane-type sesquiterpenoids from Indonesian, Tahitian and Japanese liverworts. Natural Product Communications, 2011, 6, 303-9.	0.5	14
76	Chemosystematics of Porella (Marchantiophyta, Porellaceae). Natural Product Communications, 2011, 6, 315-21.	0.5	14
77	Cytotoxic, radical scavenging and antimicrobial activities of sesquiterpenoids from the Tahitian liverwort <i>Mastigophora diclados</i> (Brid.) Nees (Mastigophoraceae). Journal of Natural Medicines, 2010, 64, 417-422.	2.3	40
78	Chemical constituents of the Vietnamese liverwort <i>Porella densifolia</i> . FÄtotterapÄ, 2010, 81, 659-661.	2.2	16
79	Insecticidal Constituents from the Argentine Liverwort <i>Plagiochila bursata</i> . Chemistry and Biodiversity, 2010, 7, 1855-1861.	2.1	15
80	Zierane sesquiterpene lactone, cembrane and fusicoccane diterpenoids, from the Tahitian liverwort <i>Chandonanthus hirtellus</i> . Phytochemistry, 2010, 71, 1387-1394.	2.9	39
81	Volatile Components of Selected Liverworts, and Cytotoxic, Radical Scavenging and Antimicrobial Activities of Their Crude Extracts. Natural Product Communications, 2010, 5, 1934578X1000500.	0.5	5
82	Biotransformation of Sesquiterpenoids from Liverworts by Fungi and Mammals. Natural Product Communications, 2010, 5, 1934578X1000500.	0.5	3
83	Biotransformation of Sesquiterpenoids., 2010, , 803-892.		6
84	Sesqui- and diterpenoids from three New Zealand liverworts: <i>Bazzania novae-zelandiae</i> , <i>Gackstroemiasp.</i> and <i>Dendromastigophorasp.</i> . Natural Product Research, 2010, 24, 68-75.	1.8	9
85	Biotransformation of Monoterpenoids. , 2010, , 669-801.		20
86	Biotransformation of sesquiterpenoids from liverworts by fungi and mammals. Natural Product Communications, 2010, 5, 695-707.	0.5	2
87	Volatile Components from Selected Tahitian Liverworts. Natural Product Communications, 2009, 4, 1934578X0900401.	0.5	11
88	Antimitotic activity of two macrocyclic bis(bibenzyls), isoplagiochins A and B from the Liverwort <i>Plagiochila fruticosa</i> . Bioorganic and Medicinal Chemistry Letters, 2009, 19, 493-496.	2.2	37
89	Synthesis of riccardin C and its seven analogues. Part 1: The role of their phenolic hydroxy groups as LXRÎ± agonists. Bioorganic and Medicinal Chemistry Letters, 2009, 19, 738-741.	2.2	36
90	Bryophytes: Bio- and Chemical Diversity, Bioactivity and Chemosystematics. Heterocycles, 2009, 77, 99.	0.7	70

#	ARTICLE	IF	CITATIONS
91	Pungent Aromatic Compound from New Zealand Liverwort <i>Hymenophyton flabellatum</i> . Chemical and Pharmaceutical Bulletin, 2009, 57, 1015-1018.	1.3	8
92	Biotransformation of Sesquiterpenoids, Ionones, Damascones, Adamantanes, and Aromatic Compounds by Green Algae, Fungi, and Mammals., , 2009, , 737-841.		1
93	Volatile components from selected Tahitian liverworts. Natural Product Communications, 2009, 4, 1387-92.	0.5	13
94	Chapter Five: Distribution of Terpenoids and Aromatic Compounds in Selected Southern Hemispheric Liverworts. Fieldiana Botany, 2008, 47, 37.	0.3	31
95	Secondary Metabolites of <i>&lt; i&gt;Cinnamosma madagascariensis&lt;/i&gt;</i> and Their $\beta$ -Glucosidase Inhibitory Properties. Journal of Natural Products, 2008, 71, 123-126.	3.0	21
96	Insect Antifeedant Sesquiterpene Acetals from the Liverwort <i>&lt; i&gt;Lepidolaena clavigera&lt;/i&gt;</i> . 2. Structures, Artifacts, and Activity. Journal of Natural Products, 2008, 71, 258-261.	3.0	28
97	Expedient Synthetic Transformation of Ptychantins into Forskolin. Synlett, 2008, 2008, 929-931.	1.8	10
98	Marchantiophyta (Liverworts): Rich Sources of Macroyclic Bis(bibenzyls). Heterocycles, 2008, 76, 99.	0.7	29
99	New ent-Verticillane Diterpenoids from the Japanese Liverwort <i>Jackiella javanica</i> . Chemical and Pharmaceutical Bulletin, 2008, 56, 1184-1188.	1.3	12
100	The Novel Compounds That Activate Farnesoid X Receptor: the Diversity of Their Effects on Gene Expression. Journal of Pharmacological Sciences, 2008, 107, 285-294.	2.5	33
101	Recent Advances of Biologically Active Substances from the Marchantiophyta. Natural Product Communications, 2008, 3, 1934578X0800300.	0.5	6
102	Chemical Constituents of Selected Japanese and New Zealand Liverworts. Natural Product Communications, 2008, 3, 1934578X0800300.	0.5	10
103	Volatile Components from Selected Mexican, Ecuadorian, Greek, German and Japanese Liverworts. Natural Product Communications, 2008, 3, 1934578X0800300.	0.5	17
104	Cinnamacrins A $\beta$ C, Cinnafragrin D, and Cytostatic Metabolites with $\beta$ -Glucosidase Inhibitory Activity from <i>Cinnamosma macrocarpa</i> . Journal of Natural Products, 2007, 70, 277-282.	3.0	29
105	Chemical Constituents of Malagasy Liverworts. 6. A Myltaylane Caffeate with Nitric Oxide Inhibitory Activity from <i>Bazzania nitida</i> . Journal of Natural Products, 2007, 70, 856-858.	3.0	13
106	Biologically active compounds from bryophytes. Pure and Applied Chemistry, 2007, 79, 557-580.	1.9	175
107	Malagasy Liverworts, Source of New and Biologically Active Compounds. Natural Product Communications, 2007, 2, 1934578X0700200.	0.5	8
108	Guy Ourisson 1926–2006. Phytochemistry, 2007, 68, 1350-1351.	2.9	1

#	ARTICLE	IF	CITATIONS
109	Bioactive compounds from liverworts: Inhibition of lipopolysaccharide-induced inducible NOS mRNA in RAW 264.7 cells by Herbertenoids and Cuparenoids. <i>Phytomedicine</i> , 2007, 14, 486-491.	5.3	23
110	Phytochemistry of three selected liverworts: <i>Conocephalum conicum</i> , <i>Plagiochila barteri</i> and <i>P. terebrans</i> . <i>Arkivoc</i> , 2007, 2007, 22-29.	0.5	7
111	Synthetic Transformation of Ptychanin into Forskolin and 1,9-Dideoxyforskolin. <i>Journal of Organic Chemistry</i> , 2006, 71, 4619-4624.	3.2	24
112	Inhibition of Nitric Oxide Production in RAW 264.7 Cells by Azaphilones from Xylariaceous Fungi. <i>Biological and Pharmaceutical Bulletin</i> , 2006, 29, 34-37.	1.4	50
113	Chemical Constituents of Malagasy Liverworts, Part V: Prenyl Bibenzyls and Clerodane Diterpenoids with Nitric Oxide Inhibitory Activity from <i>Radula appressa</i> and <i>Thysananthus spathulistipus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 1046-1049.	1.3	36
114	Diterpenoids and Aromatic Compounds from the Three New Zealand Liverworts <i>Jamesoniella kirkii</i> , <i>Balantiopsis rosea</i> , and <i>Radula</i> Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 902-906.	1.3	22
115	Hydnellins A and B, Nitrogen-Containing Terphenyls from the Mushrooms <i>Hydnellum suaveolens</i> and <i>Hydnellum geogericum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 912-914.	1.3	11
116	Bazzanane Sesquiterpenoids from the New Zealand Liverwort <i>Frullania falciloba</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2006, 54, 1347-1349.	1.3	12
117	Inedible mushrooms: a good source of biologically active substances. <i>Chemical Record</i> , 2006, 6, 79-99.	5.8	74
118	Grifolin derivatives from <i>Albatrellus caeruleoporus</i> , new inhibitors of nitric oxide production in RAW 264.7 cells. <i>Bioorganic and Medicinal Chemistry</i> , 2006, 14, 164-168.	3.0	62
119	Chemical constituents of Malagasy liverworts: Cyclomyltaylanoids from <i>Bazzania madagassa</i> . <i>Phytochemistry</i> , 2006, 67, 2616-2622.	2.9	17
120	Inhibitory activity of nitric oxide production in RAW 264.7 cells of daldinals A-C from the fungus <i>Daldinia childiae</i> and other metabolites isolated from inedible mushrooms. <i>Journal of Natural Medicines</i> , 2006, 60, 303-307.	2.3	15
121	Changes in secondary metabolism during stromatal ontogeny of <i>Hypoxylon fragiforme</i> . <i>Mycological Research</i> , 2006, 110, 811-820.	2.5	54
122	Chemical constituents of the Vietnamese inedible mushroom <i>Xylaria intracolorata</i> . <i>Natural Product Research</i> , 2006, 20, 317-321.	1.8	21
123	<i>Penicillium sclerotiorum</i> Catalyzes the Conversion of Herbertenediol into Its Dimers: Mastigophorenes A and B. <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 256-257.	1.3	16
124	Pregnane-Type Steroids from the Inedible Mushroom <i>Thelephora terrestris</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 309-312.	1.3	5
125	Chemical Constituents of Malagasy Liverworts, Part III: Sesquiterpenoids from <i>Bazzania decrescens</i> and <i>Bazzania madagassa</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 515-518.	1.3	19
126	Cohaerins A and B, azaphilones from the fungus <i>Hypoxylon cohaerens</i> , and comparison of HPLC-based metabolite profiles in <i>Hypoxylon</i> sect. <i>Annulata</i> . <i>Phytochemistry</i> , 2005, 66, 797-809.	2.9	67

#	ARTICLE	IF	CITATIONS
127	Composition of the essential oil of the liverwort <i>Radula perrottetii</i> of Japanese origin. <i>Phytochemistry</i> , 2005, 66, 941-949.	2.9	32
128	Terrestrins A-G: p-Terphenyl derivatives from the inedible mushroom <i>Thelephora terrestris</i> . <i>Phytochemistry</i> , 2005, 66, 1052-1059.	2.9	32
129	ent-Verticillane-type diterpenoids from the Japanese liverwort <i>Jackiella javanica</i> . <i>Phytochemistry</i> , 2005, 66, 1662-1670.	2.9	35
130	Lanostane triterpenoids from the inedible mushroom <i>Fomitopsis spraguei</i> . <i>Phytochemistry</i> , 2005, 66, 1656-1661.	2.9	19
131	Sassafrins A-D, new antimicrobial azaphilones from the fungus <i>Creosphaeria sassafras</i> . <i>Tetrahedron</i> , 2005, 61, 1743-1748.	1.9	45
132	Novel cytotoxic kaurane-type diterpenoids from the New Zealand Liverwort <i>Jungermannia</i> species. <i>Tetrahedron</i> , 2005, 61, 4531-4544.	1.9	50
133	Dimeric azaphilones from the xylariaceous ascomycete <i>Hypoxylon rutilum</i> . <i>Tetrahedron</i> , 2005, 61, 8451-8455.	1.9	22
134	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 412.	1.5	0
135	p-Terphenyl Compounds Possessing Antioxidative Activity from Japanese Inedible Mushrooms. <i>International Journal of Medicinal Mushrooms</i> , 2005, 7, 410-411.	1.5	0
136	Activation of p38 Mitogen-Activated Protein Kinase during gent-11 $\pm$ -Hydroxy-16-kauren-15-one-Induced Apoptosis in Human Leukemia HL-60 Cells. <i>Planta Medica</i> , 2005, 71, 275-277.	1.3	13
137	Induction of Apoptosis by Newent-Kaurene-Type Diterpenoids Isolated from the New Zealand Liverwort Jungermannia Species. <i>Planta Medica</i> , 2005, 71, 1005-1009.	1.3	21
138	Antimicrobial Azaphilones from the Fungus <i>Hypoxylon multiforme</i> . <i>Planta Medica</i> , 2005, 71, 1058-1062.	1.3	38
139	Isolation, Synthesis and Biological Activity of Grifolic Acid Derivatives from the Inedible Mushroom <i>Albatrellus dispansus</i> . <i>Heterocycles</i> , 2005, 65, 2431.	0.7	30
140	Bis(bibenzyls) from Liverworts Inhibit Lipopolysaccharide-Induced Inducible NOS in RAW 264.7 Cells: A Study of Structure-Activity Relationships and Molecular Mechanism. <i>Journal of Natural Products</i> , 2005, 68, 1779-1781.	3.0	46
141	Riccardin C: A natural product that functions as a liver X receptor (LXR) $\beta$ agonist and an LXR $\alpha$ antagonist. <i>FEBS Letters</i> , 2005, 579, 5299-5304.	2.8	66
142	Anent-Kaurene Diterpene Enhances Apoptosis Induced by Tumor Necrosis Factor in Human Leukemia Cells. <i>Planta Medica</i> , 2004, 70, 723-727.	1.3	35
143	A Comparison of Apoptosis and Necrosis Induced by ent-Kaurene-Type Diterpenoids in HL-60 Cells. <i>Planta Medica</i> , 2004, 70, 401-406.	1.3	26
144	Constituents of Isotachis aubertii (Isotachidaceae) collected in Madagascar. <i>Biochemical Systematics and Ecology</i> , 2004, 32, 1073-1078.	1.3	15

#	ARTICLE	IF	CITATIONS
145	Cyclic azaphilones daldinins E and F from the ascomycete fungus Hypoxylon fuscum (Xylariaceae). Phytochemistry, 2004, 65, 469-473.	2.9	50
146	Chemosystematics of the Hepaticae. ChemInform, 2004, 35, no.	0.0	1
147	Efficient synthesis of isoplagiochin D, a macrocyclic bis(bibenzyls), by utilizing an intramolecular Suzuki-Miyaura reaction. Tetrahedron Letters, 2004, 45, 6941-6945.	1.4	35
148	Chemosystematics of the Hepaticae. Phytochemistry, 2004, 65, 623-669.	2.9	206
149	Thelephantins I-N p-terphenyl derivatives from the inedible mushroom <i>Hydnellum caeruleum</i> . Phytochemistry, 2004, 65, 1179-1184.	2.9	23
150	New Azaphilones from the Inedible Mushroom <i>Hypoxylon rubiginosum</i> . Journal of Natural Products, 2004, 67, 1152-1155.	3.0	47
151	Tyromycic Acids A-E, New Lanostane Triterpenoids from the Mushroom <i>Tyromyces fissilis</i> . Journal of Natural Products, 2004, 67, 148-151.	3.0	16
152	Kaurene Diterpene Induces Apoptosis in Human Leukemia Cells Partly through a Caspase-8-Dependent Pathway. Journal of Pharmacology and Experimental Therapeutics, 2004, 311, 115-122.	2.5	38
153	Africane-Type Sesquiterpenoids from the Argentine Liverwort <i>Porella swartziana</i> and Their Antibacterial Activity. Journal of Natural Products, 2004, 67, 31-36.	3.0	35
154	Chemical Constituents of Malagasy Liverworts, Part II: Mastigophoric Acid Methyl Ester of Biogenetic Interest from <i>Mastigophora diclados</i> (Lepicoleaceae Subf. Mastigophoroideae). Chemical and Pharmaceutical Bulletin, 2004, 52, 1382-1384.	1.3	37
155	New Humulane-Type Sesquiterpenes from the Liverworts <i>Tylimanthus tenellus</i> and <i>Marchantia emarginata</i> subsp. <i>tosana</i> . Chemical and Pharmaceutical Bulletin, 2004, 52, 481-484.	1.3	29
156	Terpenoids and Aromatic Compounds from the New Zealand Liverworts <i>Plagiochila</i> , <i>Schistochila</i> , and <i>Heteroscyphus</i> Species. Chemical and Pharmaceutical Bulletin, 2004, 52, 556-560.	1.3	28
157	New Sesquiterpenoids from the New Zealand Liverwort <i>Chiloscyphus subporosus</i> . Chemical and Pharmaceutical Bulletin, 2004, 52, 949-952.	1.3	24
158	ent-Isopimarane-type diterpenoids from the New Zealand liverwort <i>Trichocolea mollissima</i> . Phytochemistry, 2003, 64, 1319-1325.	2.9	24
159	First synthesis of 1,9-dideoxyforskolin from ptychanin A. Tetrahedron Letters, 2003, 44, 2305-2306.	1.4	15
160	Thelephantins A, B and C: three benzoyl p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . Phytochemistry, 2003, 62, 109-113.	2.9	36
161	Volatile components of selected species of the liverwort genera <i>Frullania</i> and <i>Schusterella</i> (Frullaniaceae) from New Zealand, Australia and South America: a chemosystematic approach. Phytochemistry, 2003, 62, 439-452.	2.9	38
162	Sesquiterpene constituents from the liverwort <i>Bazzania japonica</i> . Phytochemistry, 2003, 63, 581-587.	2.9	18

#	ARTICLE	IF	CITATIONS
163	Thelephantins H: five p-terphenyl derivatives from the inedible mushroom <i>Thelephora aurantiotincta</i> . <i>Phytochemistry</i> , 2003, 63, 919-924.	2.9	36
164	Curtisiains E-H: four p-terphenyl derivatives from the inedible mushroom <i>Paxillus curtisii</i> . <i>Phytochemistry</i> , 2003, 64, 649-654.	2.9	23
165	Global phytochemistry: research in Japan. <i>Phytochemistry</i> , 2003, 64, 909-912.	2.9	1
166	Occurrence of a High Concentration of Spider Pheromones in the Ascomycete Fungus <i>Hypoxylon truncatum</i> . <i>Journal of Natural Products</i> , 2003, 66, 1613-1614.	3.0	14
167	Apoptosis-Inducing Properties of ent-Kaurene-Type Diterpenoids from the Liverwort <i>Jungermannia truncata</i> . <i>Planta Medica</i> , 2003, 69, 377-379.	1.3	38
168	Antioxidant Activity of Curtisiains I - L from the Inedible Mushroom <i>Paxillus curtisii</i> . <i>Planta Medica</i> , 2003, 69, 1063-1066.	1.3	48
169	Curtisiains M-Q: Five Novel p-Terphenyl Derivatives from the Mushroom <i>Paxillus curtisii</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1064-1067.	1.3	26
170	Tyromycic Acids F and G: Two New Triterpenoids from the Mushroom <i>Tyromyces fissionis</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1441-1443.	1.3	15
171	New ent-Kaurene-Type Diterpenoids Possessing Cytotoxicity from the New Zealand Liverwort <i>Jungermannia Species</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1189-1192.	1.3	41
172	A Novel Dimeric Lactone Bis-osmundalactone from the Japanese Inedible Mushroom <i>Paxillus atromentosus</i> var. <i>bambusinus</i> . <i>Heterocycles</i> , 2002, 56, 581.	0.7	9
173	Cytotoxic and Apoptosis-Inducing < i>ent</i>-Kaurane-Type Diterpenoids from the Japanese Liverwort < i>Jungermannia truncata</i> N< small>EES</small>. <i>Chemical and Pharmaceutical Bulletin</i> , 2002, 50, 808-813.	1.3	47
174	New Bibenzyl Cannabinoid from the New Zealand Liverwort <i>Radula marginata</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2002, 50, 1390-1392.	1.3	90
175	Chemical Constituents of the Ascomycete <i>Daldinia concentrica</i> . <i>Journal of Natural Products</i> , 2002, 65, 1869-1874.	3.0	88
176	Eremophilanolides and other constituents from the Argentine liverwort <i>Frullania brasiliensis</i> . <i>Phytochemistry</i> , 2002, 59, 205-213.	2.9	35
177	Neogrifolin derivatives possessing anti-oxidative activity from the mushroom <i>Albatrellus ovinus</i> . <i>Phytochemistry</i> , 2002, 59, 731-737.	2.9	65
178	Iridoid glucosides from roots of Vietnamese <i>Paederia scandens</i> . <i>Phytochemistry</i> , 2002, 60, 505-514.	2.9	45
179	Concentriols B, C and D, three squalene-type triterpenoids from the ascomycete <i>Daldinia concentrica</i> . <i>Phytochemistry</i> , 2002, 61, 345-353.	2.9	32
180	Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> . <i>Journal of Natural Products</i> , 2001, 64, 1309-1317.	3.0	33

#	ARTICLE	IF	CITATIONS
181	New Chroman Derivatives from the Japanese Liverworts <i>Metacalypogeia cordifolia</i> and <i>Cephalozia otaruensis</i> .. Chemical and Pharmaceutical Bulletin, 2001, 49, 924-926.	1.3	14
182	Occurrence of Polygodial and 1-(2,4,6-Trimethoxyphenyl)-but-2-en-1-one from Some Ferns and Liverworts: Role of Pungent Components in Bryophytes and Pteridophytes Evolution.. Chemical and Pharmaceutical Bulletin, 2001, 49, 1380-1381.	1.3	28
183	Sesquiterpene hydrocarbons with trifarane backbone in the liverwort <i>Trocholejeunea sandvicensis</i> . Phytochemistry, 2001, 57, 499-506.	2.9	10
184	Sesqui- and diterpenoids from two Japanese and three European liverworts. Phytochemistry, 2001, 56, 347-352.	2.9	36
185	secocuparane-type sesquiterpenoid from the Japanese liverwort <i>Jungermannia infusca</i> . Phytochemistry, 2001, 56, 807-810.	2.9	9
186	Recent advances in phytochemistry of bryophytes-acetogenins, terpenoids and bis(bibenzyl)s from selected Japanese, Taiwanese, New Zealand, Argentinean and European liverworts. Phytochemistry, 2001, 56, 297-312.	2.9	159
187	First syntheses of 1,13- and 1,15-dihydroxyherbertenes, and herbertenolide by applying intramolecular Heck reaction for the construction of adjacent quaternary centers. Tetrahedron, 2001, 57, 9299-9307.	1.9	29
188	Molecular chemotaxonomy of <i>Daldinia</i> and other Xylariaceae. Mycological Research, 2001, 105, 1191-1205.	2.5	47
189	Sesquiterpene Lactones and Acetogenin Lactones from the Hepaticae and Chemosystematics of the Liverworts <i>Frullania</i> , <i>Plagiochila</i> and <i>Porella</i> . Heterocycles, 2001, 54, 1057.	0.7	23
190	Clerodane-type Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. Chemical and Pharmaceutical Bulletin, 2000, 48, 1818-1821.	1.3	10
191	Pinguisane and dimeric pinguisane-type sesquiterpenoids from the Japanese liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . Phytochemistry, 2000, 53, 593-604.	2.9	50
192	Highly oxygenated sesquiterpenes from the liverwort <i>Trocholejeunea sandvicensis</i> . Phytochemistry, 2000, 53, 271-276.	2.9	19
193	New Chlorinated Cyclic Bis(bibenzyls) from the Liverworts <i>Herbertus sakuraii</i> and <i>Mastigophora diclados</i> . Tetrahedron, 2000, 56, 3153-3159.	1.9	57
194	Syntheses of isocitric acid derivatives and biological evaluation. Tetrahedron Letters, 2000, 41, 3095-3098.	1.4	5
195	Herbertane-type sesquiterpenoids from the liverwort <i>Herbertus sakuraii</i> . Phytochemistry, 2000, 55, 247-253.	2.9	46
196	Isolation and Structure of Striatic Acid from Liverwort <i>Cheilolejeunea serpentina</i> and the Absolute Configuration by Synthesis. Tetrahedron, 2000, 56, 1655-1659.	1.9	11
197	Synthesis of cryptoporic acid A methyl ester. Tetrahedron Letters, 2000, 41, 3099-3102.	1.4	13
198	Chemical structures of macrocyclic bis(bibenzyls) isolated from liverworts (Hepaticae). Spectroscopy, 2000, 14, 149-175.	0.8	86

#	ARTICLE	IF	CITATIONS
199	Phytochemistry of Bryophytes., 1999, , 319-342.		25
200	Biosynthesis of cyclic bis(bibenzyls) in <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1999, 50, 589-598.	2.9	64
201	A 7-nordumortenone and other dumortane derivatives from the Argentine liverwort <i>Dumontiera hirsuta</i> . <i>Phytochemistry</i> , 1999, 51, 281-287.	2.9	24
202	Sesqui- and diterpenoids from <i>Ptilidium ciliare</i> and <i>Barbilophozia</i> species (liverworts). <i>Phytochemistry</i> , 1999, 51, 563-566.	2.9	16
203	Ptychantols A-C, macrocyclic bis(bibenzyls), possessing a trans-stilbene structure from the liverwort <i>Ptychanthus striatus</i> . <i>Phytochemistry</i> , 1999, 52, 501-509.	2.9	28
204	A revision of the positive sign of the optical rotation and its maximum value of $\pm$ -eudesmol. <i>Phytochemistry</i> , 1999, 52, 689-694.	2.9	16
205	Cytochromes P-450 catalyze the formation of marchantins A and C in <i>Marchantia polymorpha</i> . <i>Phytochemistry</i> , 1999, 52, 1195-1202.	2.9	48
206	Sesquiterpenoids, hopanoids and bis(bibenzyls) from the Argentine liverwort <i>Plagiochasma rupestre</i> . <i>Phytochemistry</i> , 1999, 52, 1323-1329.	2.9	30
207	New acorane- and cuparane-type sesqui- and new labdane- and seco-labdanne-type diterpenoids from the Japanese liverwort <i>Jungermannia infusca</i> (Mitt.) Steph.. <i>Tetrahedron</i> , 1999, 55, 9117-9132.	1.9	22
208	A Novel Skeletal Diterpenoid from the German Liverwort <i>Barbilophozia hatcheri</i> (Evans) Loeske.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 138-139.	1.3	14
209	Occurrence of a Bis-benzyl Derivative in the Japanese Fern <i>Hymenophyllum barbatum</i> : First Isolation and Identification of Perrottetin H from the Pteridophytes.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 297-298.	1.3	25
210	Novel ent-vibsane- and dolabellane-type diterpenoids from the liverwort <i>Odontoschisma denudatum</i> . <i>Tetrahedron Letters</i> , 1998, 39, 579-582.	1.4	25
211	Two novel Diels-Alder reaction-type dimeric pinguisane sesquiterpenoids and related compounds from the liverwort <i>Porella acutifolia</i> subsp. <i>tosana</i> . <i>Tetrahedron Letters</i> , 1998, 39, 2977-2980.	1.4	23
212	Two novel skeletal diterpenoids, neodenudatenones A and B, from the liverwort <i>Odontoschisma denudatum</i> . <i>Tetrahedron Letters</i> , 1998, 39, 3791-3794.	1.4	13
213	Triterpenoid constituents of the moss <i>Floribundaria aurea</i> subsp. <i>Nipponica</i> . <i>Phytochemistry</i> , 1998, 48, 297-299.	2.9	23
214	Diterpenoids from the Japanese liverwort jungermannia INFUSCA fn1 fn1This paper is dedicated, with best wishes, to Professor G. H. Neil Towers on the occasion of his 75th birthday. in honour of professor G. H. Neil Towers 75TH birthday. <i>Phytochemistry</i> , 1998, 49, 601-608.	2.9	19
215	Cytotoxic 2,3-Secoiridoid-Type Sesquiterpenoids from the Liverwort <i>Plagiochila ovalifolia</i> . <i>Planta Medica</i> , 1998, 64, 462-464.	1.3	35
216	Biologically Active Substances of Japanese Inedible Mushrooms. <i>Heterocycles</i> , 1998, 47, 1067.	0.7	62

#	ARTICLE	IF	CITATIONS
217	Novel skeletal diterpenoids from the Japanese liverwort <i>Pallavicinia subciliata</i> .. Chemical and Pharmaceutical Bulletin, 1998, 46, 178-180.	1.3	38
218	New Sesqui- and Diterpenoids from the Japanese Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. Chemical and Pharmaceutical Bulletin, 1998, 46, 1184-1185.	1.3	16
219	Occurrence of ent-Sesquiterpene in the Japanese Moss- <i>Plagiomnium acutum</i> : First Isolation and Identification of the ent-Sesqui- and Dolabellane-type Diterpenoids from the Musci.. Chemical and Pharmaceutical Bulletin, 1998, 46, 1488-1489.	1.3	19
220	Eudesmane-Type Sesquiterpenoids from Japanese Liverwort <i>Frullania tamarisci</i> subsp. <i>obscura</i> .. Chemical and Pharmaceutical Bulletin, 1998, 46, 542-544.	1.3	17
221	Chemical Conversion of Labdane-Type Diterpenoid Isolated from the Liverwort <i>Porella perrottetiana</i> into (-)-Ambrox. Heterocycles, 1998, 49, 315.	0.7	20
222	Dumortenols, Novel Skeletal Sesquiterpenoids from the Argentinian Liverwort <i>Dumontiera hirsuta</i> .. Chemical and Pharmaceutical Bulletin, 1997, 45, 2119-2121.	1.3	24
223	Isolation, structure, and synthesis of chenopodanol and the absolute configuration of chenopodene and chenopodanol. Canadian Journal of Chemistry, 1997, 75, 634-640.	1.1	15
224	Isolation, Structure Elucidation, and Chemical Derivatization of a New Cyclic Bisbibenzyl Dimer, Pusilatin E, from the Liverwort <i>Riccardia multifida</i> subsp. <i>decrescens</i> . Journal of Natural Products, 1997, 60, 145-147.	3.0	19
225	Scapaundulins A and B, two novel dimeric labdane diterpenoids, and related compounds from the Japanese liverwort <i>Scapania undulata</i> (L.) Dum.. Tetrahedron Letters, 1997, 38, 1975-1978.	1.4	28
226	Eudesmane-type sesquiterpene lactones from the Japanese liverwort <i>Frullania densiloba</i> . Phytochemistry, 1997, 45, 555-558.	2.9	24
227	Sesquiterpene and other constituents of the liverwort <i>Dumontiera hirsuta</i> . Phytochemistry, 1997, 44, 293-298.	2.9	37
228	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia rotundata</i> . Phytochemistry, 1997, 44, 653-657.	2.9	11
229	Volatile components of the liverworts <i>Archilejeunea olivacea</i> , <i>cheilolejeunea imbricata</i> and <i>Leptolejeunea elliptica</i> . Phytochemistry, 1997, 44, 1261-1264.	2.9	28
230	A comparative study on three chemo-types of the liverwort <i>Conocephalum conicum</i> using volatile constituents. Phytochemistry, 1997, 44, 1265-1270.	2.9	48
231	Sesqui- and diterpenoids from the japanese liverwort <i>Jungermannia hattoriana</i> . Phytochemistry, 1997, 45, 353-363.	2.9	30
232	Sesquiterpenoids from the three Japanese liverworts <i>Lejeunea aquatica</i> , <i>L. flava</i> and <i>L. japonica</i> . Phytochemistry, 1997, 46, 145-150.	2.9	30
233	Butenolides from <i>Marchantia paleacea</i> subspecies <i>diptera</i> . Phytochemistry, 1997, 46, 293-296.	2.9	30
234	Terpenoids from the Japanese liverworts <i>Jackiella javanica</i> and <i>Jungermannia infusca</i> . Phytochemistry, 1997, 46, 1203-1208.	2.9	36

#	ARTICLE	IF	CITATIONS
235	Heterocyclic Compounds Found in Bryophytes. <i>Heterocycles</i> , 1997, 46, 795.	0.7	24
236	Total Synthesis and Absolute Configuration of Riccardiphenols A and B, Isolated from the Liverwort <i>Riccardia crassa</i> . <i>Journal of Organic Chemistry</i> , 1996, 61, 5362-5370.	3.2	21
237	Isoprenyl Phenyl Ethers from Liverworts of the Genus <i>Trichocolea</i> : Cytotoxic Activity, Structural Corrections, and Synthesis. <i>Journal of Natural Products</i> , 1996, 59, 729-733.	3.0	45
238	A New Monocyclic Diterpene from the Liverwort <i>Jungermannia infusca</i> (MITT.) STEPH.. <i>Chemical and Pharmaceutical Bulletin</i> , 1996, 44, 1628-1630.	1.3	11
239	Isoplagiochins C and D, New Type of Macroyclic Bis(bibenzyls), Having Two Biphenyl Linkages from the Liverwort <i>Plagiochila fruticosa</i> . <i>Chemistry Letters</i> , 1996, 25, 741-742.	1.3	45
240	Sesquiterpenoids from a Cell Suspension Culture of the Liverwort <i>Porella vernicosa</i> Lindb. , 1996, 11, 53-56.		14
241	Novel sesquiterpenoids from the Colombian liverwort <i>Porella swartziana</i> . <i>Tetrahedron</i> , 1996, 52, 6339-6354.	1.9	18
242	Phenolic constituents of the liverwort: Four novel cyclic bisbibenzyl dimers from <i>Blasia pusilla</i> L. <i>Tetrahedron</i> , 1996, 52, 14487-14500.	1.9	69
243	(+)-Cavicularin: A novel optically active cyclic bibenzyl-dihydrophenanthrene derivative from the liverwort <i>Cavicularia densa</i> Steph. <i>Tetrahedron Letters</i> , 1996, 37, 4745-4748.	1.4	54
244	Terpenoids and aromatic compounds from six liverworts. <i>Phytochemistry</i> , 1996, 41, 207-211.	2.9	50
245	Terpenoid constituents of the liverwort <i>Heteroscyphus coalitus</i> . <i>Phytochemistry</i> , 1996, 41, 575-580.	2.9	27
246	Cytochalasins from a <i>Daldinia</i> sp. of fungus. <i>Phytochemistry</i> , 1996, 41, 821-828.	2.9	45
247	Eudesmane-type sesquiterpenoids from the liverwort <i>Lepidozia vitrea</i> . <i>Phytochemistry</i> , 1996, 41, 833-836.	2.9	18
248	Ent-kaurane-type diterpenoids from the liverwort <i>Jungermannia exsertifolia</i> ssp. <i>cordifolia</i> . <i>Phytochemistry</i> , 1996, 41, 1129-1141.	2.9	19
249	Phytol esters and phaeophytins from the hornwort <i>Megaceros flagellaris</i> . <i>Phytochemistry</i> , 1996, 41, 1373-1376.	2.9	20
250	Sesqui- and di-terpenoids from the liverwort <i>Jungermannia vulcanicola</i> . <i>Phytochemistry</i> , 1996, 42, 93-96.	2.9	16
251	A 10-phenyl-[11]-cytochalasan from a species of <i>Daldinia</i> . <i>Phytochemistry</i> , 1996, 42, 173-176.	2.9	28
252	Sesquiterpenoids from the liverworts <i>Bazzania trilobata</i> and <i>Porella canariensis</i> . <i>Phytochemistry</i> , 1996, 42, 1361-1366.	2.9	47

#	ARTICLE	IF	CITATIONS
253	Nudenoic acid: A novel tricyclic sesquiterpenoid from the Taiwanese liverwort <i>Mylia nuda</i> . <i>Tetrahedron Letters</i> , 1996, 37, 9307-9308.	1.4	12
254	Terpenoid constituents of the new zealand liverwort <i>Jamesoniella tasmanica</i> . <i>Phytochemistry</i> , 1996, 43, 1057-1064.	2.9	18
255	Africane- and monocyclofarnesane-type sesquiterpenoids from the liverwort <i>Porella subobtusa</i> . <i>Phytochemistry</i> , 1996, 43, 1285-1291.	2.9	13
256	A phenethyl glycoside from <i>Conocephalum conicum</i> . <i>Phytochemistry</i> , 1996, 43, 1087-1088.	2.9	14
257	The absolute structures of new 1.BETA.-hydroxysacculatane-type diterpenoids with piscidal activity from the liverwort <i>Pellia endiviifolia</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1995, 43, 2030-2032.	1.3	27
258	Sesquiterpenoids from the liverwort <i>dicranolejeunea yoshinaga</i> (HATT.) MIZUT.. <i>Chemical and Pharmaceutical Bulletin</i> , 1995, 43, 714-716.	1.3	15
259	Structures of Four Novel Highly Oxygenated Labdane-Type Diterpenoids, Ptychantins F-I, from the Liverwort <i>Ptychanthus striatus</i> . <i>Chemistry Letters</i> , 1995, 24, 481-482.	1.3	9
260	Epi-neoverrucosane- and ent-clerodane-type diterpenoids and ent-2,3-secoaromadendrane- and calamenene-type sesquiterpenoids from the liverwort <i>heteroscyphus planus</i> . <i>Phytochemistry</i> , 1995, 38, 119-127.	2.9	33
261	Sesquiterpenes and a phenolic compound from the liverwort <i>Omphalanthus filiformis</i> . <i>Phytochemistry</i> , 1995, 38, 651-653.	2.9	16
262	Terpenoids from the liverworts <i>Symphyogyna brasiliensis</i> and unidentified <i>Frullania</i> species. <i>Phytochemistry</i> , 1995, 39, 99-103.	2.9	18
263	Five 10-phenyl-[11]-cytochalasans from a <i>Daldinia</i> fungal species. <i>Phytochemistry</i> , 1995, 40, 135-140.	2.9	36
264	Trifarienols A-E, trifarane-type sesquiterpenoids from the Malaysian liverwort <i>Cheilolejeunea trifaria</i> . <i>Phytochemistry</i> , 1995, 40, 171-176.	2.9	14
265	Clerodane- and halimane-type diterpenoids from the liverwort <i>Jungermannia hyalina</i> . <i>Phytochemistry</i> , 1995, 40, 209-212.	2.9	19
266	(+)-Osmundalactone, $\beta$ -lactones and spiromentins from the fungus <i>Paxillus atrotomentosus</i> . <i>Phytochemistry</i> , 1995, 40, 1251-1257.	2.9	53
267	Structure of conocephalenol, a sesquiterpenoid alcohol from the European liverwort <i>Conocephalum conicum</i> : determination of the absolute configuration by total synthesis. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1995, , 593.	0.9	15
268	Synthesis and the absolute configuration of the sesquiterpene aldehyde tridensenol from the Taiwanese liverwort <i>Bazzania tridens</i> . <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1995, , 1513.	0.9	12
269	Structures of five new highly oxygenated labdane-type diterpenoids, ptychantins A-E, closely related to forskolin from the liverwort <i>Ptychanthus striatus</i> . <i>Tetrahedron Letters</i> , 1994, 35, 5457-5460.	1.4	21
270	Structures of four novel macrocyclic bis(bibenzyl) dimers, pusilatins A-D from the liverwort <i>Blasia pusilla</i> . <i>Tetrahedron Letters</i> , 1994, 35, 909-910.	1.4	30

#	ARTICLE	IF	CITATIONS
271	Two novel macrocyclic bis(Bibenzyls), Isoplagiochins A and B from the liverwort <i>Plagiochila Fruticosa</i> . <i>Tetrahedron Letters</i> , 1994, 35, 911-912.	1.4	24
272	Trifarienols A and B, Isolated from the liverwort <i>Cheilolejeunea trifaria</i> . Sesquiterpenes having a new carbon skeleton, trifarane. <i>Tetrahedron Letters</i> , 1994, 35, 4787-4788.	1.4	16
273	Sesquiterpene esters from the liverwort <i>Plagiochila poreloides</i> . <i>Phytochemistry</i> , 1994, 37, 1091-1093.	2.9	12
274	Sesqui- and diterpenoids from <i>Plagiochila</i> species. <i>Phytochemistry</i> , 1994, 36, 1425-1430.	2.9	27
275	Sesquiterpenoids from the liverwort <i>Marsupella aquatica</i> . <i>Phytochemistry</i> , 1994, 37, 777-779.	2.9	10
276	Rearranged ent-eudesmane- and ent-eremophilane-type sesquiterpenoids from the liverwort <i>Frullania dilatata</i> . <i>Phytochemistry</i> , 1994, 37, 1317-1321.	2.9	15
277	Cadinane-type sesquiterpenoids from the liverwort <i>Scapania undulata</i> . <i>Phytochemistry</i> , 1994, 37, 1323-1325.	2.9	31
278	Chenopodene, marchantin P and riccardin G from the liverwort <i>Marchantia chenopoda</i> . <i>Phytochemistry</i> , 1994, 36, 73-76.	2.9	31
279	Volatile constituent of the liverwort <i>chiloscyphus pallidus</i> (mitt.) engel & schuster. <i>Flavour and Fragrance Journal</i> , 1994, 9, 237-240.	2.6	17
280	Naphthalene and isocoumarin derivatives from the liverwort <i>Wettsteinia schusterana</i> . <i>Phytochemistry</i> , 1994, 37, 233-235.	2.9	11
281	Sesqui- and diterpenoids from the panamanian liverwort <i>Bryopteris filicina</i> . <i>Phytochemistry</i> , 1994, 37, 433-439.	2.9	22
282	Ent-longipinane-type sesquiterpenoid from the liverwort <i>Marsupella emarginata</i> . <i>Phytochemistry</i> , 1994, 37, 1767-1768.	2.9	10
283	Structures of a Novel Binaphthyl and Three Novel Benzophenone Derivatives with Plant-Growth Inhibitory Activity from the Fungus <i>Daldinia concentrica</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1528-1530.	1.3	53
284	Structures of Daldinins A-C, Three Novel Azaphilone Derivatives from Ascomycetous Fungus <i>Daldinia concentrica</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2397-2399.	1.3	46
285	New Highly Oxygenated 6,7-Seco-kaurane- and Bis-kaurane-Type Diterpenoids from The Liverwort <i>Jungermannia exsertifolia</i> STEPH. ssp. <i>cordifolia</i> (DUM.) VANA.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 2656-2658.	1.3	9
286	Marchantin A Trimethyl Ether: Its Molecular Structure and Tubocurarine-like Skeletal Muscle Relaxation Activity.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 52-56.	1.3	38
287	Stereostructure of Plagiochiline A and Conversion of Plagiochiline A and Stearylvelutinal into Hot-Tasting Compounds by Human Saliva.. <i>Chemical and Pharmaceutical Bulletin</i> , 1994, 42, 1542-1544.	1.3	18
288	Highlights in phytochemistry of hepaticae-biologically active terpenoids and aromatic compounds. <i>Pure and Applied Chemistry</i> , 1994, 66, 2193-2196.	1.9	34

#	ARTICLE	IF	CITATIONS
289	Fusicorragatol from the Venezuelan liverwort <i>Plagiochila corrugata</i> .. Chemical and Pharmaceutical Bulletin, 1994, 42, 2650-2652.	1.3	13
290	Drimane-type sesquiterpenoids from <i>Cryptoporus volvatus</i> infected by <i>Paecilomyces varioti</i> . Phytochemistry, 1993, 33, 1055-1059.	2.9	20
291	Terpenoids from some German and Russian liverworts. Phytochemistry, 1993, 33, 1445-1448.	2.9	28
292	Bisbibenzyl and sesquiterpenoids from the liver wort <i>Jubula japonica</i> . Phytochemistry, 1993, 34, 1135-1137.	2.9	12
293	Terpenoids of the liverwort <i>Frullanoides densifolia</i> and <i>Trocholejeunea sandvicensis</i> . Phytochemistry, 1993, 32, 335-348.	2.9	52
294	Setiformenol, isolated from the liverwort <i>tetralophozia setiformis</i> , the first example of cembrane-type diterpene from bryophytes. Tetrahedron Letters, 1993, 34, 643-644.	1.4	11
295	Terpenoids from six lophoziaeae liverworts. Phytochemistry, 1993, 34, 181-190.	2.9	33
296	Inhibitory Effect of Cryptoporic Acid E, a Product from Fungus <i>&lt; i&gt;Cryptoporus volvatus&lt;/i&gt;</i> , on Colon Carcinogenesis Induced with N-methyl-N-nitrosourea in Rats and with 1,2-dimethylhydrazine in Mice. Japanese Journal of Cancer Research, 1992, 83, 830-834.	1.7	21
297	Sesquiterpene derivatives and a norsesquiterpenoid from the liverworts <i>Riccardia crassa</i> and <i>Porella caespitans</i> var. <i>Setigera</i> . Phytochemistry, 1992, 32, 137-140.	2.9	15
298	Cryptoporic acids A-G, drimane-type sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . Phytochemistry, 1992, 31, 579-592.	2.9	51
299	Studies on the liverwort sesquiterpene alcohol tamariscol. Synthesis and absolute configuration. Journal of the Chemical Society Perkin Transactions 1, 1991, , 435.	0.9	23
300	Novel neurotrophic isocuparane-type sesquiterpene dimers, mastigophorenanes A, B, C and D, isolated from the liverwort <i>Mastigophora diclados</i> . Journal of the Chemical Society Perkin Transactions 1, 1991, , 2737.	0.9	103
301	Neurotrophic secoaromadendrane-type sesquiterpenes from the liverwort <i>Plagiochila fruticosa</i> . Phytochemistry, 1991, 30, 4061-4065.	2.9	23
302	Geographical distribution of tamariscol, a mossy odorous sesquiterpene alcohol, in the liverwort <i>Frullania tamarisci</i> and related species. Phytochemistry, 1991, 30, 2295-2300.	2.9	17
303	Sesquiterpenoids from the liverwort <i>Porella acutifolia</i> subsp. <i>Tosana</i> . Phytochemistry, 1991, 30, 567-573.	2.9	33
304	Terpenoids and aromatic compounds from selected Ecuadorian liverworts*. Phytochemistry, 1991, 30, 215-217.	2.9	35
305	Prenyl bibenzyls from the liverwort <i>Radula kojana</i> . Phytochemistry, 1991, 30, 219-234.	2.9	47
306	Prenyl bibenzyls from the liverworts <i>Radula perrottetii</i> and <i>Radula complanata</i> . Phytochemistry, 1991, 30, 235-251.	2.9	89

#	ARTICLE	IF	CITATIONS
307	Cyclopropanochroman derivatives from the liverwort <i>Radula javanica</i> . <i>Phytochemistry</i> , 1991, 30, 325-328.	2.9	28
308	Bis(bibenzyl) ethers from <i>Pellia endiviifolia</i> . <i>Phytochemistry</i> , 1991, 30, 1523-1530.	2.9	14
309	Differential effects of cryptoporic acids D and E, inhibitors of superoxide anion radical release, on tumor promotion of okadaic acid in mouse skin. <i>Carcinogenesis</i> , 1991, 12, 1129-1131.	2.8	17
310	Ent-sesquiterpenoids and cyclic bis(bibenzyls) from the german liverwort <i>Marchantia Polymorpha</i> . <i>Phytochemistry</i> , 1990, 29, 1577-1584.	2.9	55
311	Bitter kaurane-type diterpene glucosides from the liverwort <i>Jungermannia infusca</i> . <i>Phytochemistry</i> , 1990, 29, 1619-1623.	2.9	18
312	Homomonoo- and sesquiterpenoids from the liverwort <i>Lophocolea heterophylla</i> . <i>Phytochemistry</i> , 1990, 29, 2334-2337.	2.9	39
313	Fusicoccane-, dolabellane- and rearranged labdane-type diterpenoids from the liverwort <i>Pleurozia gigantea</i> . <i>Phytochemistry</i> , 1990, 29, 2597-2603.	2.9	46
314	The superoxide release inhibitors, cryptoporic acids C, D, and E; dimeric drimane sesquiterpenoid ethers of isocitric acid from the fungus <i>Cryptoporus volvatus</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1989, , 258.	2.0	31
315	A Highly efficient preparation of lunularic acid and some biological activities of stilbene and dihydrostilbene derivatives. <i>Phytochemistry</i> , 1988, 27, 109-113.	2.9	34
316	Distribution of cyclic bis(bibenzyls) in the South African liverwort <i>Marchantia polymorpha</i> †. <i>Phytochemistry</i> , 1988, 27, 161-163.	2.9	24
317	Labdane type diterpenoids from the liverwort <i>Frullania hamachiloba</i> . <i>Phytochemistry</i> , 1988, 27, 1789-1793.	2.9	26
318	Fatty acids and cyclic bis(bibenzyls) from the New Zealand liverwort <i>Monoclea forsteri</i> . <i>Phytochemistry</i> , 1988, 27, 2603-2608.	2.9	35
319	Mastigophorenes: novel dimeric isocuparane-type sesquiterpenoids from the liverwort <i>Mastigophora diclados</i> . <i>Journal of the Chemical Society Chemical Communications</i> , 1988, , 1341.	2.0	24
320	Plagiochins A, B, C, and D, new type of macrocyclic bis(bibenzyls) having a biphenyl linkage between the ortho positions to the benzyl methylenes, from the liverwort subsp.. <i>Tetrahedron Letters</i> , 1987, 28, 6295-6298.	1.4	37
321	Cryptoporic acids A and B, novel bitter drimane sesquiterpenoid ethers of isocitric acid, from the fungus. <i>Tetrahedron Letters</i> , 1987, 28, 6303-6304.	1.4	29
322	Cyclic bis(bibenzyls) and related compounds from the liverworts <i>Marchantia polymorpha</i> and <i>Marchantia palmata</i> . <i>Phytochemistry</i> , 1987, 26, 1811-1816.	2.9	91
323	Bibenzyl derivatives from the Australian liverwort <i>Frullania falciloba</i> . <i>Phytochemistry</i> , 1987, 26, 1023-1025.	2.9	36
324	Bibenzyl derivatives from <i>Frullania</i> Species. <i>Phytochemistry</i> , 1987, 26, 1117-1122.	2.9	26

#	ARTICLE	IF	CITATIONS
325	Terpenoids from the french liverwort targonia hypophylla. <i>Phytochemistry</i> , 1986, 25, 2555-2556.	2.9	22
326	Isotachin c and balantiolide two aromatic compounds from the new zealand liverwort balantiopsis rosea. <i>Phytochemistry</i> , 1986, 25, 2543-2546.	2.9	25
327	Occurrence of a potent piscicidal diterpenedial in the liverwort Riccardia lobata var. Yakushimensis. <i>Phytochemistry</i> , 1985, 24, 261-262.	2.9	25
328	Total assignment of <sup>1</sup> H and <sup>13</sup> C NMR spectra of marchantins isolated from liverworts and its application to structure determination of two new macrocyclic bis(bibenzyls) from and. <i>Tetrahedron Letters</i> , 1985, 26, 4735-4738.	1.4	51
329	Perrottetins E, F, and G from (liverwort)-isolation, structure determination, and synthesis of perrottetin e. <i>Tetrahedron Letters</i> , 1985, 26, 6097-6100.	1.4	58
330	Isotachin A and isotachin B, two sulphur-containing acrylates from the liverwort isotachis japonica. <i>Phytochemistry</i> , 1985, 24, 1505-1508.	2.9	40
331	Distribution of novel cyclic bisbibenzyls in Marchantia and Riccardia species. <i>Phytochemistry</i> , 1983, 22, 1413-1415.	2.9	53
332	Sesquiterpenoids from Chiloscyphus, Clasmatocolea and Frullania species. <i>Phytochemistry</i> , 1983, 22, 961-964.	2.9	33
333	Riccardin A and riccardin B, two novel cyclic bis(bibenzyls) possessing cytotoxicity from the liverwort Riccardia multifida (L.) S. Gray. <i>Journal of Organic Chemistry</i> , 1983, 48, 2164-2167.	3.2	64
334	Riccardin C, a novel cyclic bibenzyl derivative from Reboulia hemisphaerica. <i>Phytochemistry</i> , 1982, 21, 2143-2144.	2.9	74
335	Terpenoids and bibenzyls from some New Zealand liverworts. <i>Phytochemistry</i> , 1982, 21, 2663-2667.	2.9	22
336	Cuparane- and isocuparane-type sesquiterpenoids in liverworts of the genus Herbertus. <i>Phytochemistry</i> , 1982, 21, 2471-2473.	2.9	24
337	Novel bibenzyl derivatives and ent-cuparene-type sesquiterpenoids from Radula species. <i>Phytochemistry</i> , 1982, 21, 2481-2490.	2.9	46
338	Terpenoids and bibenzyls of 25 liverwort Frullania species. <i>Phytochemistry</i> , 1981, 20, 2187-2194.	2.9	42
339	Bibenzyls from Radula tokiensis and R. Japonica. <i>Phytochemistry</i> , 1981, 20, 858-859.	2.9	24
340	Sesquiterpenes from Japanese liverworts. <i>Phytochemistry</i> , 1981, 20, 2359-2366.	2.9	49
341	A potent cytotoxic warburganal and related drimane-type sesquiterpenoids from Polygonum hydropiper. <i>Phytochemistry</i> , 1980, 21, 2895-2898.	2.9	65
342	Seven new bibenzyls and a dihydrochalcone from Radula variabilis. <i>Phytochemistry</i> , 1978, 17, 2005-2010.	2.9	51

#	ARTICLE	IF	CITATIONS
343	New bibenzyls from <i>Radula complanata</i> . <i>Phytochemistry</i> , 1978, 17, 2115-2117.	2.9	34
344	Sesquiterpenes from <i>Porella</i> species. <i>Phytochemistry</i> , 1978, 17, 457-460.	2.9	53
345	Sesquiterpenes of six <i>Porella</i> species (hepaticae). <i>Phytochemistry</i> , 1976, 15, 1929-1931.	2.9	40