

Jung Yoon Pyo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3289299/publications.pdf>

Version: 2024-02-01

174
papers

1,751
citations

331670

21
h-index

434195

31
g-index

176
all docs

176
docs citations

176
times ranked

2654
citing authors

#	ARTICLE	IF	CITATIONS
1	Birmingham Vasculitis Activity Score und der Short Form 36-Item Health Survey als Prädiktoren aktueller depressiver Störungen bei Patienten mit antineutrophile-zytoplasmatische-Antikörper-assoziiierter Vaskulitis während der SARS-CoV-2-Pandemie. Zeitschrift Fur Rheumatologie, 2024, 83, 222-229.	1.0	1
2	Reclassification of previously diagnosed GPA patients using the 2022 ACR/EULAR classification criteria. Rheumatology, 2023, 62, 1179-1186.	1.9	8
3	2019 American College of Rheumatology/European League Against Rheumatism classification criteria for IgG4-related disease by Wallace et al. Annals of the Rheumatic Diseases, 2022, 81, e179-e179.	0.9	10
4	Increased prevalence rate of metabolic syndrome is an independent predictor of cardiovascular disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Rheumatology International, 2022, 42, 291-302.	3.0	7
5	Rheumatoid factor positivity in antineutrophil cytoplasmic antibody-associated vasculitis: a distinct clinical entity or innocent bystander?. Rheumatology, 2022, 61, 1366-1375.	1.9	3
6	Nutrition Risk Index Score at Diagnosis Can Effectively Predict Poor Prognosis in Patients With Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. , 2022, 32, 423-431.		4
7	Drug Survival of Biologic Therapy in Elderly Patients With Rheumatoid Arthritis Compared With Nonelderly Patients. Journal of Clinical Rheumatology, 2022, 28, e81-e88.	0.9	4
8	A retrospective analysis of antineutrophil cytoplasmic antibody-associated vasculitis aiming for an equation prediction end-stage renal disease. Clinical Rheumatology, 2022, 41, 773-781.	2.2	5
9	Clinical application of low erythrocyte sedimentation rate/high C-reactive protein to antineutrophil cytoplasmic antibody-associated vasculitis. Journal of Clinical Laboratory Analysis, 2022, 36, e24237.	2.1	2
10	Aortic valve surgery in patients with Takayasu's arteritis: a nationwide analysis of 1,197 patients during a 9-year period. Clinical and Experimental Rheumatology, 2022, 40, 568-574.	0.8	0
11	New body mass index for predicting prognosis in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Journal of Clinical Laboratory Analysis, 2022, , e24357.	2.1	4
12	Clinical Features of Anti-Synthetase Syndrome Associated with Prognosis in Patients with Dermatomyositis and Polymyositis. Journal of Clinical Medicine, 2022, 11, 2052.	2.4	2
13	Incidence, prevalence and risk of stroke in patients with Takayasu arteritis: a nationwide population-based study in South Korea. Stroke and Vascular Neurology, 2022, 7, 149-157.	3.3	7
14	Anti-Citrullinated Peptide Antibody Expression and Its Association with Clinical Features and Outcomes in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. Medicina (Lithuania), 2022, 58, 558.	2.0	2
15	Modified Body Mass Index at Diagnosis is a Useful Predictor of Mortality in Patients With Antineutrophil Cytoplasmic Antibody-associated Vasculitis. Journal of Rheumatic Diseases, 2022, 29, 154-161.	1.1	0
16	Comparison of the analytical and clinical performances of four anti-cyclic citrullinated peptide antibody assays for diagnosing rheumatoid arthritis. Clinical Rheumatology, 2021, 40, 565-573.	2.2	5
17	Clinical and imaging findings suggestive of histopathological immunoglobulin G4-related disease: a single-center retrospective study. Clinical Rheumatology, 2021, 40, 1423-1430.	2.2	2
18	Serological Biomarkers and Indices for the Current Activity and Prognosis of ANCA-Associated Vasculitis: Experience in a Single Centre in Korea. Yonsei Medical Journal, 2021, 62, 279.	2.2	8

#	ARTICLE	IF	CITATIONS
19	Systemic inflammation response index predicts all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Urology and Nephrology</i> , 2021, 53, 1631-1638.	1.4	4
20	Association Between Serum Alarmin Levels and Disease-specific Indices in Patients With Anti-neutrophil Cytoplasmic Antibody-associated Vasculitis. <i>In Vivo</i> , 2021, 35, 1761-1768.	1.3	1
21	Fibrosis-5 predicts end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis without substantial liver diseases. <i>Clinical and Experimental Medicine</i> , 2021, 21, 399-406.	3.6	4
22	Fibrinogen to albumin ratio reflects the activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23731.	2.1	7
23	Serum chitinase-3-like 1 protein is a useful biomarker to assess disease activity in ANCA-associated vasculitis: an observational study. <i>Arthritis Research and Therapy</i> , 2021, 23, 77.	3.5	8
24	Correlation between serum cysteine-rich protein 61 and disease activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2021, 40, 3703-3710.	2.2	1
25	Risk of Stroke in Systemic Necrotizing Vasculitis: A Nationwide Study Using the National Claims Database. <i>Frontiers in Immunology</i> , 2021, 12, 629902.	4.8	8
26	Reclassification of Korean patients with polymyositis and dermatomyositis based on the Bohan and Peter criteria by the 2017 European League Against Rheumatism/American College of Rheumatology classification criteria for adult and juvenile idiopathic inflammatory myopathies. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 441-446.	1.7	3
27	Unclassifiable repeated antineutrophil cytoplasmic antibody (ANCA) positivity in diseases other than ANCA-associated vasculitis. <i>Zeitschrift Fur Rheumatologie</i> , 2021, , 1.	1.0	4
28	Association between follistatin-related protein 1 and the functional status of patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Chinese Medical Journal</i> , 2021, 134, 1168-1174.	2.3	0
29	Clinical features of Korean elderly patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 731-741.	1.7	3
30	Novel mortality-predicting index at diagnosis can effectively predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e23885.	2.1	4
31	Clinical significance of large unstained cell count in estimating the current activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e14512.	1.7	3
32	Predictive Ability of Serum IL-27 Level for Assessing Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Mediators of Inflammation</i> , 2021, 2021, 1-8.	3.0	2
33	Serum adipokine profiles in patients with microscopic polyangiitis and granulomatosis with polyangiitis: An exploratory analysis. <i>PLoS ONE</i> , 2021, 16, e0254226.	2.5	1
34	A mass-like lesion encasing the aortic arch and descending aorta: immunoglobulin G4-related periaortitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1256-1257.	1.7	0
35	Evaluation of body composition using computed tomography in patients with anti-neutrophil cytoplasmic antibody-associated vasculitis. <i>Korean Journal of Internal Medicine</i> , 2021, 36, 1221-1232.	1.7	1
36	The novel fibrosis index at diagnosis may predict all-cause mortality in patients with antineutrophil cytoplasmic antibody-associated vasculitis without substantial liver diseases. <i>Clinics</i> , 2021, 76, e2501.	1.5	2

#	ARTICLE	IF	CITATIONS
37	Male Sex Is a Significant Predictor of All-cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis. <i>Journal of Korean Medical Science</i> , 2021, 36, e120.	2.5	8
38	The Efficacy of Mycophenolate Mofetil in Remission Maintenance Therapy for Microscopic Polyangiitis and Granulomatosis with Polyangiitis. <i>Yonsei Medical Journal</i> , 2021, 62, 494.	2.2	2
39	Efficacy of the fibrosis index for predicting end-stage renal disease in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>International Journal of Clinical Practice</i> , 2021, 75, e13929.	1.7	4
40	Serum Clusterin Level Could Reflect the Current Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2021, 62, 1016.	2.2	3
41	Serum progranulin as a predictive marker for high activity of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2021, 35, e24048.	2.1	1
42	A new index for distinguishing hypereosinophilic syndrome and antineutrophil cytoplasmic antibody-negative eosinophilic granulomatosis with polyangiitis. <i>Asian Pacific Journal of Allergy and Immunology</i> , 2021, , .	0.4	3
43	Serum granzyme B is associated with otorhinolaryngological, pulmonary, and renal involvement of antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Investigative Medicine</i> , 2021, 69, 91-95.	1.6	0
44	Aortic valve surgery in patients with Takayasu's arteritis: a nationwide analysis of 1,197 patients during a 9-year period. <i>Clinical and Experimental Rheumatology</i> , 2021, , .	0.8	0
45	Clinical impact of proteinase 3-antineutrophil cytoplasmic antibody positivity in eosinophilic granulomatosis with polyangiitis. <i>Korean Journal of Internal Medicine</i> , 2021, , .	1.7	1
46	Metabolic Syndrome Severity Score, Comparable to Serum Creatinine, Could Predict the Occurrence of End-Stage Kidney Disease in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Journal of Clinical Medicine</i> , 2021, 10, 5744.	2.4	5
47	Fibrosis-4 index at diagnosis can predict all-cause mortality in patients with rheumatoid arthritis: A retrospective monocentric study. <i>Modern Rheumatology</i> , 2020, 30, 70-77.	1.8	17
48	Multivariable index for assessing the activity and predicting all-cause mortality in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23022.	2.1	13
49	Can antineutrophil cytoplasmic antibody positivity at diagnosis predict the poor outcomes of Sjögren's syndrome?. <i>Rheumatology International</i> , 2020, 40, 1063-1070.	3.0	4
50	Clinical implication of chronic paranasal sinusitis for the classification of microscopic polyangiitis. <i>International Journal of Clinical Practice</i> , 2020, 74, e13431.	1.7	2
51	Double positivity for antineutrophil cytoplasmic antibody (ANCA) and anti-glomerular basement membrane antibody could predict end-stage renal disease in ANCA-associated vasculitis: a monocentric pilot study. <i>Clinical Rheumatology</i> , 2020, 39, 831-840.	2.2	2
52	Punicalagin Ameliorates Lupus Nephritis via Inhibition of PAR2. <i>International Journal of Molecular Sciences</i> , 2020, 21, 4975.	4.1	14
53	Incidence of Tuberculosis in Systemic Necrotizing Vasculitides: A Population-Based Study From an Intermediate-Burden Country. <i>Frontiers in Medicine</i> , 2020, 7, 550004.	2.6	0
54	Atherogenic index of plasma predicts cerebrovascular accident occurrence in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Lipids in Health and Disease</i> , 2020, 19, 184.	3.0	7

#	ARTICLE	IF	CITATIONS
55	Serum Amyloid A Is a Biomarker of Disease Activity and Health-Related Quality-of-Life in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Disease Markers</i> , 2020, 2020, 1-9.	1.3	4
56	Definite IgG4-related disease had no overlap with eosinophilic granulomatosis with polyangiitis in Korean patients: a pilot study in one centre. <i>Clinical Rheumatology</i> , 2020, 39, 3009-3015.	2.2	3
57	Association between the antineutrophil cytoplasmic antibody and late coronary arterial occlusive disease in patients with Takayasu arteritis. <i>Journal of Cardiology</i> , 2020, 76, 407-412.	1.9	3
58	Clinical implication of plasma exchange on life-threatening antineutrophil cytoplasmic antibody-associated vasculitis. <i>BMC Pulmonary Medicine</i> , 2020, 20, 147.	2.0	2
59	Clinical characteristics and long-term outcomes of Libmanâ€™Sacks endocarditis in patients with systemic lupus erythematosus. <i>Lupus</i> , 2020, 29, 1115-1120.	1.6	11
60	Non-histologic factors discriminating proliferative lupus nephritis from membranous lupus nephritis. <i>Arthritis Research and Therapy</i> , 2020, 22, 138.	3.5	5
61	Comparison of clinical features and outcomes between patients with early and delayed lupus nephritis. <i>BMC Nephrology</i> , 2020, 21, 258.	1.8	5
62	Hyperuricemia is associated with decreased renal function and occurrence of end-stage renal disease in patients with microscopic polyangiitis and granulomatosis with polyangiitis: a retrospective study. <i>Rheumatology International</i> , 2020, 40, 1089-1099.	3.0	7
63	Clinical characteristics associated with drug-free sustained remission in patients with rheumatoid arthritis: Data from Korean Intensive Management of Early Rheumatoid Arthritis (KIMERA). <i>Seminars in Arthritis and Rheumatism</i> , 2020, 50, 1414-1420.	3.4	3
64	Serum interleukin-16 significantly correlates with the Vasculitis Damage Index in antineutrophil cytoplasmic antibody-associated vasculitis. <i>Arthritis Research and Therapy</i> , 2020, 22, 73.	3.5	6
65	Serum Mannose-Binding Lectin Levels Are Correlated with the Disease Activity of Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Single-Center Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020, 251, 117-123.	1.2	7
66	Worse Renal Presentation and Prognosis in Initial-Onset Lupus Nephritis than Early-Onset Lupus Nephritis. <i>Yonsei Medical Journal</i> , 2020, 61, 951.	2.2	4
67	Rituximab Biosimilar Prevents Poor Outcomes of Microscopic Polyangiitis and Granulomatosis with Polyangiitis as Effectively as Rituximab Originator. <i>Yonsei Medical Journal</i> , 2020, 61, 712.	2.2	5
68	Soluble Lectin-Like Oxidized Low-Density Lipoprotein Receptor 1 Is Inversely Correlated with the Activity of ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2020, 61, 720.	2.2	2
69	Novel PAR2 antagonist ameliorates progression of lupus nephritis in NZB/Z F1 mice. <i>FASEB Journal</i> , 2020, 34, 1-1.	0.5	0
70	Pregnancy Morbidities in Korean Patients with Takayasu Arteritis: A Monocentric Pilot Study. <i>Yonsei Medical Journal</i> , 2020, 61, 970.	2.2	3
71	MiR-451 suppresses inflammatory responses in ankylosing spondylitis by targeting macrophage migration inhibitory factor. <i>Clinical and Experimental Rheumatology</i> , 2020, 38, 275-281.	0.8	4
72	Will the HALP score help to assess the activity and predict the prognosis of antineutrophil cytoplasmic antibody-associated vasculitis?. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 124, 236-237.	0.8	2

#	ARTICLE	IF	CITATIONS
73	Application of the 2019 classification criteria for systemic lupus erythematosus to patients with established ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2020, 38 Suppl 124, 243-244.	0.8	0
74	Clinical significance of ANCA positivity in patients with IgA vasculitis: a retrospective monocentric study. <i>Rheumatology International</i> , 2019, 39, 1927-1936.	3.0	4
75	Atializumab, a humanized anti-aminocyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1) antibody significantly improves nephritis in (NZB/NZW) F1 mice. <i>Biomaterials</i> , 2019, 220, 119408.	11.4	7
76	Predictor of depressive disorders in patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical Rheumatology</i> , 2019, 38, 3485-3491.	2.2	7
77	Evaluation of macrophage activation syndrome in hospitalised patients with Kikuchi-Fujimoto disease based on the 2016 EULAR/ACR/PRINTO classification criteria. <i>PLoS ONE</i> , 2019, 14, e0219970.	2.5	8
78	Could hypereosinophilia at diagnosis estimate the current activity or predict relapse in systemic immunosuppressive drug-naïve patients with eosinophilic granulomatosis with polyangiitis?. <i>Rheumatology International</i> , 2019, 39, 1899-1905.	3.0	3
79	Serum Aminoacyl-tRNA Synthetase-Interacting Multifunctional Protein-1 Can Predict Severe Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: A Pilot Monocentric Study. <i>BioMed Research International</i> , 2019, 2019, 1-6.	1.9	7
80	Comparison of the Clinical Implications among Five Different Nutritional Indices in Patients with Lupus Nephritis. <i>Nutrients</i> , 2019, 11, 1456.	4.1	14
81	Risk of Cancers in Antineutrophil Cytoplasmic Antibody-Associated Vasculitis: Results from the Korea National Health Insurance Claims Database 2010â€“2018. <i>Journal of Clinical Medicine</i> , 2019, 8, 1871.	2.4	10
82	Clinical characteristics of patients with systemic lupus erythematosus showing a false-positive result of syphilis screening. <i>Rheumatology International</i> , 2019, 39, 1859-1866.	3.0	4
83	Anti-Smith antibody is associated with disease activity in patients with new-onset systemic lupus erythematosus. <i>Rheumatology International</i> , 2019, 39, 1937-1944.	3.0	26
84	Fibrosis-4 index at diagnosis is associated with all-cause mortality in patients with microscopic polyangiitis and granulomatosis with polyangiitis. <i>BMC Gastroenterology</i> , 2019, 19, 90.	2.0	13
85	Anti-phospholipid antibody syndrome occurrence in patients with persistent anti-phospholipid antibodies. <i>Rheumatology International</i> , 2019, 39, 1359-1367.	3.0	4
86	Thyroid Dysfunction in Patients with Antineutrophil Cytoplasmic Antibody-associated Vasculitis: A Monocentric Retrospective Study. <i>Journal of Rheumatology</i> , 2019, 46, 1248-1250.	2.0	2
87	Should nasal biopsy inevitably be performed for classifying granulomatosis with polyangiitis in patients with rhinosinusitis? A retrospective chart review study. <i>Rheumatology International</i> , 2019, 39, 885-892.	3.0	3
88	Ex Vivo Interferon Gamma Production by Peripheral Immune Cells Predicts Survival in Lung Adenocarcinoma. <i>Clinical Lung Cancer</i> , 2019, 20, e299-e308.	2.6	2
89	Serum interleukin-21 positivity could indicate the current activity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical Rheumatology</i> , 2019, 38, 1685-1690.	2.2	5
90	Prognostic nutritional index is associated with disease severity and relapse in ANCA-associated vasculitis. <i>International Journal of Rheumatic Diseases</i> , 2019, 22, 797-804.	1.9	16

#	ARTICLE	IF	CITATIONS
91	ANCA positivity at the time of renal biopsy is associated with chronicity index of lupus nephritis. <i>Rheumatology International</i> , 2019, 39, 879-884.	3.0	9
92	Persistent antiphospholipid antibodies are associated with thrombotic events in ANCA-associated vasculitis: A retrospective monocentric study. <i>Nefrologia</i> , 2019, 39, 395-401.	0.4	6
93	125â€¦Treatment outcome in lupus nephritis patients treated with mycophenolate mofetil: from a real-world clinical practice. , 2019, , .		0
94	Clinical role of albumin to globulin ratio in microscopic polyangiitis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 487-494.	2.2	11
95	Low serum complement 3 level is associated with severe ANCA-associated vasculitis at diagnosis. <i>Clinical and Experimental Nephrology</i> , 2019, 23, 223-230.	1.6	20
96	Systemic immuneâ€inflammation index could estimate the crossâ€sectional high activity and the poor outcomes in immunosuppressive drugâ€naïve patients with antineutrophil cytoplasmic antibodyâ€associated vasculitis. <i>Nephrology</i> , 2019, 24, 711-717.	1.6	42
97	No overlap between IgG4-related disease and microscopic polyangiitis and granulomatosis with polyangiitis despite elevated serum IgG4 at diagnosis: a retrospective monocentric study. <i>Clinical Rheumatology</i> , 2019, 38, 1147-1154.	2.2	21
98	Antineutrophil Cytoplasmic Antibody-Associated Vasculitis in Korea: A Narrative Review. <i>Yonsei Medical Journal</i> , 2019, 60, 10.	2.2	27
99	Controlling Nutritional Status Score is Associated with All-Cause Mortality in Patients with Antineutrophil Cytoplasmic Antibody-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2019, 60, 1164.	2.2	8
100	Eosinophilic Granulomatosis with Polyangiitis: Experiences in Korean Patients. <i>Yonsei Medical Journal</i> , 2019, 60, 705.	2.2	11
101	Subclinical but significant liver fibrosis in patients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 26-31.	0.8	1
102	Serum soluble programmed cell death protein 1 could predict the current activity and severity of antineutrophil cytoplasmic antibody-associated vasculitis: a monocentric prospective study. <i>Clinical and Experimental Rheumatology</i> , 2019, 37 Suppl 117, 116-121.	0.8	2
103	Intranuclear delivery of the transcription modulation domain of Tbet-improved lupus nephritis in (NZB/NZW) F1 lupus-prone mice. <i>Kidney International</i> , 2018, 93, 1118-1130.	5.2	9
104	Birmingham vasculitis activity and chest manifestation at diagnosis can predict hospitalised infection in ANCA-associated vasculitis. <i>Clinical Rheumatology</i> , 2018, 37, 2133-2141.	2.2	12
105	Mean platelet volume can estimate the current vasculitis activity of microscopic polyangiitis. <i>Rheumatology International</i> , 2018, 38, 1095-1101.	3.0	9
106	Increased serum interleukinâ€32 levels in patients with Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 2167-2174.	1.9	20
107	Decreased muscle mass is independently associated with knee pain in female patients with radiographically mild osteoarthritis: a nationwide cross-sectional study (KNHANES 2010â€2011). <i>Clinical Rheumatology</i> , 2018, 37, 1333-1340.	2.2	15
108	Rheumatoid factor false positivity in patients with ANCA-associated vasculitis not having medical conditions producing rheumatoid factor. <i>Clinical Rheumatology</i> , 2018, 37, 2771-2779.	2.2	10

#	ARTICLE	IF	CITATIONS
109	Risk factors associated with inadequate control of disease activity in elderly patients with rheumatoid arthritis: Results from a nationwide KOran College of Rheumatology BIOlogics (KOBIO) registry. <i>PLoS ONE</i> , 2018, 13, e0205651.	2.5	13
110	Application of 2012 EULAR/ACR criteria for polymyalgia rheumatica to Korean patients previously classified by Chuang and Hunder criteria or Healey criteria. <i>International Journal of Rheumatic Diseases</i> , 2018, 21, 1838-1843.	1.9	2
111	HBsAg-negative and anti-HBc-positive in eosinophilic granulomatosis with polyangiitis: a retrospective pilot study. <i>Rheumatology International</i> , 2018, 38, 1531-1538.	3.0	5
112	Neutrophil to lymphocyte ratio at diagnosis can estimate vasculitis activity and poor prognosis in patients with ANCA-associated vasculitis: a retrospective study. <i>BMC Nephrology</i> , 2018, 19, 187.	1.8	32
113	Safety of Tocilizumab in Rheumatoid Arthritis Patients with Resolved Hepatitis B Virus Infection: Data from Real-World Experience. <i>Yonsei Medical Journal</i> , 2018, 59, 452.	2.2	27
114	Delta Neutrophil Index Is Associated with Vasculitis Activity and Risk of Relapse in ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 397.	2.2	16
115	Extravascular manifestations of Takayasu arteritis: focusing on the features shared with spondyloarthritis. <i>Arthritis Research and Therapy</i> , 2018, 20, 142.	3.5	20
116	Platelet to lymphocyte ratio is associated with the current activity of ANCA-associated vasculitis at diagnosis: a retrospective monocentric study. <i>Rheumatology International</i> , 2018, 38, 1865-1871.	3.0	28
117	C-Reactive Protein to Serum Albumin Ratio Is an Independent Predictor of All-Cause Mortality in Patients with ANCA-Associated Vasculitis. <i>Yonsei Medical Journal</i> , 2018, 59, 865.	2.2	36
118	Serum leucine-rich α_2 -glycoprotein is elevated in patients with systemic lupus erythematosus and correlates with disease activity. <i>Clinica Chimica Acta</i> , 2018, 486, 253-258.	1.1	10
119	Red blood cell distribution width is useful in discriminating adult onset Still's disease and sepsis within 24 hours after hospitalization. <i>Korean Journal of Internal Medicine</i> , 2018, 33, 1234-1240.	1.7	8
120	Reclassification of polyarteritis nodosa based on the 1990 ACR criteria using the 2007 EMA algorithm modified by the 2012 CHCC definitions. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 165-166.	0.8	2
121	The utility of the ACR/EULAR 2017 provisional classification criteria for granulomatosis with polyangiitis in Korean patients with antineutrophil cytoplasmic antibody-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 85-87.	0.8	14
122	Renal outcome of kidney-transplantation in Korean recipients with ANCA-associated vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 115-120.	0.8	2
123	Serum aminoacyl-tRNA synthetase-interacting multifunctional protein-1 (AIMP1), a novel disease activity predictive biomarker of systemic lupus erythematosus. <i>Clinical and Experimental Rheumatology</i> , 2018, 36, 533-539.	0.8	2
124	Cancer development in Korean patients with ANCA-associated vasculitis: a single centre study. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 73-77.	0.8	3
125	The initial predictors of death in 153 patients with ANCA-associated vasculitis in a single Korean centre. <i>Clinical and Experimental Rheumatology</i> , 2018, 36 Suppl 111, 65-72.	0.8	9
126	Glycated albumin increases with disease activity in rheumatoid factor positive rheumatoid arthritis patients with normal fasting glucose and HbA1c. <i>Joint Bone Spine</i> , 2017, 84, 115-118.	1.6	5

#	ARTICLE	IF	CITATIONS
127	Serum anti-lysozyme is associated with disease activity of Behçet's disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 261-268.	1.9	8
128	Five factor score of more than 1 is associated with relapse during the first 2 year follow up in patients with eosinophilic granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1261-1268.	1.9	12
129	In-hospital mortality in febrile lupus patients based on 2016 EULAR/ACR/PRINTO classification criteria for macrophage activation syndrome. <i>Seminars in Arthritis and Rheumatism</i> , 2017, 47, 216-221.	3.4	28
130	Evaluation of Spleen Glucose Metabolism Using ¹⁸ F-FDG PET/CT in Patients with Febrile Autoimmune Disease. <i>Journal of Nuclear Medicine</i> , 2017, 58, 507-513.	5.0	33
131	Application of the 2016 EULAR/ACR/PRINTO Classification Criteria for Macrophage Activation Syndrome in Patients with Adult-onset Still Disease. <i>Journal of Rheumatology</i> , 2017, 44, 996-1003.	2.0	43
132	Delta neutrophil index contributes to the differential diagnosis between acute gout attack and cellulitis within 24 hours after hospitalization. <i>Rheumatology</i> , 2017, 56, kew471.	1.9	11
133	Malignancies in Korean patients with immunoglobulin G4-related disease. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1028-1035.	1.9	35
134	Birmingham vasculitis activity score at diagnosis is a significant predictor of relapse of polyarteritis nodosa. <i>Rheumatology International</i> , 2017, 37, 685-694.	3.0	11
135	Echocardiographic features in patients with ANCA-associated vasculitis within 3 months before and after diagnosis. <i>Clinical Rheumatology</i> , 2017, 36, 2751-2759.	2.2	9
136	Cell-penetrating interactomic inhibition of nuclear factor-kappa B in a mouse model of postoperative cognitive dysfunction. <i>Scientific Reports</i> , 2017, 7, 13482.	3.3	27
137	Birmingham vasculitis activity score of more than 9.5 at diagnosis is an independent predictor of refractory disease in granulomatosis with polyangiitis. <i>International Journal of Rheumatic Diseases</i> , 2017, 20, 1593-1605.	1.9	17
138	Defective autophagy activity and its association with spinal damage in patients with ankylosing spondylitis. <i>Joint Bone Spine</i> , 2017, 84, 583-587.	1.6	12
139	Fever as an initial manifestation of spondyloarthritis: A retrospective study. <i>PLoS ONE</i> , 2017, 12, e0184323.	2.5	5
140	Decreased ex vivo production of interferon-gamma is associated with severity and poor prognosis in patients with lupus. <i>Arthritis Research and Therapy</i> , 2017, 19, 193.	3.5	14
141	Predictors of mortality in autoimmune disease patients with concurrent cytomegalovirus infections detected by quantitative real-time PCR. <i>PLoS ONE</i> , 2017, 12, e0181590.	2.5	9
142	Chest and renal involvements, Birmingham vascular activity score more than 13.5 and five factor score (1996) more than 1 at diagnosis are significant predictors of relapse of microscopic polyangiitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 47-54.	0.8	11
143	Clinical and prognostic features of Korean patients with MPO-ANCA, PR3-ANCA and ANCA-negative vasculitis. <i>Clinical and Experimental Rheumatology</i> , 2017, 35 Suppl 103, 111-118.	0.8	16
144	Anti-Sm is associated with the early poor outcome of lupus nephritis. <i>International Journal of Rheumatic Diseases</i> , 2016, 19, 897-902.	1.9	22

#	ARTICLE	IF	CITATIONS
145	Clinical predictors of silent but substantial liver fibrosis in primary Sjogren's syndrome. <i>Modern Rheumatology</i> , 2016, 26, 576-582.	1.8	6
146	Red nail folds and Gottron's sign in a patient with leukemic infiltration of pelvic muscles. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015, 13, 581-582.	0.8	0
147	Rote Nagelfalze und Gottron-Zeichen bei einem Patienten mit leukämischer Infiltration der Beckenmuskulatur. <i>JDDG - Journal of the German Society of Dermatology</i> , 2015, 13, 581-583.	0.8	0
148	A Case of Aortitis with Systemic Lupus Erythematosus. <i>Journal of Rheumatic Diseases</i> , 2015, 22, 205.	1.1	1
149	The antibody atliximab attenuates collagen-induced arthritis by neutralizing AIMP1, an inflammatory cytokine that enhances osteoclastogenesis. <i>Biomaterials</i> , 2015, 44, 45-54.	11.4	16
150	Application of the 2013 ACR/EULAR classification criteria for systemic sclerosis to patients with Raynaud's phenomenon. <i>Arthritis Research and Therapy</i> , 2015, 17, 77.	3.5	20
151	Disease duration and Medsger's severity score are associated with significant liver fibrosis in patients with systemic sclerosis. <i>Clinical and Experimental Rheumatology</i> , 2015, 33, S68-74.	0.8	5
152	Delta Neutrophil Index as an Early Marker for Differential Diagnosis of Adult-Onset Still's Disease and Sepsis. <i>Yonsei Medical Journal</i> , 2014, 55, 753.	2.2	26
153	Differences in Clinical Manifestations and Outcomes between Adult and Child Patients with Henoch-Schönlein Purpura. <i>Journal of Korean Medical Science</i> , 2014, 29, 198.	2.5	109
154	Usefulness of Serum Leucine-Rich Alpha-2 Glycoprotein as a Disease Activity Biomarker in Patients with Rheumatoid Arthritis. <i>Journal of Korean Medical Science</i> , 2014, 29, 1199.	2.5	37
155	ROR γ -specific transcriptional interaction suppresses autoimmunity associated with T _H 17 cells. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2014, 111, 18673-18678.	7.1	33
156	Inhibition of glycogen synthase kinase-3 β suppresses inflammatory responses in rheumatoid arthritis fibroblast-like synoviocytes and collagen-induced arthritis. <i>Joint Bone Spine</i> , 2014, 81, 240-246.	1.6	29
157	Secreted frizzled-related protein 5 suppresses inflammatory response in rheumatoid arthritis fibroblast-like synoviocytes through down-regulation of c-Jun N-terminal kinase. <i>Rheumatology</i> , 2014, 53, 1704-1711.	1.9	24
158	Proper time to initiate antiosteoporotic treatment in rheumatoid arthritis with or without glucocorticoid use. <i>Korean Journal of Internal Medicine</i> , 2014, 29, 434.	1.7	0
159	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013, 23, 509-515.	1.8	4
160	Soluble Receptor for Advanced Glycation End Products Alleviates Nephritis in (NZB/NZW)F1 Mice. <i>Arthritis and Rheumatism</i> , 2013, 65, 1902-1912.	6.7	23
161	Ventricular Tachyarrhythmias in a Patient with Andersen-Tawil Syndrome. <i>Korean Circulation Journal</i> , 2013, 43, 62.	1.9	2
162	Adjusted neutropenia is associated with early serious infection in systemic lupus erythematosus. <i>Modern Rheumatology</i> , 2013, 23, 509-515.	1.8	2

#	ARTICLE	IF	CITATIONS
163	Takayasu arteritis associated with ulcerative colitis and optic neuritis: first case in Korea. Korean Journal of Internal Medicine, 2013, 28, 491.	1.7	5
164	Attenuation of nephritis in lupus-prone mice by thalidomide. Rheumatology, 2012, 51, 2131-2140.	1.9	12
165	Alleviation of rheumatoid arthritis by cell-transducible methotrexate upon transcutaneous delivery. Biomaterials, 2012, 33, 1563-1572.	11.4	29
166	A Case of Systemic Lupus Erythematosus Manifested as Anemia and Thrombocytopenia during Antithyroid Drug Treatment. Korean Journal of Medicine, 2012, 82, 232.	0.3	0
167	Encephalitis due to Acute Human Immunodeficiency Virus Infection. Infection and Chemotherapy, 2012, 44, 540.	2.3	0
168	A Case of Ankylosing Spondylitis with Follicular Lymphoma. Journal of Rheumatic Diseases, 2011, 18, 315.	1.1	0
169	Clinical Manifestations of Korean Adult Patients with Henoch-Schönlein Purpura. The Journal of the Korean Rheumatism Association, 2010, 17, 133.	0.1	3
170	Comparison of therapeutic efficacy between bortezomib and combination treatment of prednisolone and mycophenolate mofetil on nephritis in NZB/WF1 mice. Clinical and Experimental Rheumatology, 2010, 28, 393-6.	0.8	7
171	E/E' ratio is more sensitive than E/A ratio for detection of left ventricular diastolic dysfunction in patients with systemic sclerosis. Clinical and Experimental Rheumatology, 2010, 28, S12-7.	0.8	17
172	Measurement of the serum leptin level could assist disease activity monitoring in rheumatoid arthritis. Rheumatology International, 2007, 27, 537-540.	3.0	89
173	Malignancies in Korean Patients with Inflammatory Myopathy. Yonsei Medical Journal, 2006, 47, 519.	2.2	34
174	Predicting the depressive status using empirical dietary inflammatory index in patients with antineutrophil cytoplasmic antibody-associated vasculitis. Journal of Clinical Laboratory Analysis, 0, ,	2.1	1