Christa Scheidt-Nave

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3306663/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	The challenge of comprehensively mapping children's health in a nation-wide health survey: Design of the German KiGGS-Study. BMC Public Health, 2008, 8, 196.	2.9	428
2	German health interview and examination survey for adults (DEGS) - design, objectives and implementation of the first data collection wave. BMC Public Health, 2012, 12, 730.	2.9	327
3	Vitamin D and mortality: Individual participant data meta-analysis of standardized 25-hydroxyvitamin D in 26916 individuals from a European consortium. PLoS ONE, 2017, 12, e0170791.	2.5	219
4	Higher Prevalence of Vitamin D Deficiency Is Associated with Immigrant Background among Children and Adolescents in Germany. Journal of Nutrition, 2008, 138, 1482-1490.	2.9	171
5	Simply no time? Barriers to GPs' participation in primary health care research. Family Practice, 2008, 25, 105-112.	1.9	161
6	Vitamin D status among adults in Germany – results from the German Health Interview and Examination Survey for Adults (DEGS1). BMC Public Health, 2015, 15, 641.	2.9	107
7	Prevalence and correlates of frailty among older adults: findings from the German health interview and examination survey. BMC Geriatrics, 2015, 15, 22.	2.7	93
8	Prevalence, trends, patterns and associations of analgesic use in Germany. BMC Pharmacology & Toxicology, 2015, 16, 28.	2.4	81
9	Fall incidence in Germany: results of two population-based studies, and comparison of retrospective and prospective falls data collection methods. BMC Geriatrics, 2014, 14, 105.	2.7	74
10	The Prevalence of Renal Failure. Deutsches Ärzteblatt International, 2016, 113, 85-91.	0.9	74
11	HuSKY: a healthy nutrition score based on food intake of children and adolescents in Germany. British Journal of Nutrition, 2009, 102, 610.	2.3	64
12	Changes in diabetes care indicators: findings from German National Health Interview and Examination Surveys 1997–1999 and 2008–2011. BMJ Open Diabetes Research and Care, 2015, 3, e000135.	2.8	57
13	Participation in a quality of care study and consequences for generalizability of general practice research. Family Practice, 2005, 22, 458-464.	1.9	48
14	Association Between Hemoglobin A1c and All-Cause Mortality: Results of the Mortality Follow-up of the German National Health Interview and Examination Survey 1998. Diabetes Care, 2015, 38, 249-256.	8.6	46
15	Type 2 Diabetes. Deutsches Ärzteblatt International, 2013, 110, 331-7.	0.9	45
16	Implications of standardization of serum 25-hydroxyvitamin D data for the evaluation of vitamin D status in Germany, including a temporal analysis. BMC Public Health, 2018, 18, 845.	2.9	44
17	Association Between Chronic Stress and Blood Pressure. Psychosomatic Medicine, 2015, 77, 575-582.	2.0	43
18	Time Trends in Cardiometabolic Risk Factors in Adults. Deutsches Ärzteblatt International, 2016, 113, 712-719.	0.9	40

CHRISTA SCHEIDT-NAVE

#	Article	IF	CITATIONS
19	All-cause mortality in adults with and without type 2 diabetes: findings from the national health monitoring in Germany. BMJ Open Diabetes Research and Care, 2017, 5, e000451.	2.8	39
20	Validation of the German Diabetes Risk Score among the general adult population: findings from the German Health Interview and Examination Surveys. BMJ Open Diabetes Research and Care, 2016, 4, e000280.	2.8	38
21	Modifiable cardiovascular risk factors in adults aged 40–79 years in Germany with and without prior coronary heart disease or stroke. BMC Public Health, 2015, 15, 701.	2.9	30
22	Differences in Menopausal Hormone Therapy Use among Women in Germany between 1998 and 2003. BMC Women's Health, 2007, 7, 19.	2.0	29
23	BURDEN 2020—Burden of disease in Germany at the national and regional level. Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz, 2018, 61, 1159-1166.	7.2	28
24	Major comorbid conditions in asthma and association with asthma-related hospitalizations and emergency department admissions in adults: results from the German national health telephone interview survey (GEDA) 2010. BMC Pulmonary Medicine, 2013, 13, 46.	2.0	26
25	Perceived diabetes risk and related determinants in individuals with high actual diabetes risk: results from a nationwide population-based survey. BMJ Open Diabetes Research and Care, 2019, 7, e000680.	2.8	25
26	â€~We can do only what we have the means for' general practitioners' views of primary care for older people with complex health problems. BMC Family Practice, 2015, 16, 35.	2.9	24
27	Age, maturation and serum lipid parameters: findings from the German Health Survey for Children and Adolescents. BMC Public Health, 2019, 19, 1627.	2.9	24
28	Changes in mean serum lipids among adults in Germany: results from National Health Surveys 1997-99 and 2008-11. BMC Public Health, 2016, 16, 240.	2.9	21
29	Chronische Erkrankungen – Epidemiologische Entwicklung und die Bedeutung für die Öffentliche Gesundheit. Public Health Forum, 2010, 18, 2-4.	0.2	20
30	Algorithm-Based Cholesterol Monitoring in Children with Type 1 Diabetes. Journal of Pediatrics, 2014, 164, 1079-1084.e2.	1.8	20
31	Operationalizing multimorbidity and autonomy for health services research in aging populations - the OMAHA study. BMC Health Services Research, 2011, 11, 47.	2.2	18
32	Longitudinal change in health-related quality of life in people with prevalent and incident type 2 diabetes compared to diabetes-free controls. PLoS ONE, 2017, 12, e0176895.	2.5	17
33	Physical health-related quality of life in relation to metabolic health and obesity among men and women in Germany. Health and Quality of Life Outcomes, 2017, 15, 122.	2.4	14
34	Psychometric properties of a nationwide survey for adults with and without diabetes: the "disease knowledge and information needs – diabetes mellitus (2017)―survey. BMC Public Health, 2020, 20, 192.	2.9	14
35	Excess mortality in adults with documented diabetes in Germany: routine data analysis of all insurance claims in Germany 2013–2014. BMJ Open, 2021, 11, e041508.	1.9	14
36	Care for the chronically ill in Germany - The challenges during the COVID-19 pandemic , 2021, 5, 2-27.		14

#	Article	IF	CITATIONS
37	Indicators of Overweight and Cardiovascular Disease Risk Factors among 11- to 17-Year-Old Boys and Girls in Germany. Obesity Facts, 2011, 4, 379-385.	3.4	13
38	Temporal changes in predicted risk of type 2 diabetes in Germany: findings from the German Health Interview and Examination Surveys 1997–1999 and 2008–2011. BMJ Open, 2017, 7, e013058.	1.9	12
39	Determinants of Change in Glycemic Status in Individuals with Prediabetes: Results from a Nationwide Cohort Study in Germany. Journal of Diabetes Research, 2018, 2018, 1-12.	2.3	11
40	Health App Use and Its Correlates Among Individuals With and Without Type 2 Diabetes: Nationwide Population-Based Survey. JMIR Diabetes, 2020, 5, e14396.	1.9	11
41	Impact of the estimation equation for GFR on population-based prevalence estimates of kidney dysfunction. BMC Nephrology, 2017, 18, 341.	1.8	10
42	Factors associated with undiagnosed type 2 diabetes in Germany: results from German Health Interview and Examination Survey for Adults 2008–2011. BMJ Open Diabetes Research and Care, 2020, 8, e001707.	2.8	10
43	Metabolic Health in Relation to Body Size: Changes in Prevalence over Time between 1997-99 and 2008-11 in Germany. PLoS ONE, 2016, 11, e0167159.	2.5	9
44	Asthma management practices in adults – findings from the German Health Update (GEDA) 2010 and the German National Health Interview and Examination Survey (DEGS1) 2008–2011. Journal of Asthma, 2016, 53, 50-61.	1.7	9
45	Are depressive symptoms associated with quality of care in diabetes? Findings from a nationwide population-based study. BMJ Open Diabetes Research and Care, 2021, 9, e001804.	2.8	9
46	Health situation of adults in Germany - Results for selected indicators from GEDA 2019/2020-EHIS , 2021, 6, 3-25.		8
47	Association between random glucose and all-cause mortality: findings from the mortality follow-up of the German National Health Interview and Examination Survey 1998. BMC Endocrine Disorders, 2018, 18, 95.	2.2	7
48	Indicator-based public health monitoring in old age in OECD member countries: a scoping review. BMC Public Health, 2019, 19, 1068.	2.9	7
49	Individual and area-level determinants associated with C-reactive protein as a marker of cardiometabolic risk among adults: Results from the German National Health Interview and Examination Survey 2008-2011. PLoS ONE, 2019, 14, e0211774.	2.5	7
50	Estimating the impact of tax policy interventions on the projected number and prevalence of adults with type 2 diabetes in Germany between 2020 and 2040. BMJ Open Diabetes Research and Care, 2021, 9, e001813.	2.8	7
51	Cardiovascular reactivity is independently associated with better mental health. Blood Pressure Monitoring, 2016, 21, 215-223.	0.8	6
52	Changes in Waist Circumference among German Adults over Time - Compiling Results of Seven Prospective Cohort Studies. Obesity Facts, 2016, 9, 332-343.	3.4	6
53	First results from the study 'Disease knowledge and information needs - Diabetes mellitus (2017)' , 2018, 3, 22-60.		6
54	Predicting risk of substantial weight gain in German adults—a multi-center cohort approach. European Journal of Public Health, 2017, 27, ckw216.	0.3	5

CHRISTA SCHEIDT-NAVE

#	Article	IF	CITATIONS
55	Non-utilisation of health care services during the COVID-19 pandemic: Results of the CoMoLo study , 2022, 7, 2-17.		5
56	Utilisation of outpatient medical services by people with diagnosed diabetes during the COVID-19 pandemic in Germany , 2021, 6, 51-58.		4
57	Older people at the beginning of the COVID-19 pandemic: A scoping review , 2021, 6, 2-37.		4
58	National public health system responses to diabetes and other important noncommunicable diseases. Bundesgesundheitsblatt - Gesundheitsforschung - Gesundheitsschutz, 2018, 61, 1300-1306.	7.2	3
59	Diabetes-Surveillance am Robert Koch-Institut – Modellprojekt für den Aufbau einer NCD-Surveillance in Deutschland. Public Health Forum, 2021, 29, 277-281.	0.2	3
60	Limitations in activities of daily living in old age in Germany and the EU - Results from the European Health Interview Survey (EHIS) 2 , 2019, 4, 48-56.		3
61	Sexâ€specific impact of major depressive disorder on 12â€year change in glycaemic status: Results from a nationwide cohort study of adults without diabetes in Germany. Diabetic Medicine, 2022, 39, e14767.	2.3	3
62	Social inequality and diabetes mellitus - developments over time among the adult population in Germany , 2019, 4, 11-28.		2
63	Gestational diabetes in Germany: Development of screening participation and prevalence , 2021, 6, 3-18.		2
64	An international comparison of noncommunicable disease reporting: the case of diabetes mellitus , 2019, 4, 64-85.		1
65	Providing actionable evidence in Public Health - The 2018 international workshop on evidence-based public health at the Robert Koch Institute, Berlin , 2020, 5, 3-6.		1
66	Addressing health and ageing in the German national health monitoring system. BMC Proceedings, 2013, 7, S12.	1.6	0
67	Self-assessed quality of care among adults with diagnosed diabetes in Germany , 2021, 6, 36-42.		0

Diabetes surveillance - status and perspectives.. , 2018, 3, 10.

0