

Siqing Fu

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3321238/publications.pdf>

Version: 2024-02-01

171
papers

7,947
citations

53794

45
h-index

60623

81
g-index

177
all docs

177
docs citations

177
times ranked

12627
citing authors

#	ARTICLE	IF	CITATIONS
1	Validation of Prognostic Scores in Patients With Metastatic Urothelial Cancer Enrolling in Phase I Targeted Therapy or Next Generation Immunotherapy Trials. <i>Clinical Genitourinary Cancer</i> , 2022, 20, e16-e24.	1.9	1
2	Supportive care for the prevention of nausea, vomiting and anorexia in a phase 1B study of selinexor in advanced cancer patients: an exploratory study. <i>Investigational New Drugs</i> , 2022, 40, 124-133.	2.6	2
3	Phase 1 trial of ADI-PEG 20 and liposomal doxorubicin in patients with metastatic solid tumors. <i>Cancer Medicine</i> , 2022, 11, 340-347.	2.8	13
4	Phase Ib Study of Navicixizumab Plus Paclitaxel in Patients With Platinum-Resistant Ovarian, Primary Peritoneal, or Fallopian Tube Cancer. <i>Journal of Clinical Oncology</i> , 2022, 40, 2568-2577.	1.6	18
5	Clinical characteristics and outcomes of phase I cancer patients with CCNE1 amplification: MD Anderson experiences. <i>Scientific Reports</i> , 2022, 12, .	3.3	7
6	Longitudinal Monitoring of Circulating Tumor DNA to Predict Treatment Outcomes in Advanced Cancers. <i>JCO Precision Oncology</i> , 2022, , .	3.0	15
7	Molecular Profiling of Metastatic Bladder Cancer Early-Phase Clinical Trial Participants Predicts Patient Outcomes. <i>Molecular Cancer Research</i> , 2021, 19, 395-402.	3.4	7
8	Phase I Study of Everolimus, Letrozole, and Trastuzumab in Patients with Hormone Receptor ⁺ positive Metastatic Breast Cancer or Other Solid Tumors. <i>Clinical Cancer Research</i> , 2021, 27, 1247-1255.	7.0	5
9	Phase 1 trial of ADI-PEG20 plus cisplatin in patients with pretreated metastatic melanoma or other advanced solid malignancies. <i>British Journal of Cancer</i> , 2021, 124, 1533-1539.	6.4	20
10	Associations between the gut microbiome and fatigue in cancer patients. <i>Scientific Reports</i> , 2021, 11, 5847.	3.3	24
11	Precision medicine: preliminary results from the Initiative for Molecular Profiling and Advanced Cancer Therapy 2 (IMPACT2) study. <i>Npj Precision Oncology</i> , 2021, 5, 21.	5.4	12
12	Overview of Ocular Side Effects of Selinexor. <i>Oncologist</i> , 2021, 26, 619-623.	3.7	5
13	Patient-Reported Out-of-Pocket Costs and Financial Toxicity During Early-Phase Oncology Clinical Trials. <i>Oncologist</i> , 2021, 26, 588-596.	3.7	42
14	Implementation of a Novel Web-Based Lesion Selection Tool to Improve Acquisition of Tumor Biopsy Specimens. <i>Journal of Immunotherapy and Precision Oncology</i> , 2021, 4, 45-52.	1.4	5
15	A phase I study of the WT2725 dosing emulsion in patients with advanced malignancies. <i>Scientific Reports</i> , 2021, 11, 22355.	3.3	5
16	Evaluating the psychometric properties of the Immunotherapy module of the MD Anderson Symptom Inventory. , 2020, 8, e000931.		11
17	Cell-free Circulating Tumor DNA Variant Allele Frequency Associates with Survival in Metastatic Cancer. <i>Clinical Cancer Research</i> , 2020, 26, 1924-1931.	7.0	50
18	Phase I studies of vorinostat with ixazomib or pazopanib imply a role of antiangiogenesis-based therapy for TP53 mutant malignancies. <i>Scientific Reports</i> , 2020, 10, 3080.	3.3	10

#	ARTICLE	IF	CITATIONS
19	Dual EGFR blockade with cetuximab and erlotinib combined with anti-VEGF antibody bevacizumab in advanced solid tumors: a phase 1 dose escalation triplet combination trial. <i>Experimental Hematology and Oncology</i> , 2020, 9, 7.	5.0	11
20	Pharmacokinetics of the Investigational Aurora A Kinase Inhibitor Alisertib in Adult Patients With Advanced Solid Tumors or Relapsed/Refractory Lymphoma With Varying Degrees of Hepatic Dysfunction. <i>Journal of Clinical Pharmacology</i> , 2019, 59, 1204-1215.	2.0	3
21	Long-term overall survival and prognostic score predicting survival: the IMPACT study in precision medicine. <i>Journal of Hematology and Oncology</i> , 2019, 12, 145.	17.0	35
22	Cancer-Related Internet Use and Its Association With Patient Decision Making and Trust in Physicians Among Patients in an Early Drug Development Clinic: A Questionnaire-Based Cross-Sectional Observational Study. <i>Journal of Medical Internet Research</i> , 2019, 21, e10348.	4.3	13
23	Development of a prognostic scoring system for patients with advanced cancer enrolled in immune checkpoint inhibitor phase 1 clinical trials. <i>British Journal of Cancer</i> , 2018, 118, 763-769.	6.4	28
24	Predicting outcomes in patients with advanced non-small cell lung cancer enrolled in early phase immunotherapy trials. <i>Lung Cancer</i> , 2018, 120, 137-141.	2.0	29
25	Phase I study of nab-paclitaxel, gemcitabine, and bevacizumab in patients with advanced cancers. <i>British Journal of Cancer</i> , 2018, 118, 1419-1424.	6.4	7
26	Liquid Biopsies Using Plasma Exosomal Nucleic Acids and Plasma Cell-Free DNA Compared with Clinical Outcomes of Patients with Advanced Cancers. <i>Clinical Cancer Research</i> , 2018, 24, 181-188.	7.0	127
27	Phase I study of the combination of crizotinib (as a MET inhibitor) and dasatinib (as a c-SRC inhibitor) in patients with advanced cancer. <i>Investigational New Drugs</i> , 2018, 36, 416-423.	2.6	17
28	Radiomics to predict immunotherapy-induced pneumonitis: proof of concept. <i>Investigational New Drugs</i> , 2018, 36, 601-607.	2.6	90
29	Phase I Dose-Escalation Study of Anti-CTLA-4 Antibody Ipilimumab and Lenalidomide in Patients with Advanced Cancers. <i>Molecular Cancer Therapeutics</i> , 2018, 17, 671-676.	4.1	33
30	Incidence of immune-related adverse events and its association with treatment outcomes: the MD Anderson Cancer Center experience. <i>Investigational New Drugs</i> , 2018, 36, 638-646.	2.6	149
31	A phase I clinical trial of hepatic arterial infusion of oxaliplatin and oral capecitabine, with or without intravenous bevacizumab, in patients with advanced cancer and predominant liver involvement. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 877-885.	2.3	5
32	A phase I study of LY3164530, a bispecific antibody targeting MET and EGFR, in patients with advanced or metastatic cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2018, 82, 407-418.	2.3	46
33	Strategic development of AZD1775, a Wee1 kinase inhibitor, for cancer therapy. <i>Expert Opinion on Investigational Drugs</i> , 2018, 27, 741-751.	4.1	43
34	Outcome analysis of Phase I trial patients with metastatic KRAS and/or TP53 mutant non-small cell lung cancer. <i>Oncotarget</i> , 2018, 9, 33258-33270.	1.8	9
35	Incidence of infusion reactions to anti-neoplastic agents in early phase clinical trials: The MD Anderson Cancer Center experience. <i>Investigational New Drugs</i> , 2017, 35, 59-67.	2.6	10
36	Development and Validation of an Ultradeep Next-Generation Sequencing Assay for Testing of Plasma Cell-Free DNA from Patients with Advanced Cancer. <i>Clinical Cancer Research</i> , 2017, 23, 5648-5656.	7.0	50

#	ARTICLE	IF	CITATIONS
37	Insurance Clearance for Early-Phase Oncology Clinical Trials Following the Affordable Care Act. <i>Clinical Cancer Research</i> , 2017, 23, 4155-4162.	7.0	4
38	Replication Stress Leading to Apoptosis within the S-phase Contributes to Synergism between Vorinostat and AZD1775 in HNSCC Harboring High-Risk <i>TP53</i> Mutation. <i>Clinical Cancer Research</i> , 2017, 23, 6541-6554.	7.0	27
39	Outcomes of patients with sarcoma enrolled in clinical trials of pazopanib combined with histone deacetylase, mTOR, Her2, or MEK inhibitors. <i>Scientific Reports</i> , 2017, 7, 15963.	3.3	21
40	Phase I trial of MEK 1/2 inhibitor pimasertib combined with mTOR inhibitor temsirolimus in patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2017, 35, 616-626.	2.6	22
41	Phase I clinical trial of combination imatinib and ipilimumab in patients with advanced malignancies. , 2017, 5, 35.		58
42	Post-Discharge Survival Outcomes of Patients with Advanced Cancer from the University of Texas MD Anderson Cancer Center Investigational Cancer Therapeutics (Phase I Trials) Inpatient Unit. <i>Oncology</i> , 2017, 92, 14-20.	1.9	3
43	Outcomes of patients ≥65 years old with advanced cancer treated on phase I trials at MD ANDERSON CANCER CENTER. <i>International Journal of Cancer</i> , 2017, 140, 208-215.	5.1	7
44	Ipilimumab with Stereotactic Ablative Radiation Therapy: Phase I Results and Immunologic Correlates from Peripheral T Cells. <i>Clinical Cancer Research</i> , 2017, 23, 1388-1396.	7.0	261
45	Characteristics and outcomes of patients with advanced sarcoma enrolled in early phase immunotherapy trials. , 2017, 5, 100.		114
46	Initiative for Molecular Profiling and Advanced Cancer Therapy (IMPACT): An MD Anderson Precision Medicine Study. <i>JCO Precision Oncology</i> , 2017, 2017, 1-18.	3.0	107
47	First-in-human trial of multikinase VEGF inhibitor regorafenib and anti-EGFR antibody cetuximab in advanced cancer patients. <i>JCI Insight</i> , 2017, 2, .	5.0	26
48	Antiangiogenesis and gene aberration-related therapy may improve overall survival in patients with concurrent KRAS and TP53 hotspot mutant cancer. <i>Oncotarget</i> , 2017, 8, 33796-33806.	1.8	5
49	Outcomes of phase I clinical trials for patients with advanced pancreatic cancer: update of the MD Anderson Cancer Center experience. <i>Oncotarget</i> , 2017, 8, 87163-87173.	1.8	0
50	Evaluation of Novel Targeted Therapies in Aggressive Biology Sarcoma Patients after progression from US FDA approved Therapies. <i>Scientific Reports</i> , 2016, 6, 35448.	3.3	12
51	Survival of patients with metastatic leiomyosarcoma: the MD Anderson Clinical Center for targeted therapy experience. <i>Cancer Medicine</i> , 2016, 5, 3437-3444.	2.8	20
52	Cancer Therapy Directed by Comprehensive Genomic Profiling: A Single Center Study. <i>Cancer Research</i> , 2016, 76, 3690-3701.	0.9	203
53	Phase I clinical trial of lenalidomide in combination with bevacizumab in patients with advanced cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 1097-1102.	2.3	5
54	Phase I dose escalation study of temsirolimus in combination with metformin in patients with advanced/refractory cancers. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 973-977.	2.3	34

#	ARTICLE	IF	CITATIONS
55	Pharmacokinetics of ixazomib, an oral proteasome inhibitor, in solid tumour patients with moderate or severe hepatic impairment. <i>British Journal of Clinical Pharmacology</i> , 2016, 82, 728-738.	2.4	38
56	<i>TP53</i> Alterations Correlate with Response to VEGF/VEGFR Inhibitors: Implications for Targeted Therapeutics. <i>Molecular Cancer Therapeutics</i> , 2016, 15, 2475-2485.	4.1	73
57	Sleep quality and its association with fatigue, symptom burden, and mood in patients with advanced cancer in a clinic for early-phase oncology clinical trials. <i>Cancer</i> , 2016, 122, 3401-3409.	4.1	50
58	Phase IB Study of Vemurafenib in Combination with Irinotecan and Cetuximab in Patients with Metastatic Colorectal Cancer with <i>BRAF</i> V600E Mutation. <i>Cancer Discovery</i> , 2016, 6, 1352-1365.	9.4	192
59	<i>BRAF</i> Mutation Testing in Cell-Free DNA from the Plasma of Patients with Advanced Cancers Using a Rapid, Automated Molecular Diagnostics System. <i>Molecular Cancer Therapeutics</i> , 2016, 15, 1397-1404.	4.1	78
60	Phase I clinical trial of lenalidomide in combination with 5-fluorouracil, leucovorin, and oxaliplatin in patients with advanced cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 575-581.	2.3	7
61	Pharmacokinetic evaluation of nanoparticle albumin-bound paclitaxel delivered via hepatic arterial infusion in patients with predominantly hepatic metastases. <i>Cancer Chemotherapy and Pharmacology</i> , 2016, 77, 357-364.	2.3	4
62	An overview of tyrosine kinase inhibitors for the treatment of epithelial ovarian cancer. <i>Expert Opinion on Investigational Drugs</i> , 2016, 25, 15-30.	4.1	8
63	Phase I dose-escalation study of the mTOR inhibitor sirolimus and the HDAC inhibitor vorinostat in patients with advanced malignancy. <i>Oncotarget</i> , 2016, 7, 67521-67531.	1.8	44
64	Advanced malignancies treated with a combination of the VEGF inhibitor bevacizumab, anti-EGFR antibody cetuximab, and the mTOR inhibitor temsirolimus. <i>Oncotarget</i> , 2016, 7, 23227-23238.	1.8	23
65	Continuous anti-angiogenic therapy after tumor progression in patients with recurrent high-grade epithelial ovarian cancer: phase I trial experience. <i>Oncotarget</i> , 2016, 7, 35132-35143.	1.8	9
66	Characteristics and outcomes for patients with advanced vaginal or vulvar cancer referred to a phase I clinical trials program: the MD Anderson cancer center experience. <i>Gynecologic Oncology Research and Practice</i> , 2015, 2, 10.	3.6	11
67	Actionable mutations in plasma cell-free DNA in patients with advanced cancers referred for experimental targeted therapies. <i>Oncotarget</i> , 2015, 6, 12809-12821.	1.8	86
68	Targeting drug transport mechanisms for improving platinum-based cancer chemotherapy. <i>Expert Opinion on Therapeutic Targets</i> , 2015, 19, 1307-1317.	3.4	36
69	Retreatment with anti-EGFR based therapies in metastatic colorectal cancer: impact of intervening time interval and prior anti-EGFR response. <i>BMC Cancer</i> , 2015, 15, 713.	2.6	43
70	The Prevalence and Impact of Hyperglycemia and Hyperlipidemia in Patients With Advanced Cancer Receiving Combination Treatment With the Mammalian Target of Rapamycin Inhibitor Temsirolimus and Insulin Growth Factor-Receptor Antibody Cixutumumab. <i>Oncologist</i> , 2015, 20, 737-741.	3.7	11
71	Olanzapine for cachexia in patients with advanced cancer: an exploratory study of effects on weight and metabolic cytokines. <i>Supportive Care in Cancer</i> , 2015, 23, 2649-2654.	2.2	26
72	Phase I study of pazopanib and vorinostat: a therapeutic approach for inhibiting mutant p53-mediated angiogenesis and facilitating mutant p53 degradation. <i>Annals of Oncology</i> , 2015, 26, 1012-1018.	1.2	56

#	ARTICLE	IF	CITATIONS
73	Dual antiangiogenic inhibition: a phase I dose escalation and expansion trial targeting VEGF-A and VEGFR in patients with advanced solid tumors. <i>Investigational New Drugs</i> , 2015, 33, 215-224.	2.6	8
74	A phase I trial of combination trastuzumab, lapatinib, and bevacizumab in patients with advanced cancer. <i>Investigational New Drugs</i> , 2015, 33, 177-186.	2.6	25
75	Phase I Dose-Escalation Study of the Multikinase Inhibitor Lenvatinib in Patients with Advanced Solid Tumors and in an Expanded Cohort of Patients with Melanoma. <i>Clinical Cancer Research</i> , 2015, 21, 4801-4810.	7.0	63
76	Phase I combination of pazopanib and everolimus in PIK3CA mutation positive/PTEN loss patients with advanced solid tumors refractory to standard therapy. <i>Investigational New Drugs</i> , 2015, 33, 700-709.	2.6	12
77	Phase I trial of valproic acid and lenalidomide in patients with advanced cancer. <i>Cancer Chemotherapy and Pharmacology</i> , 2015, 75, 869-874.	2.3	26
78	Xilonix, a novel true human antibody targeting the inflammatory cytokine interleukin-1 alpha, in non-small cell lung cancer. <i>Investigational New Drugs</i> , 2015, 33, 621-631.	2.6	63
79	Phase I study of azacitidine and oxaliplatin in patients with advanced cancers that have relapsed or are refractory to any platinum therapy. <i>Clinical Epigenetics</i> , 2015, 7, 29.	4.1	13
80	Dose-finding study of hepatic arterial infusion of irinotecan-based treatment in patients with advanced cancers metastatic to the liver. <i>Investigational New Drugs</i> , 2015, 33, 911-920.	2.6	7
81	MET Abnormalities in Patients With Genitourinary Malignancies and Outcomes With c-MET Inhibitors. <i>Clinical Genitourinary Cancer</i> , 2015, 13, e19-e26.	1.9	18
82	Prospective study comparing outcomes in patients with advanced malignancies on molecular alteration-matched versus non-matched therapy. <i>Journal of Clinical Oncology</i> , 2015, 33, 11019-11019.	1.6	8
83	Clinical next generation sequencing to identify actionable aberrations in a phase I program. <i>Oncotarget</i> , 2015, 6, 20099-20110.	1.8	41
84	A first-in-human study of AMG 208, an oral MET inhibitor, in adult patients with advanced solid tumors. <i>Oncotarget</i> , 2015, 6, 18693-18706.	1.8	24
85	Exploring response signals and targets in aggressive unresectable hepatocellular carcinoma: an analysis of targeted therapy phase 1 trials. <i>Oncotarget</i> , 2015, 6, 28453-28462.	1.8	9
86	<i>BRAF</i> mutation testing with a rapid, fully integrated molecular diagnostics system. <i>Oncotarget</i> , 2015, 6, 26886-26894.	1.8	45
87	Relative bioavailability of a prototype oral solution of the Aurora A kinase inhibitor alisertib (MLN8237) in patients with advanced solid tumors. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2015, 53, 563-572.	0.6	8
88	Triple-Negative Breast Cancer Patients Treated at MD Anderson Cancer Center in Phase I Trials: Improved Outcomes with Combination Chemotherapy and Targeted Agents. <i>Molecular Cancer Therapeutics</i> , 2014, 13, 3175-3184.	4.1	31
89	Analysis of 1,115 Patients Tested for <i>MET</i> Amplification and Therapy Response in the MD Anderson Phase I Clinic. <i>Clinical Cancer Research</i> , 2014, 20, 6336-6345.	7.0	70
90	Synergy Between VEGF/VEGFR Inhibitors and Chemotherapy Agents in the Phase I Clinic. <i>Clinical Cancer Research</i> , 2014, 20, 5956-5963.	7.0	10

#	ARTICLE	IF	CITATIONS
91	Incidence of Mucositis in Patients Treated With Temsirolimus-Based Regimens and Correlation to Treatment Response. <i>Oncologist</i> , 2014, 19, 426-428.	3.7	12
92	Assessing PIK3CA and PTEN in Early-Phase Trials with PI3K/AKT/mTOR Inhibitors. <i>Cell Reports</i> , 2014, 6, 377-387.	6.4	210
93	Phase I clinical trial of lenalidomide in combination with sorafenib in patients with advanced cancer. <i>Investigational New Drugs</i> , 2014, 32, 279-286.	2.6	14
94	Phase I study of anti-VEGF monoclonal antibody bevacizumab and histone deacetylase inhibitor valproic acid in patients with advanced cancers. <i>Cancer Chemotherapy and Pharmacology</i> , 2014, 73, 495-501.	2.3	46
95	Phase I dose-escalating study of TAS-106 in combination with carboplatin in patients with solid tumors. <i>Investigational New Drugs</i> , 2014, 32, 154-159.	2.6	8
96	Analysis of MET Genetic Aberrations in Patients With Breast Cancer at MD Anderson Phase I Unit. <i>Clinical Breast Cancer</i> , 2014, 14, 468-474.	2.4	29
97	Personalized Medicine for Patients with Advanced Cancer in the Phase I Program at MD Anderson: Validation and Landmark Analyses. <i>Clinical Cancer Research</i> , 2014, 20, 4827-4836.	7.0	186
98	Exploratory study of carboplatin plus the copper-lowering agent trientine in patients with advanced malignancies. <i>Investigational New Drugs</i> , 2014, 32, 465-472.	2.6	31
99	Evaluation of a novel blood pressure scoring method and its association with clinical response in cancer patients treated with anti-vascular endothelial growth factor therapy. <i>Investigational New Drugs</i> , 2014, 32, 717-722.	2.6	3
100	Dual inhibition of the vascular endothelial growth factor pathway: A phase 1 trial evaluating bevacizumab and AZD2171 (cediranib) in patients with advanced solid tumors. <i>Cancer</i> , 2014, 120, 2164-2173.	4.1	27
101	MABp1, a first-in-class true human antibody targeting interleukin-1 β in refractory cancers: an open-label, phase 1 dose-escalation and expansion study. <i>Lancet Oncology</i> , The, 2014, 15, 656-666.	10.7	178
102	Phase I Clinical Trial of Bendamustine and Bevacizumab for Patients With Advanced Cancer. <i>Journal of the National Comprehensive Cancer Network: JNCCN</i> , 2014, 12, 194-203.	4.9	4
103	Dual EGFR Inhibition in combination with anti-VEGF treatment in colorectal cancer. <i>Oncoscience</i> , 2014, 1, 540-549.	2.2	22
104	Anastrozole and everolimus in advanced gynecologic and breast malignancies: activity and molecular alterations in the PI3K/AKT/mTOR pathway. <i>Oncotarget</i> , 2014, 5, 3029-3038.	1.8	40
105	<i>MET</i> aberrations and c-MET inhibitors in patients with gastric and esophageal cancers in a phase I unit. <i>Oncotarget</i> , 2014, 5, 1837-1845.	1.8	27
106	Advanced gynecologic malignancies treated with a combination of the VEGF inhibitor bevacizumab and the mTOR inhibitor temsirolimus. <i>Oncotarget</i> , 2014, 5, 1846-1855.	1.8	28
107	Unique molecular signatures as a hallmark of patients with metastatic breast cancer: Implications for current treatment paradigms. <i>Oncotarget</i> , 2014, 5, 2349-2354.	1.8	54
108	Characteristics and survival of patients with advanced cancer and p53 mutations. <i>Oncotarget</i> , 2014, 5, 3871-3879.	1.8	11

#	ARTICLE	IF	CITATIONS
109	Outcomes of patients with advanced cancer and KRAS mutations in phase I clinical trials. <i>Oncotarget</i> , 2014, 5, 8937-8946.	1.8	6
110	Targeted PI3K/AKT/mTOR therapy for metastatic carcinomas of the cervix: A phase I clinical experience. <i>Oncotarget</i> , 2014, 5, 11168-11179.	1.8	53
111	Outcomes of patients with metastatic cervical cancer in a phase I clinical trials program. <i>Anticancer Research</i> , 2014, 34, 2349-55.	1.1	8
112	Dose-finding study of hepatic arterial infusion of oxaliplatin-based treatment in patients with advanced solid tumors metastatic to the liver. <i>Cancer Chemotherapy and Pharmacology</i> , 2013, 71, 389-397.	2.3	13
113	<i>PIK3CA</i> Mutation H1047R Is Associated with Response to PI3K/AKT/mTOR Signaling Pathway Inhibitors in Early-Phase Clinical Trials. <i>Cancer Research</i> , 2013, 73, 276-284.	0.9	262
114	Methylation and histone deacetylase inhibition in combination with platinum treatment in patients with advanced malignancies. <i>Investigational New Drugs</i> , 2013, 31, 1192-1200.	2.6	51
115	Retreatment after Secondary Resistance or Mixed Response: A Pilot Study. <i>Oncology</i> , 2013, 85, 350-355.	1.9	4
116	Weekly <i>nab</i> -Rapamycin in Patients with Advanced Nonhematologic Malignancies: Final Results of a Phase I Trial. <i>Clinical Cancer Research</i> , 2013, 19, 5474-5484.	7.0	72
117	Barriers to Study Enrollment in Patients With Advanced Cancer Referred to a Phase I Clinical Trials Unit. <i>Oncologist</i> , 2013, 18, 1315-1320.	3.7	20
118	Target-Based Therapeutic Matching in Early-Phase Clinical Trials in Patients with Advanced Colorectal Cancer and <i>PIK3CA</i> Mutations. <i>Molecular Cancer Therapeutics</i> , 2013, 12, 2857-2863.	4.1	42
119	Reply to M. Rouanne et al. <i>Journal of Clinical Oncology</i> , 2013, 31, 818-818.	1.6	0
120	Combining Erlotinib and Cetuximab Is Associated with Activity in Patients with Non-Small Cell Lung Cancer (Including Squamous Cell Carcinomas) and Wild-Type EGFR or Resistant Mutations. <i>Molecular Cancer Therapeutics</i> , 2013, 12, 2167-2175.	4.1	33
121	Enhanced Cytotoxic Effects of Combined Valproic Acid and the Aurora Kinase Inhibitor VE465 on Gynecologic Cancer Cells. <i>Frontiers in Oncology</i> , 2013, 3, 58.	2.8	23
122	MET nucleotide variations and amplification in advanced ovarian cancer: characteristics and outcomes with c-Met inhibitors. <i>Oncoscience</i> , 2013, 1, 5-13.	2.2	25
123	Germline <i>PTPRD</i> Mutations in Ewing Sarcoma: Biologic and Clinical Implications. <i>Oncotarget</i> , 2013, 4, 884-889.	1.8	24
124	Revisiting Clinical Trials Using EGFR Inhibitor-Based Regimens in Patients with Advanced Non-Small Cell Lung Cancer: A Retrospective Analysis of an MD Anderson Cancer Center Phase I Population. <i>Oncotarget</i> , 2013, 4, 772-784.	1.8	16
125	Dual EGFR inhibition in combination with anti-VEGF treatment: A phase I clinical trial in non-small cell lung cancer. <i>Oncotarget</i> , 2013, 4, 118-127.	1.8	33
126	Targeted Therapy of Advanced Gallbladder Cancer and Cholangiocarcinoma with Aggressive Biology: Eliciting Early Response Signals from Phase I trials. <i>Oncotarget</i> , 2013, 4, 153-162.	1.8	31

#	ARTICLE	IF	CITATIONS
127	P53 Mutations in Advanced Cancers: Clinical Characteristics, Outcomes, and Correlation between Progression-Free Survival and Bevacizumab-Containing Therapy. <i>Oncotarget</i> , 2013, 4, 705-714.	1.8	96
128	Survival of 1,181 Patients in a Phase I Clinic: The MD Anderson Clinical Center for Targeted Therapy Experience. <i>Clinical Cancer Research</i> , 2012, 18, 2922-2929.	7.0	78
129	Aurora kinase inhibitor VE 465 synergistically enhances cytotoxicity of carboplatin in ovarian cancer cells through induction of apoptosis and downregulation of histone 3. <i>Cancer Biology and Therapy</i> , 2012, 13, 1034-1041.	3.4	13
130	Role of the Human High-Affinity Copper Transporter in Copper Homeostasis Regulation and Cisplatin Sensitivity in Cancer Chemotherapy. <i>Cancer Research</i> , 2012, 72, 4616-4621.	0.9	85
131	Personalized Medicine in a Phase I Clinical Trials Program: The MD Anderson Cancer Center Initiative. <i>Clinical Cancer Research</i> , 2012, 18, 6373-6383.	7.0	458
132	Mechanistic Basis for Overcoming Platinum Resistance Using Copper Chelating Agents. <i>Molecular Cancer Therapeutics</i> , 2012, 11, 2483-2494.	4.1	67
133	Advance Care Planning in Patients With Cancer Referred to a Phase I Clinical Trials Program: The MD Anderson Cancer Center Experience. <i>Journal of Clinical Oncology</i> , 2012, 30, 2891-2896.	1.6	25
134	Overcoming Platinum Resistance through the Use of a Copper-Lowering Agent. <i>Molecular Cancer Therapeutics</i> , 2012, 11, 1221-1225.	4.1	70
135	Phase I Study of the Antiangiogenic Antibody Bevacizumab and the mTOR/Hypoxia-Inducible Factor Inhibitor Temsirolimus Combined with Liposomal Doxorubicin: Tolerance and Biological Activity. <i>Clinical Cancer Research</i> , 2012, 18, 5796-5805.	7.0	77
136	Outcomes in 144 Patients With Colorectal Cancer Treated in a Phase I Clinic: The MD Anderson Cancer Center Experience. <i>Clinical Colorectal Cancer</i> , 2012, 11, 297-303.	2.3	6
137	Evaluation of the Clinical Relevance of Body Composition Parameters in Patients With Cancer Metastatic to the Liver Treated With Hepatic Arterial Infusion Chemotherapy. <i>Nutrition and Cancer</i> , 2012, 64, 206-217.	2.0	29
138	Insulin Growth Factor-Receptor (IGF-1R) Antibody Cixutumumab Combined with the mTOR Inhibitor Temsirolimus in Patients with Refractory Ewing's Sarcoma Family Tumors. <i>Clinical Cancer Research</i> , 2012, 18, 2625-2631.	7.0	184
139	PI3K/AKT/mTOR Inhibitors in Patients With Breast and Gynecologic Malignancies Harboring PI3KCA Mutations. <i>Journal of Clinical Oncology</i> , 2012, 30, 777-782.	1.6	414
140	Safety, pharmacokinetics, and activity of EZN2208, a novel conjugate of polyethylene glycol and SN38, in patients with advanced malignancies. <i>Cancer</i> , 2012, 118, 6144-6151.	4.1	42
141	Perifosine plus docetaxel in patients with platinum and taxane resistant or refractory high-grade epithelial ovarian cancer. <i>Gynecologic Oncology</i> , 2012, 126, 47-53.	1.4	74
142	Validation of the royal marsden hospital prognostic score in patients treated in the phase I clinical trials program at the MD Anderson Cancer Center. <i>Cancer</i> , 2012, 118, 1422-1428.	4.1	88
143	KRASness and PI3KANess in Patients with Advanced Colorectal Cancer: Outcome after Treatment with Early-Phase Trials with Targeted Pathway Inhibitors. <i>PLoS ONE</i> , 2012, 7, e38033.	2.5	44
144	PIK3CA Mutations in Advanced Cancers: Characteristics and Outcomes. <i>Oncotarget</i> , 2012, 3, 1566-1575.	1.8	79

#	ARTICLE	IF	CITATIONS
145	Intraperitoneal and intravenous chemotherapy in peritoneal carcinomatosis. <i>Hepato-Gastroenterology</i> , 2012, 59, 960-4.	0.5	2
146	PIK3CA Mutations Frequently Coexist with RAS and BRAF Mutations in Patients with Advanced Cancers. <i>PLoS ONE</i> , 2011, 6, e22769.	2.5	174
147	PIK3CA Mutations in Patients with Advanced Cancers Treated with PI3K/AKT/mTOR Axis Inhibitors. <i>Molecular Cancer Therapeutics</i> , 2011, 10, 558-565.	4.1	311
148	Phase I clinical trial of hepatic arterial infusion of paclitaxel in patients with advanced cancer and dominant liver involvement. <i>Cancer Chemotherapy and Pharmacology</i> , 2011, 68, 247-253.	2.3	22
149	Phase I study to reverse platinum resistance through use of a hypomethylating agent, azacitidine, in patients with platinum-resistant or platinum-refractory epithelial ovarian cancer. <i>Cancer</i> , 2011, 117, 1661-1669.	4.1	156
150	Prevalence of complementary medicine use in a phase 1 clinical trials program. <i>Cancer</i> , 2011, 117, 5142-5150.	4.1	53
151	Phase I Trial of Hepatic Arterial Infusion of Nanoparticle Albumin-Bound Paclitaxel: Toxicity, Pharmacokinetics, and Activity. <i>Molecular Cancer Therapeutics</i> , 2011, 10, 1300-1307.	4.1	22
152	Outcome Analyses After the First Admission to an Intensive Care Unit in Patients With Advanced Cancer Referred to a Phase I Clinical Trials Program. <i>Journal of Clinical Oncology</i> , 2011, 29, 3547-3552.	1.6	28
153	Outcomes of Research Biopsies in Phase I Clinical Trials: The MD Anderson Cancer Center Experience. <i>Oncologist</i> , 2011, 16, 1292-1298.	3.7	60
154	Phase I clinical trial of hepatic arterial infusion of cisplatin in combination with intravenous liposomal doxorubicin in patients with advanced cancer and dominant liver involvement. <i>Cancer Chemotherapy and Pharmacology</i> , 2010, 66, 1087-1093.	2.3	17
155	A phase 1 study of hepatic arterial infusion of oxaliplatin in combination with systemic 5-fluorouracil, leucovorin, and bevacizumab in patients with advanced solid tumors metastatic to the liver. <i>Cancer</i> , 2010, 116, 4086-4094.	4.1	26
156	Development of curcumin as an epigenetic agent. <i>Cancer</i> , 2010, 116, 4670-4676.	4.1	146
157	Patients with Advanced Head and Neck Cancers Have Similar Progression-Free Survival on Phase I Trials and Their Last Food and Drug Administration-Approved Treatment. <i>Clinical Cancer Research</i> , 2010, 16, 4031-4037.	7.0	15
158	Exploratory Study of Hepatic Arterial Infusion Oxaliplatin With Systemic 5-Fluorouracil/Bevacizumab in Patients With Refractory Solid Tumor and Extensive Liver Metastases. <i>Clinical Colorectal Cancer</i> , 2010, 9, 311-314.	2.3	8
159	Nuclear cyclin B1 is overexpressed in low-malignant-potential ovarian tumors but not in epithelial ovarian cancer. <i>American Journal of Obstetrics and Gynecology</i> , 2009, 201, 367.e1-367.e6.	1.3	8
160	The changing face of Phase 1 cancer clinical trials. <i>Cancer</i> , 2009, 115, 1592-1597.	4.1	19
161	Azacitidine enhances sensitivity of platinum-resistant ovarian cancer cells to carboplatin through induction of apoptosis. <i>American Journal of Obstetrics and Gynecology</i> , 2009, 200, 177.e1-177.e9.	1.3	50
162	Abstract B134: PIK3CA mutations in patients with advanced cancers treated in phase I clinical trials. , 2009, , .		2

#	ARTICLE	IF	CITATIONS
163	Diammine Dicarboxylic Acid Platinum Enhances Cytotoxicity in Platinum-Resistant Ovarian Cancer Cells through Induction of Apoptosis and S-Phase Cell Arrest. <i>Pharmaceutical Research</i> , 2008, 25, 2272-2282.	3.5	6
164	Update on Aurora Kinase Inhibitors in Gynecologic Malignancies. <i>Recent Patents on Anti-Cancer Drug Discovery</i> , 2008, 3, 162-177.	1.6	5
165	Pegylated liposomal doxorubicin treatment in recurrent gynecologic cancer patients with renal dysfunction. <i>Gynecologic Oncology</i> , 2007, 106, 375-380.	1.4	9
166	Proteomics in Gynecologic Malignancies. <i>American Journal of Cancer</i> , 2006, 5, 299-317.	0.4	0
167	Clinical application of oxaliplatin in epithelial ovarian cancer. <i>International Journal of Gynecological Cancer</i> , 2006, 16, 1717-1732.	2.5	26
168	Targeting Aurora kinases in ovarian cancer. <i>Expert Opinion on Therapeutic Targets</i> , 2006, 10, 77-85.	3.4	23
169	Molecular approaches to the diagnosis and treatment of cancer. <i>Stem Cells</i> , 1993, 11, 129-130.	3.2	1
170	Use of Retroviral Markers to Identify Efficacy of Purging and Origin of Relapse Following Autologous Bone Marrow and Peripheral Blood Cell Transplantation in Indolent B Cell Neoplasms (Follicular) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 46 Anderson Cancer Center. <i>Human Gene Therapy</i> , 1993, 4, 821-834.	2.7	9
171	Genetic Therapy of Human Neoplastic Disease. <i>Stem Cells and Development</i> , 1993, 2, 373-375.	1.0	0