

# Verena Jantsch

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3343056/publications.pdf>

Version: 2024-02-01

46  
papers

2,359  
citations

218677

26  
h-index

233421

45  
g-index

54  
all docs

54  
docs citations

54  
times ranked

1956  
citing authors

#	ARTICLE	IF	CITATIONS
1	Release of CHK-2 from PPM-1.D anchorage schedules meiotic entry. <i>Science Advances</i> , 2022, 8, eabl8861.	10.3	5
2	DNA topoisomerase 3 is required for efficient germ cell quality control. <i>Journal of Cell Biology</i> , 2021, 220, .	5.2	8
3	Putting organelles in their place. <i>ELife</i> , 2021, 10, .	6.0	1
4	<i>Caenorhabditis elegans</i> RMI2 functional homolog-2 (RMIF-2) and RMI1 (RMH-1) have both overlapping and distinct meiotic functions within the BTR complex. <i>PLoS Genetics</i> , 2021, 17, e1009663.	3.5	5
5	Poly(ADP-ribose) glycohydrolase coordinates meiotic DNA double-strand break induction and repair independent of its catalytic activity. <i>Nature Communications</i> , 2020, 11, 4869.	12.8	16
6	"The nuclear envelope, a meiotic jack-of-all-trades". <i>Current Opinion in Cell Biology</i> , 2020, 64, 34-42.	5.4	25
7	PLK-1 promotes the merger of the parental genome into a single nucleus by triggering lamina disassembly. <i>ELife</i> , 2020, 9, .	6.0	20
8	Of funding and finches. <i>Genome Biology</i> , 2019, 20, 176.	8.8	0
9	Meiotic chromosome movement: whatâ€™s lamin got to do with it?. <i>Nucleus</i> , 2019, 10, 1-6.	2.2	6
10	Meiotic chromosomes in motion: a perspective from <i>Mus musculus</i> and <i>Caenorhabditis elegans</i> . <i>Chromosoma</i> , 2019, 128, 317-330.	2.2	37
11	LEM-3 is a midbody-tethered DNA nuclease that resolves chromatin bridges during late mitosis. <i>Nature Communications</i> , 2018, 9, 728.	12.8	37
12	Transient and Partial Nuclear Lamina Disruption Promotes Chromosome Movement in Early Meiotic Prophase. <i>Developmental Cell</i> , 2018, 45, 212-225.e7.	7.0	40
13	<i>C. elegans</i> ZHP-4 is required at multiple distinct steps in the formation of crossovers and their transition to segregation competent chiasmata. <i>PLoS Genetics</i> , 2018, 14, e1007776.	3.5	24
14	BRCA1-BARD1 associate with the synaptonemal complex and pro-crossover factors and influence RAD-51 dynamics during <i>Caenorhabditis elegans</i> meiosis. <i>PLoS Genetics</i> , 2018, 14, e1007653.	3.5	44
15	The conserved LEM-3/Ankle1 nuclease is involved in the combinatorial regulation of meiotic recombination repair and chromosome segregation in <i>Caenorhabditis elegans</i> . <i>PLoS Genetics</i> , 2018, 14, e1007453.	3.5	22
16	Initiation of Meiotic Development Is Controlled by Three Post-transcriptional Pathways in <i>Caenorhabditis elegans</i> . <i>Genetics</i> , 2018, 209, 1197-1224.	2.9	38
17	Meiosis. <i>WormBook</i> , 2017, 2017, 1-43.	5.3	92
18	Separable Roles for a <i>Caenorhabditis elegans</i> RMI1 Homolog in Promoting and Antagonizing Meiotic Crossovers Ensure Faithful Chromosome Inheritance. <i>PLoS Biology</i> , 2016, 14, e1002412.	5.6	32

#	ARTICLE	IF	CITATIONS
19	UNC-84: <i>œLINC-ing</i> -chromosome movement and double strand break repair. <i>Journal of Cell Biology</i> , 2016, 215, 753-756.	5.2	3
20	Nuclear Envelope Retention of LINC Complexes Is Promoted by SUN-1 Oligomerization in the <i>Caenorhabditis elegans</i> Germ Line. <i>Genetics</i> , 2016, 203, 733-748.	2.9	8
21	A Surveillance System Ensures Crossover Formation in <i>C. elegans</i> . <i>Current Biology</i> , 2016, 26, 2873-2884.	3.9	56
22	Bisecting Galactose as a Feature of N-Glycans of Wild-type and Mutant <i>Caenorhabditis elegans</i> . <i>Molecular and Cellular Proteomics</i> , 2015, 14, 2111-2125.	3.8	32
23	Chromosome movement in meiosis I prophase of <i>Caenorhabditis elegans</i> . <i>Chromosoma</i> , 2014, 123, 15-24.	2.2	48
24	Matefin/SUN-1 Phosphorylation Is Part of a Surveillance Mechanism to Coordinate Chromosome Synapsis and Recombination with Meiotic Progression and Chromosome Movement. <i>PLoS Genetics</i> , 2013, 9, e1003335.	3.5	90
25	Combinatorial Regulation of Meiotic Holliday Junction Resolution in <i>C. elegans</i> by HIM-6 (BLM) Helicase, SLX-4, and the SLX-1, MUS-81 and XPF-1 Nucleases. <i>PLoS Genetics</i> , 2013, 9, e1003591.	3.5	88
26	Galactosylated Fucose Epitopes in Nematodes. <i>Journal of Biological Chemistry</i> , 2012, 287, 28276-28290.	3.4	43
27	Transgene-mediated cosuppression and RNA interference enhance germ-line apoptosis in <i>Caenorhabditis elegans</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 3440-3445.	7.1	17
28	Photo-sensitive hydrogels for three-dimensional laser microfabrication in the presence of whole organisms. <i>Journal of Biomedical Optics</i> , 2012, 17, 1.	2.6	117
29	Polo Kinases Establish Links between Meiotic Chromosomes and Cytoskeletal Forces Essential for Homolog Pairing. <i>Developmental Cell</i> , 2011, 21, 948-958.	7.0	104
30	A New Thermosensitive smc-3 Allele Reveals Involvement of Cohesin in Homologous Recombination in <i>C. elegans</i> . <i>PLoS ONE</i> , 2011, 6, e24799.	2.5	17
31	Mutations in <i>Caenorhabditis elegans him-19</i> Show Meiotic Defects That Worsen with Age. <i>Molecular Biology of the Cell</i> , 2010, 21, 885-896.	2.1	24
32	Leptotene/Zygotene Chromosome Movement Via the SUN/KASH Protein Bridge in <i>Caenorhabditis elegans</i> . <i>PLoS Genetics</i> , 2010, 6, e1001219.	3.5	72
33	<i>Caenorhabditis elegans</i> RNA-Directed RNA Polymerase in Sperm Development and Endogenous RNA Interference. <i>Genetics</i> , 2009, 183, 1297-1314.	2.9	80
34	SUN-domain and KASH-domain proteins during development, meiosis and disease. <i>Cellular and Molecular Life Sciences</i> , 2009, 66, 1518-1533.	5.4	87
35	Meiotic Chromosome Homology Search Involves Modifications of the Nuclear Envelope Protein Matefin/SUN-1. <i>Cell</i> , 2009, 139, 920-933.	28.9	181
36	ZHP-3 Acts at Crossovers to Couple Meiotic Recombination with Synaptonemal Complex Disassembly and Bivalent Formation in <i>C. elegans</i> . <i>PLoS Genetics</i> , 2008, 4, e1000235.	3.5	129

#	ARTICLE	IF	CITATIONS
37	Biosynthesis of Truncated N-Linked Oligosaccharides Results from Non-orthologous Hexosaminidase-mediated Mechanisms in Nematodes, Plants, and Insects. <i>Journal of Biological Chemistry</i> , 2007, 282, 27825-27840.	3.4	84
38	<i>Caenorhabditis elegans</i> <i>prom-1</i> Is Required for Meiotic Prophase Progression and Homologous Chromosome Pairing. <i>Molecular Biology of the Cell</i> , 2007, 18, 4911-4920.	2.1	34
39	The Nuclear Envelope Protein Matefin/SUN-1 Is Required for Homologous Pairing in <i>C. elegans</i> Meiosis. <i>Developmental Cell</i> , 2007, 12, 873-885.	7.0	166
40	A conserved function for a <i>Caenorhabditis elegans</i> Com1/Sae2/CtIP protein homolog in meiotic recombination. <i>EMBO Journal</i> , 2007, 26, 5071-5082.	7.8	94
41	Reconstitution in vitro of the GDP-fucose biosynthetic pathways of <i>Caenorhabditis elegans</i> and <i>Drosophila melanogaster</i> . <i>FEBS Journal</i> , 2006, 273, 2244-2256.	4.7	22
42	A Deletion in the Golgi $\alpha$ -Mannosidase II Gene of <i>Caenorhabditis elegans</i> Results in Unexpected Non-wild-type N-Glycan Structures. <i>Journal of Biological Chemistry</i> , 2006, 281, 28265-28277.	3.4	44
43	Molecular Basis of Anti-horseradish Peroxidase Staining in <i>Caenorhabditis elegans</i> . <i>Journal of Biological Chemistry</i> , 2004, 279, 49588-49598.	3.4	74
44	Targeted Gene Knockout Reveals a Role in Meiotic Recombination for ZHP-3, a Zip3-Related Protein in <i>Caenorhabditis elegans</i> . <i>Molecular and Cellular Biology</i> , 2004, 24, 7998-8006.	2.3	110
45	The <i>Caenorhabditis elegans</i> SCC-3 homologue is required for meiotic synapsis and for proper chromosome disjunction in mitosis and meiosis. <i>Experimental Cell Research</i> , 2003, 289, 245-255.	2.6	46
46	The Tpv2 family of retrotransposons of <i>Phaseolus vulgaris</i> : structure, integration characteristics, and use for genotype classification. <i>Plant Molecular Biology</i> , 1999, 39, 797-807.	3.9	28