

# Alicia Del Toro-Arreola

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3350793/publications.pdf>

Version: 2024-02-01

10  
papers

440  
citations

1040056

9  
h-index

1372567

10  
g-index

10  
all docs

10  
docs citations

10  
times ranked

950  
citing authors

#	ARTICLE	IF	CITATIONS
1	Low NKp30, NKp46 and NKG2D expression and reduced cytotoxic activity on NK cells in cervical cancer and precursor lesions. <i>BMC Cancer</i> , 2009, 9, 186.	2.6	164
2	Effect of d-limonene on immune response in BALB/c mice with lymphoma. <i>International Immunopharmacology</i> , 2005, 5, 829-838.	3.8	78
3	Augmented serum level of major histocompatibility complex class I-related chain A (MICA) protein and reduced NKG2D expression on NK and T cells in patients with cervical cancer and precursor lesions. <i>BMC Cancer</i> , 2008, 8, 16.	2.6	70
4	Protease-activated receptor-2 (PAR-2) in cervical cancer proliferation. <i>Gynecologic Oncology</i> , 2008, 108, 19-26.	1.4	26
5	The role of endothelial cells on islet function and revascularization after islet transplantation. <i>Organogenesis</i> , 2016, 12, 28-32.	1.2	22
6	Circulating soluble levels of MIF in women with breast cancer in the molecular subtypes: relationship with Th17 cytokine profile. <i>Clinical and Experimental Medicine</i> , 2019, 19, 385-391.	3.6	22
7	Preparation of PLGA/Rose Bengal colloidal particles by double emulsion and layer-by-layer for breast cancer treatment. <i>Journal of Colloid and Interface Science</i> , 2018, 518, 122-129.	9.4	20
8	Characterization of B7H6, an endogenous ligand for the NK cell activating receptor NKp30, reveals the identity of two different soluble isoforms during normal human pregnancy. <i>Immunobiology</i> , 2018, 223, 57-63.	1.9	16
9	MHC class I-related chain A and B ligands are differentially expressed in human cervical cancer cell lines. <i>Cancer Cell International</i> , 2011, 11, 15.	4.1	15
10	Association of the genetic variants (â€794 CATT5â€8 and â€173 GÂ>ÂC) of macrophage migration inhibitory factor (MIF) with higher soluble levels of MIF and TNFÎ± in women with breast cancer. <i>Journal of Clinical Laboratory Analysis</i> , 2020, 34, e23209.	2.1	7