

# Yota Murakami

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3364340/publications.pdf>

Version: 2024-02-01

41  
papers

1,151  
citations

471509

17  
h-index

414414

32  
g-index

43  
all docs

43  
docs citations

43  
times ranked

1372  
citing authors

#	ARTICLE	IF	CITATIONS
1	Histone variant H2A.Z plays multiple roles in the maintenance of heterochromatin integrity. <i>Genes To Cells</i> , 2022, 27, 93-112.	1.2	4
2	Construction and characterization of a zinc-inducible gene expression vector in fission yeast. <i>Yeast</i> , 2021, 38, 251-261.	1.7	0
3	Trimethylguanosine synthase 1 (Tgs1) is involved in Swi6/HP1-independent siRNA production and establishment of heterochromatin in fission yeast. <i>Genes To Cells</i> , 2021, 26, 203-218.	1.2	3
4	Two secured FACT recruitment mechanisms are essential for heterochromatin maintenance. <i>Cell Reports</i> , 2021, 36, 109540.	6.4	13
5	Nonmuscle myosin IIA and IIB differently suppress microtubule growth to stabilize cell morphology. <i>Journal of Biochemistry</i> , 2020, 167, 25-39.	1.7	4
6	Unprogrammed epigenetic variation mediated by stochastic formation of ectopic heterochromatin. <i>Current Genetics</i> , 2020, 66, 319-325.	1.7	2
7	Epigenetic regulation affects gene amplification in <i>Drosophila</i> development. <i>Frontiers in Bioscience - Landmark</i> , 2020, 25, 632-645.	3.0	4
8	Phosphorylation of repressive histone code readers by casein kinase 2 plays diverse roles in heterochromatin regulation. <i>Journal of Biochemistry</i> , 2019, 166, 3-6.	1.7	7
9	Regulation of ectopic heterochromatin-mediated epigenetic diversification by the JmjC family protein Epe1. <i>PLoS Genetics</i> , 2019, 15, e1008129.	3.5	23
10	Differential contributions of nonmuscle myosin IIA and IIB to cytokinesis in human immortalized fibroblasts. <i>Experimental Cell Research</i> , 2019, 376, 67-76.	2.6	19
11	Complete Genome Sequence of <i>Staphylococcus arlettae</i> Strain P2, Isolated from a Laboratory Environment. <i>Microbiology Resource Announcements</i> , 2019, 8, .	0.6	5
12	Heterochromatin suppresses gross chromosomal rearrangements at centromeres by repressing Tfs1/TFIIS-dependent transcription. <i>Communications Biology</i> , 2019, 2, 17.	4.4	24
13	Nonmuscle myosin IIA and IIB differentially contribute to intrinsic and directed migration of human embryonic lung fibroblasts. <i>Biochemical and Biophysical Research Communications</i> , 2018, 498, 25-31.	2.1	7
14	Different contributions of nonmuscle myosin IIA and IIB to the organization of stress fiber subtypes in fibroblasts. <i>Molecular Biology of the Cell</i> , 2018, 29, 911-922.	2.1	26
15	DNA replication machinery is required for development in <i>Drosophila</i> . <i>Frontiers in Bioscience - Landmark</i> , 2018, 23, 493-505.	3.0	5
16	RNAi-dependent heterochromatin assembly in fission yeast <i>Schizosaccharomyces pombe</i> requires heat-shock molecular chaperones Hsp90 and Mas5. <i>Epigenetics and Chromatin</i> , 2018, 11, 26.	3.9	11
17	Sfh1, an essential component of the RSC chromatin remodeling complex, maintains genome integrity in fission yeast. <i>Genes To Cells</i> , 2018, 23, 738-752.	1.2	5
18	DNA replication machinery contributes to development of eye in <i>Drosophila</i> . <i>Frontiers in Bioscience - Landmark</i> , 2018, 23, 506-511.	3.0	2

#	ARTICLE	IF	CITATIONS
19	Semi-retentive cytoskeletal fractionation (SERCYF): A novel method for the biochemical analysis of the organization of microtubule and actin cytoskeleton networks. <i>Biochemical and Biophysical Research Communications</i> , 2017, 488, 614-620.	2.1	4
20	Ser7 of RNAPII-CTD facilitates heterochromatin formation by linking ncRNA to RNAi. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E11208-E11217.	7.1	13
21	H3K36 methylation state and associated silencing mechanisms. <i>Transcription</i> , 2017, 8, 26-31.	3.1	12
22	Inner nuclear membrane protein Lem2 augments heterochromatin formation in response to nutritional conditions. <i>Genes To Cells</i> , 2016, 21, 812-832.	1.2	44
23	Histone H3K36 trimethylation is essential for multiple silencing mechanisms in fission yeast. <i>Nucleic Acids Research</i> , 2016, 44, 4147-4162.	14.5	44
24	A novel method for purification of the endogenously expressed fission yeast Set2 complex. <i>Protein Expression and Purification</i> , 2014, 97, 44-49.	1.3	2
25	Histone deacetylases govern heterochromatin in every phase. <i>EMBO Journal</i> , 2013, 32, 2301-2303.	7.8	12
26	Mediator Directs Co-transcriptional Heterochromatin Assembly by RNA Interference-Dependent and -Independent Pathways. <i>PLoS Genetics</i> , 2013, 9, e1003677.	3.5	28
27	Spt6 prevents transcription-coupled loss of posttranslationally modified histone H3. <i>Scientific Reports</i> , 2013, 3, 2186.	3.3	52
28	Heterochromatin protein 1 homologue Swi6 acts in concert with Ers1 to regulate RNAi-directed heterochromatin assembly. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2012, 109, 6159-6164.	7.1	30
29	A novel RNAi protein, Dsh1, assembles RNAi machinery on chromatin to amplify heterochromatic siRNA. <i>Genes and Development</i> , 2012, 26, 1811-1824.	5.9	22
30	Ran and Calcineurin Can Participate Collaboratively in the Regulation of Spermatogenesis in Scallop. <i>Marine Biotechnology</i> , 2012, 14, 479-490.	2.4	4
31	DNA-RNA hybrid formation mediates RNAi-directed heterochromatin formation. <i>Genes To Cells</i> , 2012, 17, 218-233.	1.2	94
32	Roles of the C-terminal residues of calmodulin in structure and function. <i>Biophysics (Nagoya-shi)</i> , 2012, 10, 10-15.	0.4	5
33	Phosphorylation of Swi6/HP1 regulates transcriptional gene silencing at heterochromatin. <i>Genes and Development</i> , 2009, 23, 18-23.	5.9	61
34	Fission yeast chromatin assembly factor 1 assists in the replication-coupled maintenance of heterochromatin. <i>Genes To Cells</i> , 2008, 13, 1027-1043.	1.2	41
35	Transcription factor Runx1 recruits the polyomavirus replication origin to replication factories. <i>Journal of Cellular Biochemistry</i> , 2007, 100, 1313-1323.	2.6	13
36	Transcription factors and DNA replication origin selection. <i>BioEssays</i> , 2005, 27, 1107-1116.	2.5	56

#	ARTICLE	IF	CITATIONS
37	RNA Polymerase II Is Required for RNAi-Dependent Heterochromatin Assembly. <i>Science</i> , 2005, 309, 467-469.	12.6	258
38	Fission yeast CENP-B homologs nucleate centromeric heterochromatin by promoting heterochromatin-specific histone tail modifications. <i>Genes and Development</i> , 2002, 16, 1766-1778.	5.9	97
39	Block of granulocytic differentiation of 32Dcl3 cells by AML1/ETO(MTG8) but not by highly expressed Bcl-2. <i>Oncogene</i> , 1999, 18, 4055-4062.	5.9	36
40	Context-Dependent Modulation of Replication Activity of <i>Saccharomyces cerevisiae</i> Autonomously Replicating Sequences by Transcription Factors. <i>Molecular and Cellular Biology</i> , 1999, 19, 7428-7435.	2.3	18
41	The Capacity of Polyomavirus Enhancer Binding Protein 2±B (AML1/Cbfa2) To Stimulate Polyomavirus DNA Replication Is Related to Its Affinity for the Nuclear Matrix. <i>Molecular and Cellular Biology</i> , 1998, 18, 4165-4176.	2.3	40