Su Jin Lee

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3399492/publications.pdf

Version: 2024-02-01

203 5,101 33 61 papers citations h-index g-index

203 203 203 9480 all docs docs citations times ranked citing authors

| # | Article | IF | CITATIONS |
|----|--|------|-----------|
| 1 | Postnatal regulation of B-1a cell development and survival by the CIC-PER2-BHLHE41 axis. Cell Reports, 2022, 38, 110386. | 6.4 | 7 |
| 2 | IOX1 activity as sepsis therapy and an antibiotic against multidrug-resistant bacteria. Scientific Reports, 2021, 11, 2942. | 3.3 | 4 |
| 3 | Myostatin inhibitor YK11 as a preventative health supplement for bacterial sepsis. Biochemical and Biophysical Research Communications, 2021, 543, 1-7. | 2.1 | 10 |
| 4 | Comparison of bone single-photon emission computed tomography (SPECT)/CT and bone scintigraphy in assessing knee joints. BMC Medical Imaging, 2021, 21, 60. | 2.7 | 5 |
| 5 | Early-phase 18F-FP-CIT and 18F-flutemetamol PET were significantly correlated. Scientific Reports, 2021, 11, 12297. | 3.3 | 5 |
| 6 | Skeletal muscle mass at C3 may not be a strong predictor for skeletal muscle mass at L3 in sarcopenic patients with head and neck cancer. PLoS ONE, 2021, 16, e0254844. | 2.5 | 12 |
| 7 | IOX1 impedes host inflammation in imiquimod-triggered psoriasis. Heliyon, 2021, 7, e08433. | 3.2 | O |
| 8 | Primary Subcapsular Reflux as an Etiology of Subcapsular Renal Abscess. Childhood Kidney Diseases, 2021, 25, 133-139. | 0.4 | 0 |
| 9 | Homeostasis and regulation of autoreactive B cells. Cellular and Molecular Immunology, 2020, 17, 561-569. | 10.5 | 18 |
| 10 | Cell cycle synchronisation using thiazolidinediones affects cellular glucose metabolism and enhances the therapeutic effect of 2-deoxyglucose in colon cancer. Scientific Reports, 2020, 10, 4713. | 3.3 | 4 |
| 11 | Detection of Fusion Genes Using a Targeted RNA Sequencing Panel in Gastrointestinal and Rare Cancers. Journal of Oncology, 2020, 2020, 1-8. | 1.3 | 7 |
| 12 | Synergistic Effects of Combination Therapy with AKT and mTOR Inhibitors on Bladder Cancer Cells. International Journal of Molecular Sciences, 2020, 21, 2825. | 4.1 | 11 |
| 13 | The use of regorafenib for patients with refractory metastatic colorectal cancer in clinical practice. OncoTargets and Therapy, 2019, Volume 12, 225-231. | 2.0 | 4 |
| 14 | Comprehensive molecular and clinical characterization of Asian melanoma patients treated with anti-PD-1 antibody. BMC Cancer, 2019, 19, 805. | 2.6 | 9 |
| 15 | Gastrointestinal stromal tumour with CDKN2A deletions: a report of three cases. Pathology, 2019, 51, 537-539. | 0.6 | O |
| 16 | The impact of primary tumor site on outcomes of treatment with etoposide and cisplatin in grade 3 gastroenteropancreatic neuroendocrine carcinoma. Journal of Cancer, 2019, 10, 3140-3144. | 2.5 | 5 |
| 17 | Clinical significance of radiotherapy before and/or during nivolumab treatment in hepatocellular carcinoma. Cancer Medicine, 2019, 8, 6986-6994. | 2.8 | 37 |
| 18 | Combination of Docetaxel Plus Savolitinib in Refractory Cancer Patients: A Report on Phase I Trial. Translational Oncology, 2019, 12, 597-601. | 3.7 | 8 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 19 | A retrospective feasibility study of biweekly docetaxel in patients with high-risk metastatic castration-na \tilde{A} -ve prostate cancer. BMC Urology, 2019, 19, 30. | 1.4 | 5 |
| 20 | The Impact of Primary Tumor Sidedness on the Effect of Regorafenib in Refractory Metastatic Colorectal Cancer. Journal of Cancer, 2019, 10, 1611-1615. | 2.5 | 7 |
| 21 | PD-L1 protein expression in non-small-cell lung cancer and its relationship with the hypoxia-related signaling pathways: A study based on immunohistochemistry and RNA sequencing data. Lung Cancer, 2019, 129, 41-47. | 2.0 | 24 |
| 22 | Can the basal serum thyroglobulin level be used to predict the recombinant human TSH-stimulated thyroglobulin level in differentiated patients with thyroid cancer?. Medicine (United States), 2019, 98, e18437. | 1.0 | 0 |
| 23 | Anaplastic Oligodendroglioma Found as an Incidental Photopenic Lesion on 18F-FP-CIT PET/CT Image. Clinical Nuclear Medicine, 2019, 44, e168-e169. | 1.3 | 1 |
| 24 | Risk Factors for Indeterminate Response After Radioactive Iodine Therapy in Patients With Differentiated Thyroid Cancer. Clinical Nuclear Medicine, 2019, 44, 714-718. | 1.3 | 7 |
| 25 | Intratumoral heterogeneity as measured using the tumor-stroma ratio and PET texture analyses in females with lung adenocarcinomas differs from that of males with lung adenocarcinomas or squamous cell carcinomas. Medicine (United States), 2019, 98, e14876. | 1.0 | 6 |
| 26 | LAG3 in Solid Tumors as a Potential Novel Immunotherapy Target. Journal of Immunotherapy, 2019, 42, 279-283. | 2.4 | 11 |
| 27 | A Case of Radioactive Iodine Uptake Found in Artificial Eye. Clinical Nuclear Medicine, 2019, 44, 317-318. | 1.3 | 2 |
| 28 | A Machine-Learning Approach Using PET-Based Radiomics to Predict the Histological Subtypes of Lung Cancer. Clinical Nuclear Medicine, 2019, 44, 956-960. | 1.3 | 101 |
| 29 | Clinicopathologic parameters associated with the FDG-avidity in staging of early gastric cancer using 18F-FDG PET. Medicine (United States), 2019, 98, e16690. | 1.0 | 4 |
| 30 | Multidisciplinary approach is associated with improved survival of hepatocellular carcinoma patients. PLoS ONE, 2019, 14, e0210730. | 2.5 | 64 |
| 31 | 18F-fluorodeoxyglucose positron emission tomography is correlated with the pathological necrosis and decreased microvessel density in lung adenocarcinomas. Annals of Nuclear Medicine, 2019, 33, 93-102. | 2.2 | 7 |
| 32 | Multicenter retrospective analysis of the safety and efficacy of regorafenib after progression on sorafenib in Korean patients with hepatocellular carcinoma. Investigational New Drugs, 2019, 37, 567-572. | 2.6 | 44 |
| 33 | First-in-human phase I trial of anti-hepatocyte growth factor (HGF) antibody (YYB101) in refractory solid tumor patients: Integrative pathologic-genomic analysis and the final results Journal of Clinical Oncology, 2019, 37, 3104-3104. | 1.6 | 2 |
| 34 | A Randomized, Open-Label, Phase II Study Comparing Pemetrexed Plus Cisplatin Followed by Maintenance Pemetrexed versus Pemetrexed Alone in Patients with Epidermal Growth Factor Receptor (EGFR)-Mutant Non-small Cell Lung Cancer after Failure of First-Line EGFR Tyrosine Kinase Inhibitor: KCSG-LU12-13. Cancer Research and Treatment, 2019, 51, 718-726. | 3.0 | 10 |
| 35 | Comparison of the 7th and the 8th AJCC Staging System for Non-metastatic D2-Resected Lymph Node–Positive Gastric Cancer Treated with Different Adjuvant Protocols. Cancer Research and Treatment, 2019, 51, 876-885. | 3.0 | 8 |
| 36 | The impact of primary tumor site on outcomes of treatment with etoposide and cisplatin in grade 3 gastroenteropancreatic neuroendocrine carcinoma Journal of Clinical Oncology, 2019, 37, 213-213. | 1.6 | 0 |

| # | Article | IF | Citations |
|----|--|------|-----------|
| 37 | Phase 1a study results investigating the safety and preliminary efficacy of ABL001 (NOV1501), a bispecific antibody targeting VEGF and DLL4 in metastatic gastrointestinal (GI) cancer Journal of Clinical Oncology, 2019, 37, 3023-3023. | 1.6 | 3 |
| 38 | Atezolizumab in Patients with Pretreated Urothelial Cancer: a Korean Single-Center, Retrospective Study. Cancer Research and Treatment, 2019, 51, 1269-1274. | 3.0 | 4 |
| 39 | Enhanced Expression and Functional Characterization of the Recombinant Putative Lysozyme-PMAP36 Fusion Protein. Molecules and Cells, 2019, 42, 262-269. | 2.6 | 3 |
| 40 | MCT4 Expression Is a Potential Therapeutic Target in Colorectal Cancer with Peritoneal Carcinomatosis. Molecular Cancer Therapeutics, 2018, 17, 838-848. | 4.1 | 36 |
| 41 | Phase I Trial of Anti-MET Monoclonal Antibody in MET-Overexpressed Refractory Cancer. Clinical Colorectal Cancer, 2018, 17, 140-146. | 2.3 | 17 |
| 42 | c-MET Overexpression in Colorectal Cancer: A Poor Prognostic Factor for Survival. Clinical Colorectal Cancer, 2018, 17, 165-169. | 2.3 | 71 |
| 43 | The Correlation Between Serum Chemokines and Clinical Outcome in Patients with Advanced Biliary Tract Cancer. Translational Oncology, 2018, 11, 353-357. | 3.7 | 8 |
| 44 | Clinical benefit and residual kidney function of en bloc nephrectomy for perirenal retroperitoneal sarcoma. Asia-Pacific Journal of Clinical Oncology, 2018, 14, e465-e471. | 1.1 | 4 |
| 45 | Adjuvant Chemotherapy with or without Concurrent Radiotherapy for Patients with Stage IB Gastric Cancer: a Subgroup Analysis of the Adjuvant Chemoradiotherapy in Stomach Tumors (ARTIST) Phase III Trial. Journal of Gastric Cancer, 2018, 18, 348. | 2.5 | 12 |
| 46 | Fibroblast growth factor receptor 3 (FGFR3) aberrations in muscle-invasive urothelial carcinoma. BMC Urology, 2018, 18, 68. | 1.4 | 18 |
| 47 | Necessity of adjuvant concurrent chemo-radiotherapy in D2-resected LN-positive gastric cancer. Radiotherapy and Oncology, 2018, 129, 306-312. | 0.6 | 12 |
| 48 | Pazopanib monotherapy in the treatment of pretreated, metastatic uterine sarcoma: a single-center retrospective study. Journal of Gynecologic Oncology, 2018, 29, e3. | 2.2 | 13 |
| 49 | Comprehensive molecular characterization of clinical responses to PD-1 inhibition in metastatic gastric cancer. Nature Medicine, 2018, 24, 1449-1458. | 30.7 | 1,071 |
| 50 | Pemetrexed Monotherapy as Salvage Treatment in Patients with Metastatic Colorectal Cancer Refractory to Standard Chemotherapy: A Phase II Single-arm Prospective Trial. Journal of Cancer, 2018, 9, 2910-2915. | 2.5 | 6 |
| 51 | First-in-human phase I trial of anti-hepatocyte growth factor (HGF) antibody (YYB101) in refractory solid tumor patients Journal of Clinical Oncology, 2018, 36, e14501-e14501. | 1.6 | 1 |
| 52 | Associations Between PET Textural Features and GLUT1 Expression, and the Prognostic Significance of Textural Features in Lung Adenocarcinoma. Anticancer Research, 2018, 38, 1067-1071. | 1.1 | 11 |
| 53 | Pazopanib for the Treatment of Non-clear Cell Renal Cell Carcinoma: A Single-Arm, Open-Label, Multicenter, Phase II Study. Cancer Research and Treatment, 2018, 50, 488-494. | 3.0 | 28 |
| 54 | VariantPlex panel to detect genomic aberrations in oncology patients with rare cancer type Journal of Clinical Oncology, 2018, 36, e24234-e24234. | 1.6 | 0 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Detection of targetable fusions using FusionPlex in oncology patients Journal of Clinical Oncology, 2018, 36, e24238-e24238. | 1.6 | 0 |
| 56 | Intrinsic resistance to sunitinib in patients with metastatic renal cell carcinoma. Asia-Pacific Journal of Clinical Oncology, 2017, 13, 61-67. | 1.1 | 18 |
| 57 | Clinical trial of nintedanib in patients with recurrent or metastatic salivary gland cancer of the head and neck: A multicenter phase 2 study (Korean Cancer Study Group HN14â€01). Cancer, 2017, 123, 1958-1964. | 4.1 | 44 |
| 58 | Differential expression and prognostic significance of GLUT1 according to histologic type of non-small-cell lung cancer and its association with volume-dependent parameters. Lung Cancer, 2017, 104, 31-37. | 2.0 | 54 |
| 59 | A multi-center, open-label, randomized phase III trial of first-line chemotherapy with capecitabine monotherapy versus capecitabine plus oxaliplatin in elderly patients with advanced gastric cancer. Journal of Geriatric Oncology, 2017, 8, 170-175. | 1.0 | 39 |
| 60 | Different subtypes of epithelioid sarcoma and their clinical implication: longâ€term multiâ€institutional experience with a rare sarcoma. Apmis, 2017, 125, 223-229. | 2.0 | 15 |
| 61 | Phase I trial and pharmacokinetic study of tanibirumab, a fully human monoclonal antibody to vascular endothelial growth factor receptor 2, in patients with refractory solid tumors. Investigational New Drugs, 2017, 35, 782-790. | 2.6 | 22 |
| 62 | Hepatocellular carcinoma treatment: a comparative review of emerging growth factor receptor antagonists. Expert Opinion on Emerging Drugs, 2017, 22, 191-200. | 2.4 | 3 |
| 63 | Neoagarooligosaccharides prevent septic shock by modulating A20-and cyclooxygenase-2-mediated interleukin-10 secretion in a septic-shock mouse model. Biochemical and Biophysical Research Communications, 2017, 486, 998-1004. | 2.1 | 19 |
| 64 | Intratumoural heterogeneity measured using FDG PET and MRI is associated with tumour–stroma ratio and clinical outcome in head and neck squamous cell carcinoma. Clinical Radiology, 2017, 72, 482-489. | 1.1 | 25 |
| 65 | A retrospective feasibility study of biweekly, reduced-dose docetaxel in Asian patients with castrate-resistant, metastatic prostate cancer. BMC Urology, 2017, 17, 63. | 1.4 | 7 |
| 66 | Radiolabeled Anti-Adenosine Triphosphate Synthase Monoclonal Antibody as a Theragnostic Agent Targeting Angiogenesis. Molecular Imaging, 2017, 16, 153601211773739. | 1.4 | 6 |
| 67 | Prognostic significance of lactate dehydrogenase B according to histologic type of non-small-cell lung cancer and its association with serum lactate dehydrogenase. Pathology Research and Practice, 2017, 213, 1134-1138. | 2.3 | 20 |
| 68 | Clinical usefulness of F-18 FDG PET in lymphoepithelioma-like gastric carcinoma. European Journal of Radiology, 2017, 94, 160-166. | 2.6 | 7 |
| 69 | Prognostic value of preoperative total psoas muscle area on long-term outcome in surgically treated oesophageal cancer patients. Interactive Cardiovascular and Thoracic Surgery, 2017, 24, 13-19. | 1.1 | 20 |
| 70 | Prognostic utility of FDG PET/CT and bone scintigraphy in breast cancer patients with bone-only metastasis. Medicine (United States), 2017, 96, e8985. | 1.0 | 11 |
| 71 | An investigation of the role of gene copy number variations in sorafenib sensitivity in metastatic hepatocellular carcinoma patients. Journal of Cancer, 2017, 8, 730-736. | 2.5 | 1 |
| 72 | Prognostic Significance of EDIL3 Expression and Correlation with Mesenchymal Phenotype and Microvessel Density in Lung Adenocarcinoma. Scientific Reports, 2017, 7, 8649. | 3.3 | 28 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Serosal Cavities Contain Two Populations of Innate-like integrin $\hat{l}\pm4$ highCD4+T Cells, Integrin $\hat{l}\pm4\hat{l}^21+\hat{l}\pm4\hat{l}^27\hat{a}^2$ and $\hat{l}\pm4\hat{l}^21+\hat{l}\pm6\hat{l}^21\hat{a}^2\hat{l}\pm4\hat{l}^27+$ Cells. Immune Network, 2017, 17, 392. | 3.6 | 4 |
| 74 | Direct analysis of aberrant glycosylation on haptoglobin in patients with gastric cancer. Oncotarget, 2017, 8, 11094-11104. | 1.8 | 21 |
| 75 | Postoperative change of the psoas muscle area as a predictor of survival in surgically treated esophageal cancer patients. Journal of Thoracic Disease, 2017, 9, 355-361. | 1.4 | 17 |
| 76 | The implication of FLT3 amplification for FLT targeted therapeutics in solid tumors. Oncotarget, 2017, 8, 3237-3245. | 1.8 | 20 |
| 77 | MerTK is a novel therapeutic target in gastric cancer. Oncotarget, 2017, 8, 96656-96667. | 1.8 | 23 |
| 78 | Efficacy and Safety of Regorafenib in Korean Patients with Advanced Gastrointestinal Stromal Tumor after Failure of Imatinib and Sunitinib: A Multicenter Study Based on the Management Access Program. Cancer Research and Treatment, 2017, 49, 350-357. | 3.0 | 23 |
| 79 | Programmed death (PD)-ligand 1 (L1) expression and mismatch repair (MMR) deficiency across tumor types: Candidates for checkpoint inhibitor based immunotherapy Journal of Clinical Oncology, 2017, 35, e14622-e14622. | 1.6 | 0 |
| 80 | The impact of pathologic differentiation (well/ poorly) and the degree of Ki-67 index in patients with metastatic WHO grade 3 GEP-NECs Journal of Clinical Oncology, 2017, 35, e15686-e15686. | 1.6 | 0 |
| 81 | Genomic Profiling of Metastatic Gastroenteropancreatic Neuroendocrine Tumor (GEP-NET) Patients in the Personalized-Medicine Era. Journal of Cancer, 2016, 7, 1044-1048. | 2.5 | 17 |
| 82 | The Influence of Metastatic Lymph Node Ratio on the Treatment Outcomes in the Adjuvant Chemoradiotherapy in Stomach Tumors (ARTIST) Trial: A Phase III Trial. Journal of Gastric Cancer, 2016, 16, 105. | 2.5 | 34 |
| 83 | The Impact of PD-L1 Expression in Patients with Metastatic GEP-NETs. Journal of Cancer, 2016, 7, 484-489. | 2.5 | 106 |
| 84 | Efficacy of BRAF Inhibitors in Asian Metastatic Melanoma Patients: Potential Implications of Genomic Sequencing in BRAF-Mutated Melanoma. Translational Oncology, 2016, 9, 557-564. | 3.7 | 16 |
| 85 | Clinical significance of post-treatment 18F-fluorodeoxyglucose uptake in cervical lymph nodes in patients with diffuse large B-cell lymphoma. European Radiology, 2016, 26, 4632-4639. | 4.5 | 14 |
| 86 | BEZ235 (PIK3/mTOR inhibitor) Overcomes Pazopanib Resistance in Patient-Derived Refractory Soft Tissue Sarcoma Cells. Translational Oncology, 2016, 9, 197-202. | 3.7 | 10 |
| 87 | The Highest Metabolic Activity on FDG PET Is Associated With Overall Survival in Limited-Stage Small-Cell Lung Cancer. Medicine (United States), 2016, 95, e2772. | 1.0 | 18 |
| 88 | The prognostic value of total lesion glycolysis via 18F-fluorodeoxyglucose PET–CT in surgically treated esophageal squamous cell carcinoma. Annals of Nuclear Medicine, 2016, 30, 81-88. | 2.2 | 15 |
| 89 | A Korean multi-center, real-world, retrospective study of first-line pazopanib in unselected patients with metastatic renal clear-cell carcinoma. BMC Urology, 2016, 16, 46. | 1.4 | 14 |
| 90 | Loss of Tuberous Sclerosis Complex 2 (TSC2) as a Predictive Biomarker of Response to mTOR Inhibitor Treatment in Patients with Hepatocellular Carcinoma. Translational Oncology, 2016, 9, 466-471. | 3.7 | 13 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | A Korean single-center, real-world, retrospective study of first-line weekly paclitaxel in patients with metastatic angiosarcoma. Clinical Sarcoma Research, 2016, 6, 8. | 2.3 | 10 |
| 92 | Different clinical outcomes between locally advanced hypopharyngeal and oropharyngeal cancer treated with definitive concurrent chemoradiotherapy: implication for subgroup selection for induction chemotherapy. Japanese Journal of Clinical Oncology, 2016, 46, 40-45. | 1.3 | 9 |
| 93 | Value of FGFR2 expression for advanced gastric cancer patients receiving pazopanib plus CapeOX (capecitabine and oxaliplatin). Journal of Cancer Research and Clinical Oncology, 2016, 142, 1231-1237. | 2.5 | 11 |
| 94 | Comparison of Diagnostic Performance Between Visual and Quantitative Assessment of Bone Scintigraphy Results in Patients With Painful Temporomandibular Disorder. Medicine (United States), 2016, 95, e2485. | 1.0 | 5 |
| 95 | Troglitazone Stimulates Cancer Cell Uptake of $\langle \sup 18 \langle \sup FFDG \rangle$ by Suppressing Mitochondrial Respiration and Augments Sensitivity to Glucose Restriction. Journal of Nuclear Medicine, 2016, 57, 129-135. | 5.0 | 5 |
| 96 | Phase II trial of nintedanib in patients with recurrent or metastatic salivary gland cancer: A multicenter phase II study Journal of Clinical Oncology, 2016, 34, 6090-6090. | 1.6 | 3 |
| 97 | Prospective phase II trial of pazopanib plus CapeOX (capecitabine and oxaliplatin) in previously untreated patients with advanced gastric cancer. Oncotarget, 2016, 7, 24088-24096. | 1.8 | 15 |
| 98 | Enhancing the Therapeutic Efficacy of 2-Deoxyglucose in Breast Cancer Cells Using Cell-cycle Synchronization. Anticancer Research, 2016, 36, 5975-5980. | 1.1 | 8 |
| 99 | The Role of Plasma Chromogranin A as Assessment of Treatment Response in Non-functioning Gastroenteropancreatic Neuroendocrine Tumors. Cancer Research and Treatment, 2016, 48, 153-161. | 3.0 | 11 |
| 100 | A retrospective feasibility analysis of biweekly reduced dose docetaxel in Korean patients with castration-resistant, metastatic prostate cancer Journal of Clinical Oncology, 2016, 34, e638-e638. | 1.6 | 0 |
| 101 | Adjuvant chemotherapy with or without concurrent radiotherapy in stage IB patients with gastric cancer: Subgroup analysis of the adjuvant chemoradiotherapy in stomach tumors (ARTIST) phase III trial Journal of Clinical Oncology, 2016, 34, 95-95. | 1.6 | 0 |
| 102 | Tuberous sclerosis complex 2 (TSC2) expression in hepatocellular carcinoma to predict responses to mTOR inhibitor Journal of Clinical Oncology, 2016, 34, e15628-e15628. | 1.6 | 0 |
| 103 | The clinicopathologic features and treatment of 607 hindgut neuroendocrine tumor (NET) patients at a single institution Journal of Clinical Oncology, 2016, 34, 4091-4091. | 1.6 | 0 |
| 104 | The impact of cetuximab plus AKT- or mTOR- inhibitor in patient-derived colon cancer cell model with RAS wild type and PIK3CA mutation Journal of Clinical Oncology, 2016, 34, e15153-e15153. | 1.6 | 0 |
| 105 | Relation Between F-18 FDG Uptake of PET/CT and BRAFV600E Mutation in Papillary Thyroid Cancer. Medicine (United States), 2015, 94, e2063. | 1.0 | 17 |
| 106 | The Correlations Between MRI Perfusion, Diffusion Parameters, and 18F-FDG PET Metabolic Parameters in Primary Head-and-Neck Cancer. Medicine (United States), 2015, 94, e2141. | 1.0 | 31 |
| 107 | The Utility of Segmental Analysis in Cardiac I-123 MIBG SPECT in Parkinson's Disease. Nuclear Medicine and Molecular Imaging, 2015, 49, 298-302. | 1.0 | 5 |
| 108 | Prediction of occult lymph node metastasis using volume-based PET parameters in small-sized peripheral non-small cell lung cancer. Cancer Imaging, 2015, 15, 21. | 2.8 | 43 |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Phase II Trial of Nilotinib in Patients With Metastatic Malignant Melanoma Harboring <i>KIT</i> Gene Aberration: A Multicenter Trial of Korean Cancer Study Group (UN10-06). Oncologist, 2015, 20, 1312-1319. | 3.7 | 70 |
| 110 | Capecitabine in combination with either cisplatin or weekly paclitaxel as a first-line treatment for metastatic esophageal squamous cell carcinoma: a randomized phase II study. BMC Cancer, 2015, 15, 693. | 2.6 | 23 |
| 111 | Exploratory biomarker analysis for treatment response in KRAS wild type metastatic colorectal cancer patients who received cetuximab plus irinotecan. BMC Cancer, 2015, 15, 747. | 2.6 | 10 |
| 112 | Physiologic Facial Muscle Uptake on 18F-FDG PET/CT by Chewing-Like Habitual Movement in Patient With SjĶgren Syndrome. Clinical Nuclear Medicine, 2015, 40, 268-269. | 1.3 | 6 |
| 113 | Circulating Tumor Cells are Predictive of Poor Response to Chemotherapy in Metastatic gastric cancer. International Journal of Biological Markers, 2015, 30, 382-386. | 1.8 | 25 |
| 114 | Gastrointestinal malignancies harbor actionable MET exon 14 deletions. Oncotarget, 2015, 6, 28211-28222. | 1.8 | 57 |
| 115 | Single Pulmonary Langerhans Cell Histiocytosis Mimicking Metastasis in Lung Cancer: Imaging with 18F-FDG PET/CT. Nuclear Medicine and Molecular Imaging, 2015, 49, 243-245. | 1.0 | 2 |
| 116 | Prevalence and clinical outcomes of young breast cancer (YBC) patients according to intrinsic breast cancer subtypes: Single institutional experience in Korea. Breast, 2015, 24, 213-217. | 2.2 | 23 |
| 117 | Correlation Between F-18 Fluorodeoxyglucose Positron Emission Tomography Metabolic Parameters and Dynamic Contrast-Enhanced MRI-Derived Perfusion Data in Patients with Invasive Ductal Breast Carcinoma. Annals of Surgical Oncology, 2015, 22, 3866-3872. | 1.5 | 18 |
| 118 | Tumour shrinkage at 6Âweeks predicts favorable clinical outcomes in a phase III study of gemcitabine and oxaliplatin with or without erlotinib for advanced biliary tract cancer. BMC Cancer, 2015, 15, 530. | 2.6 | 17 |
| 119 | Efficacy of pazopanib monotherapy in patients who had been heavily pretreated for metastatic soft tissue sarcoma: a retrospective case series. BMC Cancer, 2015, 15, 154. | 2.6 | 58 |
| 120 | Pemetrexed plus platinum versus pemetrexed alone in non-small cell lung cancer patients who have progressed after first-line EGFR TKls. Lung Cancer, 2015, 90, 261-266. | 2.0 | 4 |
| 121 | Characterization of Durable Responder for Capecitabine Monotherapy in Patients With Anthracycline-and Taxane-Pretreated Metastatic Breast Cancer. Clinical Breast Cancer, 2015, 15, e287-e292. | 2.4 | 10 |
| 122 | Phase I trial and pharmacokinetic study of Tanibirumab, a fully human monoclonal antibody to the vascular endothelial growth factor receptor 2 in patients with refractory solid tumors Journal of Clinical Oncology, 2015, 33, 2522-2522. | 1.6 | 2 |
| 123 | Patient-derived cell models as preclinical tools for genome-directed targeted therapy. Oncotarget, 2015, 6, 25619-25630. | 1.8 | 48 |
| 124 | PIK3CA mutation detection in metastatic biliary cancer using cell-free DNA. Oncotarget, 2015, 6, 40026-40035. | 1.8 | 15 |
| 125 | NTRK1 rearrangement in colorectal cancer patients: evidence for actionable target using patient-derived tumor cell line. Oncotarget, 2015, 6, 39028-39035. | 1.8 | 53 |
| 126 | Salivary gland scintigraphy in Sjögren's syndrome. Nuklearmedizin - NuclearMedicine, 2014, 53, 139-145. | 0.7 | 18 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 127 | The Clinical Impacts of Apparent Embolic Event and the Predictors of In-Hospital Mortality in Patients with Infective Endocarditis. Journal of Korean Medical Science, 2014, 29, 1646. | 2.5 | 2 |
| 128 | Diagnostic performance of 18F-FDG PET/CT, ultrasonography and MRI. Nuklearmedizin - NuclearMedicine, 2014, 53, 89-94. | 0.7 | 41 |
| 129 | Pazopanib, a Novel Multitargeted Kinase Inhibitor, Shows Potent <i>In Vitro</i> Antitumor Activity in Gastric Cancer Cell Lines with <i>FGFR2</i> Amplification. Molecular Cancer Therapeutics, 2014, 13, 2527-2536. | 4.1 | 34 |
| 130 | Isolated Splenic Metastasis of Ovarian Cancer Detected With 18F-FDG PET/CT. Clinical Nuclear Medicine, 2014, 39, 349-351. | 1.3 | 6 |
| 131 | Oxidized Low-Density Lipoprotein Stimulates Macrophage ¹⁸ F-FDG Uptake via Hypoxia-Inducible Factor-11± Activation Through Nox2-Dependent Reactive Oxygen Species Generation. Journal of Nuclear Medicine, 2014, 55, 1699-1705. | 5.0 | 63 |
| 132 | Successful use of pazopanib for treatment of refractory metastatic hemangiopericytoma. Clinical Sarcoma Research, 2014, 4, 13. | 2.3 | 28 |
| 133 | Positive Correlation between Baseline PET or PET/CT Findings and Clinical Parameters in Multiple Myeloma Patients. Acta Haematologica, 2014, 131, 193-199. | 1.4 | 23 |
| 134 | Incidentally Detected Juvenile-Pattern Bone Scintigraphy in a Young Man with Kallmann's Syndrome. Nuclear Medicine and Molecular Imaging, 2014, 48, 247-248. | 1.0 | 0 |
| 135 | DCEP for relapsed or refractory multiple myeloma after therapy with novel agents. Annals of Hematology, 2014, 93, 99-105. | 1.8 | 29 |
| 136 | Prognostic significance of the intratumoral heterogeneity of ¹⁸ Fâ€FDG uptake in oral cavity cancer. Journal of Surgical Oncology, 2014, 110, 702-706. | 1.7 | 38 |
| 137 | Is there any predictor for clinical outcome in EGFR mutant NSCLC patients treated with EGFR TKIs?. Cancer Chemotherapy and Pharmacology, 2014, 73, 1063-1070. | 2.3 | 18 |
| 138 | Safety and efficacy of gemcitabine or pemetrexed in combination with a platinum in patients with non-small-cell lung cancer and prior interstitial lung disease. Cancer Chemotherapy and Pharmacology, 2014, 73, 1217-1225. | 2.3 | 32 |
| 139 | Significance of Salivary Gland Radioiodine Retention on Post-ablation 1311 Scintigraphy as a Predictor of Salivary Gland Dysfunction in Patients with Differentiated Thyroid Carcinoma. Nuclear Medicine and Molecular Imaging, 2014, 48, 203-211. | 1.0 | 11 |
| 140 | Carotid Artery FDG Uptake May Serve as a Biomarker for Cardiovascular Risk Stratification in Asymptomatic Adults. Nuclear Medicine and Molecular Imaging, 2014, 48, 196-202. | 1.0 | 8 |
| 141 | Natural history of untreated patients who had metastatic biliary tract cancer (BTC) with good performance status (PS) Journal of Clinical Oncology, 2014, 32, e15106-e15106. | 1.6 | 0 |
| 142 | Adjuvant chemoradiation with 5-fluorouracil/leucovorin versus S-1 in gastric cancer patients following D2 lymph node dissection surgery: a feasibility study. Anticancer Research, 2014, 34, 6585-91. | 1.1 | 2 |
| 143 | Usefulness of 18F-fluoride PET/CT in Breast Cancer Patients with Osteosclerotic Bone Metastases. Nuclear Medicine and Molecular Imaging, 2013, 47, 27-35. | 1.0 | 33 |
| 144 | Symptomatic late-onset sialadenitis after radioiodine therapy in thyroid cancer. Annals of Nuclear Medicine, 2013, 27, 386-391. | 2,2 | 33 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 145 | Inclusion of hemoglobin level in prognostic score provides better prognostic stratification in patients with acute promyelocytic leukemia (APL). International Journal of Hematology, 2013, 97, 388-396. | 1.6 | 6 |
| 146 | Clinical implication of Time To Brain Metastasis (TTBM) according to breast cancer subtypes. SpringerPlus, 2013, 2, 136. | 1.2 | 18 |
| 147 | Phase II trial of capecitabine and everolimus (RAD001) combination in refractory gastric cancer patients. Investigational New Drugs, 2013, 31, 1580-1586. | 2.6 | 33 |
| 148 | Symptomatic brain involvement as the initial manifestation of neuromyelitis optica. Journal of Clinical Neuroscience, 2013, 20, 938-942. | 1.5 | 9 |
| 149 | Gastric cancer (GC) patients with hedgehog pathway activation: PTCH1 and GLI2 as independent prognostic factors. Targeted Oncology, 2013, 8, 271-280. | 3.6 | 25 |
| 150 | Identification of <i>ROS1</i> rearrangement in gastric adenocarcinoma. Cancer, 2013, 119, 1627-1635. | 4.1 | 108 |
| 151 | Incidental Focal Uptake in Oropharynx Due to Lingual Thyroid on Diagnostic 1311 Whole-Body Scan After Total Thyroidectomy. Clinical Nuclear Medicine, 2013, 38, 646-648. | 1.3 | 2 |
| 152 | Effect of paclitaxel/carboplatin salvage chemotherapy in noncutaneous versus cutaneous metastatic melanoma. Melanoma Research, 2013, 23, 147-151. | 1.2 | 34 |
| 153 | Decreased Metabolism in the Cerebral Cortex in Early-Stage Huntington's Disease: A Possible | | |
| | | | |

| # | Article | IF | CITATIONS |
|-----|---|-----|-----------|
| 163 | A survey of c-MET expression and amplification in 287 patients with hepatocellular carcinoma. Anticancer Research, 2013, 33, 5179-86. | 1.1 | 42 |
| 164 | Apparent Relative Hypermetabolism of Selective Brain Areas in Huntington Disease and Importance of Reference Region for Analysis. Clinical Nuclear Medicine, 2012, 37, 663-668. | 1.3 | 12 |
| 165 | Primary Malignant Fibrous Histiocytoma in Mediastinum: Imaging with 18F-FDG PET/CT. Nuclear Medicine and Molecular Imaging, 2012, 46, 304-307. | 1.0 | 9 |
| 166 | ALK inhibitor crizotinib combined with intrathecal methotrexate treatment for non-small cell lung cancer with leptomeningeal carcinomatosis. Lung Cancer, 2012, 76, 253-254. | 2.0 | 26 |
| 167 | Unique perception of clinical trials by Korean cancer patients. BMC Cancer, 2012, 12, 594. | 2.6 | 21 |
| 168 | Prognostic Value of Volume-Based < sup > 18 < /sup > F-Fluorodeoxyglucose PET/CT Parameters in Patients with Clinically Node-Negative Oral Tongue Squamous Cell Carcinoma. Korean Journal of Radiology, 2012, 13, 752. | 3.4 | 48 |
| 169 | Phase 1 trial of S-1 in combination with sorafenib for patients with advanced hepatocellular carcinoma. Investigational New Drugs, 2012, 30, 1540-1547. | 2.6 | 14 |
| 170 | Resting cerebral glucose metabolism and perfusion patterns in women with posttraumatic stress disorder related to sexual assault. Psychiatry Research - Neuroimaging, 2012, 201, 214-217. | 1.8 | 22 |
| 171 | Analysis of prognostic factors and 5-year survival rate in patients with hepatocellular carcinoma: a single-center experience. The Korean Journal of Hepatology, 2012, 18, 48. | 1.5 | 34 |
| 172 | A prospective, phase II trial of induction chemotherapy with docetaxel/cisplatin for Masaoka stage III/IV thymic epithelial tumors Journal of Clinical Oncology, 2012, 30, 7104-7104. | 1.6 | 0 |
| 173 | Non Malignant Late Complications After Allogeneic Hematopoietic Stem Cell Transplantation (allo-HSCT); Korean Experience. Blood, 2012, 120, 4462-4462. | 1.4 | 0 |
| 174 | Correlation Between Unidentified Bright Objects on Brain Magnetic Resonance Imaging (MRI) and Cerebral Glucose Metabolism in Patients with Neurofibromatosis Type 1. Journal of Genetic Medicine, 2012, 9, 84-88. | 0.2 | 0 |
| 175 | Evaluation of Agreement between 64-slice Computed Tomography Angiography and 201-TI Single Photon Emission Computed Tomography-myocardial Perfusion Imaging in the Diagnosis of Significant Coronary Artery Disease. Journal of the Korean Society of Radiology, 2011, 64, 541. | 0.2 | 0 |
| 176 | Incidental Multiple Focal Uptakes of Tc-99m MDP in Both Lungs Without Lung Parenchymal Abnormality. Clinical Nuclear Medicine, 2011, 36, 221-223. | 1.3 | 4 |
| 177 | Thymidylate synthase and thymidine phosphorylase as predictive markers of capecitabine monotherapy in patients with anthracycline- and taxane-pretreated metastatic breast cancer. Cancer Chemotherapy and Pharmacology, 2011, 68, 743-751. | 2.3 | 27 |
| 178 | Phase I trial of capecitabine plus everolimus (RAD001) in patients with previously treated metastatic gastric cancer. Cancer Chemotherapy and Pharmacology, 2011, 68, 255-262. | 2.3 | 31 |
| 179 | Salvage S-1 monotherapy in metastatic colorectal cancer patients who failed irinotecan-based or oxaliplatin-based chemotherapy. Medical Oncology, 2011, 28, 291-294. | 2.5 | 5 |
| 180 | Clinical Significance of Diffuse 18F-FDG Uptake in Residual Thyroid Gland after Unilateral Thyroid Lobectomy. Nuclear Medicine and Molecular Imaging, 2011, 45, 197-202. | 1.0 | 1 |

| # | Article | IF | Citations |
|-----|--|-----|-----------|
| 181 | Clinical presentation of carcinoma of unknown primary: 14Âyears of experience. Tumor Biology, 2011, 32, 45-51. | 1.8 | 4 |
| 182 | Retrospective multicenter study on the development of peripheral lymphocytosis following secondâ€line dasatinib therapy for chronic myeloid leukemia. American Journal of Hematology, 2011, 86, 346-350. | 4.1 | 24 |
| 183 | Analysis of Prognostic Factors of Pediatric-Type Sarcomas in Adult Patients. Oncology, 2011, 80, 21-28. | 1.9 | 9 |
| 184 | Implications of Bone-Only Metastases in Breast Cancer: Favorable Preference with Excellent Outcomes of Hormone Receptor Positive Breast Cancer. Cancer Research and Treatment, 2011, 43, 89-95. | 3.0 | 79 |
| 185 | Abstract B38: Troglitazone enhances breast cancer cell18F-FDG uptake and intracelluar lactic acid accumulation through PPAR-rindependent activation of the MAPK pathway, 2011, , . | | O |
| 186 | Prognostic Value of Metabolic Tumor Volume Measured by 18F-Fluorodeoxyglucose Positron Emission Tomography in Patients with Esophageal Carcinoma. Annals of Surgical Oncology, 2010, 17, 115-122. | 1.5 | 194 |
| 187 | Survival benefit from ovarian metastatectomy in colorectal cancer patients with ovarian metastasis: a retrospective analysis. Cancer Chemotherapy and Pharmacology, 2010, 66, 229-235. | 2.3 | 49 |
| 188 | Extraosseous osteosarcoma: Single institutional experience in Korea. Asia-Pacific Journal of Clinical Oncology, 2010, 6, 126-129. | 1.1 | 20 |
| 189 | Multicenter Retrospective Study on the Development of Peripheral Lymphocytosis During Second-Line Dasatinib Therapy for Chronic Myeloid Leukemia. Blood, 2010, 116, 2275-2275. | 1.4 | 1 |
| 190 | Treatment outcomes and clinicopathologic characteristics of tripleâ€negative breast cancer patients who received platinumâ€containing chemotherapy. International Journal of Cancer, 2009, 124, 1457-1462. | 5.1 | 69 |
| 191 | Prognostic model to predict outcomes in nonsmall cell lung cancer patients treated with gefitinib as a salvage treatment. Cancer, 2009, 115, 1518-1530. | 4.1 | 20 |
| 192 | Atypical basal ganglia germinoma presenting as cerebral hemiatrophy: diagnosis and follow-up with 11C-methionine positron emission tomography. Child's Nervous System, 2009, 25, 29-37. | 1.1 | 27 |
| 193 | Rituximab-induced acute thrombocytopenia: a case report and review of the literature. Medical Oncology, 2009, 26, 45-48. | 2.5 | 28 |
| 194 | Variation in FDG uptake on PET in patients with radiation-induced pelvic insufficiency fractures: a review of 10 cases. Annals of Nuclear Medicine, 2009, 23, 511-516. | 2.2 | 9 |
| 195 | Modified Transobturator Tape (Canal Transobturator Tape) Surgery for Female Stress Urinary Incontinence. Journal of Urology, 2009, 181, 2616-2621. | 0.4 | 11 |
| 196 | Myocardial perfusion defects and coronary risk factors in symptomatic and asymptomatic elderly women. International Journal of Cardiovascular Imaging, 2008, 24, 277-281. | 1.5 | 8 |
| 197 | Comparison of Uptake Characteristics and Prognostic Value of ²⁰¹ Tl and ¹⁸ Fâ€FDG in Esophageal Cancer. World Journal of Surgery, 2008, 32, 69-75. | 1.6 | 6 |
| 198 | Potential value of radionuclide cisternography in diagnosis and management planning of spontaneous intracranial hypotension. Clinical Neurology and Neurosurgery, 2008, 110, 657-661. | 1.4 | 37 |

Su Jin Lee

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 199 | Reversal of Vascular ¹⁸ F-FDG Uptake with Plasma High-Density Lipoprotein Elevation by Atherogenic Risk Reduction. Journal of Nuclear Medicine, 2008, 49, 1277-1282. | 5.0 | 108 |
| 200 | Neutrophil Counts Could Predict the Response to Immunosuppressive Treatment with Antithymocyte Globulin and Cyclosporine in the Patients with Severe Aplastic Anemia. Blood, 2008, 112, 4109-4109. | 1.4 | 8 |
| 201 | BCL2 Gene Polymorphism Could Predict the Treatment Outcomes in Patients with De Novo Acute Myeloid Leukemia. Blood, 2008, 112, 3973-3973. | 1.4 | O |
| 202 | Localization of Dystonic Muscles with 18F-FDG PET/CT in Idiopathic Cervical Dystonia. Journal of Nuclear Medicine, 2007, 48, 1790-1795. | 5.0 | 40 |
| 203 | Pazopanib monotherapy in the treatment of pretreated, metastatic uterine sarcoma: a single-center retrospective study. Journal of Gynecologic Oncology, 0, 29, . | 2.2 | 0 |