

Karel Kostev

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3412860/publications.pdf>

Version: 2024-02-01

477
papers

6,652
citations

109321

35
h-index

149698

56
g-index

506
all docs

506
docs citations

506
times ranked

7779
citing authors

#	ARTICLE	IF	CITATIONS
1	Basic characteristics and representativeness of the German Disease Analyzer database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 459-466.	0.6	261
2	Validity and representativeness of the "Disease Analyzer" patient database for use in pharmaco-epidemiological and pharmacoeconomic studies. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2009, 47, 617-626.	0.6	237
3	Prevalence, utilization, and costs of antiepileptic drugs for epilepsy in Germany—a nationwide population-based study in children and adults. <i>Journal of Neurology</i> , 2012, 259, 2376-2384.	3.6	116
4	GRAND: the German retrospective cohort analysis on compliance and persistence and the associated risk of fractures in osteoporotic women treated with oral bisphosphonates. <i>Osteoporosis International</i> , 2012, 23, 223-231.	3.1	114
5	Incidence of newly diagnosed diabetes after Covid-19. <i>Diabetologia</i> , 2022, 65, 949-954.	6.3	92
6	Impact of the COVID-19 Pandemic on Cancer Diagnoses in General and Specialized Practices in Germany. <i>Cancers</i> , 2021, 13, 408.	3.7	90
7	Persistence in patients with breast cancer treated with tamoxifen or aromatase inhibitors: a retrospective database analysis. <i>Breast Cancer Research and Treatment</i> , 2013, 138, 185-191.	2.5	85
8	No reduced risk of overall, colorectal, lung, breast, and prostate cancer with metformin therapy in diabetic patients: database analyses from Germany and the UK. <i>Pharmacoepidemiology and Drug Safety</i> , 2015, 24, 865-874.	1.9	81
9	Risk factors for dementia diagnosis in German primary care practices. <i>International Psychogeriatrics</i> , 2016, 28, 1059-1065.	1.0	79
10	Treatment persistence, hypoglycaemia and clinical outcomes in type 2 diabetes patients with dipeptidyl peptidase-4 inhibitors and sulphonylureas: a primary care database analysis. <i>Diabetes, Obesity and Metabolism</i> , 2013, 15, 55-61.	4.4	78
11	Amputation rate and risk factors in type 2 patients with diabetic foot syndrome under real-life conditions in Germany. <i>Primary Care Diabetes</i> , 2012, 6, 241-246.	1.8	76
12	Prevalence of diabetic foot syndrome and its risk factors in the UK. <i>Journal of Wound Care</i> , 2010, 19, 333-337.	1.2	74
13	Diabetic retinopathy at diagnosis of type 2 diabetes in the UK: a database analysis. <i>Diabetologia</i> , 2013, 56, 109-111.	6.3	73
14	Basic characteristics and representativeness of the German Disease Analyzer database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 459-466.	0.6	73
15	Effect of the COVID-19 lockdown on disease recognition and utilisation of healthcare services in the older population in Germany: a cross-sectional study. <i>Age and Ageing</i> , 2021, 50, 317-325.	1.6	71
16	Nonalcoholic Fatty Liver Disease Increases the Risk of Anxiety and Depression. <i>Hepatology Communications</i> , 2020, 4, 1293-1301.	4.3	70
17	Impact of NAFLD on the Incidence of Cardiovascular Diseases in a Primary Care Population in Germany. <i>Digestive Diseases and Sciences</i> , 2020, 65, 2112-2119.	2.3	68
18	Persistence and compliance of medications used in the treatment of osteoporosis—analysis using a large scale, representative, longitudinal German database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 315-322.	0.6	68

#	ARTICLE	IF	CITATIONS
19	Impact of tobacco smoking on the risk of developing 25 different cancers in the UK: a retrospective study of 422,010 patients followed for up to 30 years. <i>Oncotarget</i> , 2018, 9, 17420-17429.	1.8	67
20	Adherence to GLP-1 receptor agonist therapy administered by once-daily or once-weekly injection in patients with type 2 diabetes in Germany. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, Volume 9, 201-205.	2.4	65
21	Nonadherence to antiepileptic drugs in Germany. <i>Neurology</i> , 2016, 87, 466-472.	1.1	62
22	Persistence with antihypertensive treatments: results of a 3-year follow-up cohort study. <i>European Journal of Clinical Pharmacology</i> , 2007, 63, 1055-1061.	1.9	61
23	Isopropanolic black cohosh extract and recurrence-free survival after breast cancer. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2007, 45, 143-154.	0.6	60
24	Treatment patterns and low-density lipoprotein cholesterol (LDL-C) goal attainment among patients receiving high- or moderate-intensity statins. <i>Clinical Research in Cardiology</i> , 2018, 107, 380-388.	3.3	59
25	Fracture risk in patients with newly diagnosed type 2 diabetes: a retrospective database analysis in primary care. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 766-770.	2.3	56
26	Use of dipeptidyl peptidase-4 inhibitors and risk of bone fracture in patients with type 2 diabetes in Germany – A retrospective analysis of real-world data. <i>Osteoporosis International</i> , 2017, 28, 2421-2428.	3.1	52
27	Real-world treatment patterns of sacubitril/valsartan: a longitudinal cohort study in Germany. <i>European Journal of Heart Failure</i> , 2019, 21, 588-597.	7.1	49
28	Prevalence and risk factors of neuropathy in newly diagnosed type 2 diabetes in primary care practices: A retrospective database analysis in Germany and UK. <i>Primary Care Diabetes</i> , 2014, 8, 250-255.	1.8	46
29	Non-alcoholic fatty liver disease increases the risk of incident chronic kidney disease. <i>United European Gastroenterology Journal</i> , 2020, 8, 942-948.	3.8	43
30	Panic buying or good adherence? Increased pharmacy purchases of drugs from wholesalers in the last week prior to Covid-19 lockdown. <i>Journal of Psychiatric Research</i> , 2020, 130, 19-21.	3.1	43
31	Prevalence of depression, anxiety and their risk factors in German women with breast cancer in general and gynecological practices. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 447-452.	2.5	42
32	Are Sulfonylurea and Insulin Therapies Associated With a Larger Risk of Cancer Than Metformin Therapy? A Retrospective Database Analysis. <i>Diabetes Care</i> , 2015, 38, 59-65.	8.6	41
33	Depression Risk in Patients with Rheumatoid Arthritis in the United Kingdom. <i>Rheumatology and Therapy</i> , 2017, 4, 195-200.	2.3	41
34	Association Between the Use of Antihyperglycemic Drugs and Dementia Risk: A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 725-732.	2.6	40
35	Switching the manufacturer of antiepileptic drugs is associated with higher risk of seizures: A nationwide study of prescription data in Germany. <i>Annals of Neurology</i> , 2018, 84, 918-925.	5.3	38
36	Prevalence of depression in type 2 diabetes patients in German primary care practices. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 432-437.	2.3	36

#	ARTICLE	IF	CITATIONS
37	Antiemetic Medications in Children with Presumed Infectious Gastroenteritis”Pharmacoepidemiology in Europe and Northern America. <i>Journal of Pediatrics</i> , 2008, 153, 659-662.e3.	1.8	35
38	The impact of depot medroxyprogesterone acetate on fracture risk: a case-control study from the UK. <i>Osteoporosis International</i> , 2017, 28, 291-297.	3.1	35
39	Impact of Rotavirus Vaccination in Germany. <i>Pediatric Infectious Disease Journal</i> , 2014, 33, e299-e304.	2.0	33
40	Discontinuation of Treatment Using Anticholinergic Medications in Patients With Urinary Incontinence. <i>Obstetrics and Gynecology</i> , 2014, 124, 794-800.	2.4	32
41	The impact of treatment compliance on fracture risk in women with breast cancer treated with aromatase inhibitors in the United Kingdom. <i>Breast Cancer Research and Treatment</i> , 2016, 155, 151-157.	2.5	32
42	Association Between Hearing and Vision Impairment and Risk of Dementia: Results of a Case-Control Study Based on Secondary Data. <i>Frontiers in Aging Neuroscience</i> , 2019, 11, 363.	3.4	32
43	Incident Dementia in Elderly Patients with Nonalcoholic Fatty Liver Disease in Germany. <i>Digestive Diseases and Sciences</i> , 2021, 66, 3179-3185.	2.3	32
44	Impact of the coronavirus 2019 (COVID-19) pandemic on anxiety diagnosis in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021, 143, 528-533.	3.1	32
45	Fewer non-”COVID-19 respiratory tract infections and gastrointestinal infections during the COVID-19 pandemic. <i>Journal of Medical Virology</i> , 2022, 94, 298-302.	5.0	32
46	Predictors of hypoglycaemia in insulin-treated type 2 diabetes patients in primary care: A retrospective database analysis. <i>Primary Care Diabetes</i> , 2014, 8, 127-131.	1.8	31
47	Heart failure is associated with an increased incidence of cancer diagnoses. <i>ESC Heart Failure</i> , 2021, 8, 3628-3633.	3.1	31
48	Obesity and lipid metabolism disorders determine the risk for development of long COVID syndrome: a cross-sectional study from 50,402 COVID-19 patients. <i>Infection</i> , 2022, 50, 1165-1170.	4.7	31
49	Association Between Migraine Headaches and Dementia in More than 7,400 Patients Followed in General Practices in the United Kingdom. <i>Journal of Alzheimer’s Disease</i> , 2019, 71, 353-360.	2.6	30
50	Continuous treatment with antidementia drugs in Germany 2003-2013: a retrospective database analysis. <i>International Psychogeriatrics</i> , 2015, 27, 1335-1342.	1.0	29
51	Incidence of depression and anxiety among women newly diagnosed with breast or genital organ cancer in Germany. <i>Psycho-Oncology</i> , 2017, 26, 1535-1540.	2.3	28
52	Antihypertensive treatment and risk of dementia: a retrospective database study. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 195-201.	0.6	28
53	Impact of metformin on metastases in patients with breast cancer and type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1056-1059.	2.3	27
54	Long-term use of benzodiazepines in older patients in Germany: a retrospective analysis. <i>Therapeutic Advances in Psychopharmacology</i> , 2017, 7, 191-200.	2.7	27

#	ARTICLE	IF	CITATIONS
55	The tamoxifen paradox—“influence of adjuvant tamoxifen on fracture risk in pre- and postmenopausal women with breast cancer. <i>Osteoporosis International</i> , 2018, 29, 2557-2564.	3.1	27
56	Prevalence of chronic diseases among older patients in German general practices. <i>GMS German Medical Science</i> , 2016, 14, Doc03.	2.7	27
57	Prevalence of and risk factors associated with cesarean section in Lebanon —“ A retrospective study based on a sample of 29,270 women. <i>Women and Birth</i> , 2017, 30, e265-e271.	2.0	26
58	Prevalence of vulvovaginal candidiasis in gynecological practices in Germany: A retrospective study of 954,186 patients. <i>Current Medical Mycology</i> , 2018, 4, 6-11.	0.8	26
59	Antiherpetic medication and incident dementia: Observational cohort studies in four countries. <i>European Journal of Neurology</i> , 2021, 28, 1840-1848.	3.3	26
60	Association Between Inflammatory Bowel Disease and Dementia: A Retrospective Cohort Study. <i>Journal of Alzheimer's Disease</i> , 2021, 80, 1471-1478.	2.6	26
61	An elevated FIB-4 score predicts liver cancer development: A longitudinal analysis from 29,999 patients with NAFLD. <i>Journal of Hepatology</i> , 2022, 76, 247-248.	3.7	25
62	Cost comparison of outpatient treatment with granulocyte colony-stimulating factors (G-CSF) in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 281-289.	0.6	24
63	Risk of venous thrombosis in users of hormonal contraceptives in German gynaecological practices: a patient database analysis. <i>Archives of Gynecology and Obstetrics</i> , 2014, 289, 413-419.	1.7	24
64	Sociodemographic Disparities in Administration of Antiepileptic Drugs to Adults with Epilepsy in Germany: A Retrospective, Database Study of Drug Prescriptions. <i>CNS Drugs</i> , 2014, 28, 753-759.	5.9	24
65	Differences in persistency with teriparatide in patients with osteoporosis according to gender and health care provider. <i>Osteoporosis International</i> , 2014, 25, 2721-2728.	3.1	24
66	Changes in HbA1c, body weight, and systolic blood pressure in type 2 diabetes patients initiating dapagliflozin therapy: a primary care database study. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, Volume 9, 337-345.	2.4	24
67	Association between thyroid gland diseases and breast cancer: a case—“control study. <i>Breast Cancer Research and Treatment</i> , 2020, 182, 207-213.	2.5	24
68	Prescription patterns and drug costs in German patients with dementia in nursing homes and home-care settings. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 9-15.	0.6	24
69	Fracture risk in patients with type 2 diabetes under different antidiabetic treatment regimens: a retrospective database analysis in primary care. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, 9, 17.	2.4	23
70	Characteristics of diabetic patients visiting a podiatry practice in Germany. <i>Journal of Wound Care</i> , 2010, 19, 140-148.	1.2	22
71	Induction of labor in breech presentations at term: a retrospective observational study. <i>Archives of Gynecology and Obstetrics</i> , 2016, 293, 549-555.	1.7	22
72	Impact of prior breast cancer on mode of delivery and pregnancy-associated disorders: a retrospective analysis of subsequent pregnancy outcomes. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1069-1074.	2.5	22

#	ARTICLE	IF	CITATIONS
73	Prevalence of depression, anxiety, and adjustment disorders in women with spontaneous abortion in Germany – A retrospective cohort study. <i>Psychiatry Research</i> , 2017, 258, 382-386.	3.3	22
74	Impact of Osteoporosis on the Risk of Dementia in Almost 60,000 Patients Followed in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 401-407.	2.6	22
75	Reduced antibiotic use after initial treatment of acute respiratory infections with phytopharmaceuticals- a retrospective cohort study. <i>Postgraduate Medicine</i> , 2020, 132, 412-418.	2.0	22
76	Prevalence of overuse of short-acting beta-2 agonists (SABA) and associated factors among patients with asthma in Germany. <i>Respiratory Research</i> , 2021, 22, 108.	3.6	22
77	Gender- and age-related treatment compliance in patients with osteoporosis in Germany. <i>Patient Preference and Adherence</i> , 2016, Volume 10, 2379-2385.	1.8	21
78	Prevalence of medically treated children with ADHD and type 1 diabetes in Germany – Analysis of two representative databases. <i>Journal of Pediatric Endocrinology and Metabolism</i> , 2016, 29, 1293-1297.	0.9	21
79	Adherence to antiepileptic drugs in children and adolescents: A retrospective study in primary care settings in Germany. <i>Epilepsy and Behavior</i> , 2017, 75, 36-41.	1.7	21
80	Prevalence of, and factors associated with, long-term COVID-19 sick leave in working-age patients followed in general practices in Germany. <i>International Journal of Infectious Diseases</i> , 2021, 109, 203-208.	3.3	21
81	Effect of the COVID-19 pandemic on stress, anxiety, and depressive disorders in German primary care: A cross-sectional study. <i>Journal of Psychiatric Research</i> , 2021, 143, 43-49.	3.1	21
82	Tumor Incidence in Patients with Non-Alcoholic Fatty Liver Disease. <i>Deutsches A&#x0308;rztblatt International</i> , 2020, 117, 719-724.	0.9	21
83	Changes in time to insulin initiation in type 2 diabetes patients: A retrospective database analysis in Germany and UK (2005–2010). <i>Primary Care Diabetes</i> , 2013, 7, 229-233.	1.8	20
84	Impact of Disease Management Programs on HbA1c Values in Type 2 Diabetes Patients in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 117-122.	2.2	20
85	Depression risk in patients with late-onset rheumatoid arthritis in Germany. <i>Quality of Life Research</i> , 2017, 26, 437-443.	3.1	20
86	Multimorbidity and polypharmacy among elderly people followed in general practices in Germany. <i>European Journal of Internal Medicine</i> , 2018, 55, 66-68.	2.2	20
87	Association between chronic somatic conditions and depression in children and adolescents: A retrospective study of 13,326 patients. <i>Journal of Affective Disorders</i> , 2019, 245, 697-701.	4.1	20
88	Glycemic control after initiating basal insulin therapy in patients with type 2 diabetes: a primary care database analysis. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2015, 8, 45.	2.4	19
89	Impact of disease management programs on women with breast cancer in Germany. <i>Breast Cancer Research and Treatment</i> , 2015, 153, 391-395.	2.5	19
90	Depression risk in patients with coronary heart disease in Germany. <i>World Journal of Cardiology</i> , 2016, 8, 547.	1.5	19

#	ARTICLE	IF	CITATIONS
91	Association of dipeptidyl peptidase 4 inhibitors with risk of metastases in patients with type 2 diabetes and breast, prostate or digestive system cancer. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 687-692.	2.3	19
92	Association between induced abortion, spontaneous abortion, and infertility respectively and the risk of psychiatric disorders in 57,770 women followed in gynecological practices in Germany. <i>Journal of Affective Disorders</i> , 2019, 251, 107-113.	4.1	19
93	Progression of mild cognitive impairment to dementia in German specialist practices. <i>Dementia</i> , 2019, 18, 380-390.	2.0	19
94	Drug treatment for patients with bipolar disorders in psychiatric practices in Germany in 2009 and 2018. <i>Psychiatry Research</i> , 2020, 289, 112965.	3.3	19
95	Association Between Low-Density Lipoprotein Cholesterol Levels, Statin Use, and Dementia in Patients followed in German General Practices. <i>Journal of Alzheimer's Disease</i> , 2021, 79, 37-46.	2.6	19
96	Real-life treatment patterns, compliance, persistence, and medication costs in patients with hypertension in Germany. <i>Journal of Medical Economics</i> , 2012, 15, 155-165.	2.1	18
97	Trends and patterns of hormonal contraceptive prescribing for adolescents in primary care in the UK. <i>Journal of Family Planning and Reproductive Health Care</i> , 2015, 41, 216-222.	0.8	18
98	Do patients with lactose intolerance exhibit more frequent comorbidities than patients without lactose intolerance? An analysis of routine data from German medical practices. <i>Annals of Gastroenterology</i> , 2016, 29, 174-9.	0.6	18
99	Oral contraceptive use and fracture risk—a retrospective study of 12,970 women in the UK. <i>Osteoporosis International</i> , 2017, 28, 2349-2355.	3.1	18
100	Depression and sleep disorders are associated with early mortality in women with breast cancer in the United Kingdom. <i>Journal of Psychiatric Research</i> , 2021, 143, 481-484.	3.1	18
101	Consultations Decline for Stroke, Transient Ischemic Attack, and Myocardial Infarction during the COVID-19 Pandemic in Germany. <i>Neuroepidemiology</i> , 2021, 55, 70-78.	2.3	18
102	Risk of psychiatric and neurological diseases in patients with workplace mobbing experience in Germany: a retrospective database analysis. <i>GMS German Medical Science</i> , 2014, 12, Doc10.	2.7	18
103	Treatment persistence after initiating basal insulin in type 2 diabetes patients: A primary care database analysis. <i>Primary Care Diabetes</i> , 2015, 9, 377-384.	1.8	17
104	The use of proton pump inhibitors is positively associated with osteoporosis in postmenopausal women in Germany. <i>Climacteric</i> , 2016, 19, 478-481.	2.4	17
105	Hip fracture risk in patients with dementia in German primary care practices. <i>Dementia</i> , 2017, 16, 853-864.	2.0	17
106	Risk factors and outcomes in œwell-selectedœ vaginal breech deliveries: a retrospective observational study. <i>Journal of Perinatal Medicine</i> , 2017, 45, 291-297.	1.4	17
107	Prevalence of pain medication prescriptions in France, Germany, and the UK - a cross-sectional study including 4,270,142 patients. <i>Postgraduate Medicine</i> , 2018, 130, 32-36.	2.0	17
108	Association of laryngeal cancer with vocal cord leukoplakia and associated risk factors in 1,184 patients diagnosed in otorhinolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018, 8, 689-693.	1.0	17

#	ARTICLE	IF	CITATIONS
109	Impact of dronedarone on the risk of myocardial infarction and stroke in atrial fibrillation patients followed in general practices in Germany. <i>International Journal of Cardiology</i> , 2019, 278, 126-132.	1.7	17
110	Is loneliness associated with mild cognitive impairment in low- and middle-income countries?. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 1345-1353.	2.7	17
111	The risk of developing depression when suffering from neurological diseases. <i>GMS German Medical Science</i> , 2013, 11, Doc02.	2.7	17
112	Investigation of the prevalence of non-COVID-19 infectious diseases during the COVID-19 pandemic. <i>Public Health</i> , 2022, 203, 53-57.	2.9	17
113	Lower incidence of recorded cardiovascular outcomes in patients with type 2 diabetes using insulin aspart vs. those on human regular insulin: observational evidence from general practices. <i>Diabetes, Obesity and Metabolism</i> , 2013, 15, 358-363.	4.4	16
114	Treatment of depression in cancer and non-cancer patients in German neuropsychiatric practices. <i>Psycho-Oncology</i> , 2016, 25, 1324-1328.	2.3	16
115	Depression risk in female patients with osteoporosis in primary care practices in Germany. <i>Osteoporosis International</i> , 2016, 27, 2739-2744.	3.1	16
116	Diagnoses of Dizziness- and Vertigo-related Disorders in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2018, 39, 474-480.	1.3	16
117	Association between sleep disorders and the presence of breast cancer metastases in gynecological practices in Germany: a case-control study of 11,412 women. <i>Breast Cancer Research and Treatment</i> , 2018, 171, 443-448.	2.5	16
118	The impact of the coronavirus disease (COVID-19) pandemic on outpatient epilepsy care: An analysis of physician practices in Germany. <i>Epilepsy and Behavior</i> , 2021, 117, 107833.	1.7	16
119	Treatment of Urinary Tract Infections with Canephron® in Germany: A Retrospective Database Analysis. <i>Antibiotics</i> , 2021, 10, 685.	3.7	16
120	Potential delay in the diagnosis of vulvar cancer and associated risk factors in women treated in German gynecological practices. <i>Oncotarget</i> , 2018, 9, 8725-8730.	1.8	16
121	Prescription rates of cardiovascular and diabetes therapies prior to and during the COVID-19 lockdown in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 475-481.	0.6	16
122	Increase in depression and anxiety disorder diagnoses during the COVID-19 pandemic in children and adolescents followed in pediatric practices in Germany. <i>European Child and Adolescent Psychiatry</i> , 2023, 32, 873-879.	4.7	16
123	Low blood levels of high-density lipoprotein (HDL) cholesterol are positively associated with cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 3039-3046.	2.5	16
124	Rotavirus Vaccination in Germany. <i>Pediatric Infectious Disease Journal</i> , 2011, 30, e244-e247.	2.0	15
125	Discontinuation rates of menopausal hormone therapy among postmenopausal women in the post-WHI study era. <i>Climacteric</i> , 2015, 18, 737-742.	2.4	15
126	Changes in Type 2 Diabetes Mellitus Patients in German Primary Care Prior to (2006) and After (2010.) <i>TJ ETQq0 0 0 rgBT /Overlock 10 T</i>	2.25	15

#	ARTICLE	IF	CITATIONS
127	Prevalence and risk factors for delirium diagnosis in patients followed in general practices in Germany. <i>International Psychogeriatrics</i> , 2018, 30, 511-518.	1.0	15
128	Change in glucose-lowering medication regimens in individuals with type 2 diabetes mellitus during the COVID-19 pandemic in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 910-915.	4.4	15
129	To Be Continued? Long-Term Treatment Effects of Antidepressant Drug Classes and Individual Antidepressants on the Risk of Developing Dementia. <i>Journal of Clinical Psychiatry</i> , 2020, 81, .	2.2	15
130	Adherence in tamoxifen therapy after conversion to a rebate pharmaceutical in breast cancer patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013, 51, 969-975.	0.6	15
131	The association between antihypertensive therapy and the incidence of Parkinson's disease in patients followed in general practices in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 483-488.	0.6	15
132	Decrease in the number of patients diagnosed with cancer during the COVID-19 pandemic in Germany. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 3117-3123.	2.5	15
133	Drug utilization patterns and reported health status in ethnic German migrants (Aussiedler) in Germany: a cross-sectional study. <i>BMC Public Health</i> , 2011, 11, 509.	2.9	14
134	Change in glycated haemoglobin levels after initiating second-line therapy in type 2 diabetes: a primary care database study. <i>Diabetes, Obesity and Metabolism</i> , 2016, 18, 840-843.	4.4	14
135	Comorbid disorders associated with asthma in children in Germany – National analysis of pediatric primary care data. <i>Pediatric Allergy and Immunology</i> , 2016, 27, 861-866.	2.6	14
136	Compliance with vaccination against tick-borne encephalitis virus in Germany. <i>Clinical Microbiology and Infection</i> , 2017, 23, 460-463.	6.0	14
137	Relationship between induced abortion and the incidence of depression, anxiety disorder, adjustment disorder, and somatoform disorder in Germany. <i>Journal of Psychiatric Research</i> , 2019, 114, 75-79.	3.1	14
138	Incidence of epilepsy and associated factors in elderly patients in Germany. <i>Epilepsy and Behavior</i> , 2019, 90, 107-111.	1.7	14
139	Trends in valproate use in patients in Germany between 2009 and 2017. <i>Epilepsy and Behavior</i> , 2019, 92, 26-30.	1.7	14
140	Non-alcoholic fatty liver disease (NAFLD) is associated with an increased incidence of osteoporosis and bone fractures. <i>Zeitschrift Fur Gastroenterologie</i> , 2021, , .	0.5	14
141	Overweight and Obesity Determine the Risk for Gastrointestinal Cancer in a Sex-Dependent Manner: A Retrospective Cohort Study of 287,357 Outpatients in Germany. <i>Cancers</i> , 2022, 14, 931.	3.7	14
142	Persistence with bisphosphonates in patients with metastatic breast cancer: a retrospective database analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013, 139, 1149-1155.	2.5	13
143	Age-related differences in persistence in women with breast cancer treated with tamoxifen or aromatase inhibitors in Germany. <i>Journal of Geriatric Oncology</i> , 2016, 7, 169-175.	1.0	13
144	Incidence of fractures in young women with breast cancer - a retrospective cohort study. <i>Journal of Bone Oncology</i> , 2019, 18, 100254.	2.4	13

#	ARTICLE	IF	CITATIONS
145	The Relationship Between the Use of Antihypertensive Drugs and the Incidence of Dementia in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019, 70, 91-97.	2.6	13
146	Association between various thyroid gland diseases, TSH values and thyroid cancer: a caseâ€“control study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 2989-2994.	2.5	13
147	Incidence of fractures in patients with type 1 diabetes mellitusâ€“a retrospective study with 4420 patients. <i>Osteoporosis International</i> , 2020, 31, 1315-1322.	3.1	13
148	Mild cognitive impairment is associated with fall-related injury among adults aged â‰¥65 years in low- and middle-income countries. <i>Experimental Gerontology</i> , 2021, 146, 111222.	2.8	13
149	Which adverse effects influence the dropout rate in selective serotonin reuptake inhibitor (SSRI) treatment? Results for 50,824 patients. <i>GMS German Medical Science</i> , 2014, 12, Doc15.	2.7	13
150	Persistence with antidepressant drugs in patients with dementia: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 323-329.	0.6	13
151	Persistence with antipsychotics in dementia patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 835-840.	0.6	13
152	Relation of the first hypertension-associated event with medication, compliance and persistence in naïve hypertensive patients after initiating monotherapy. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2010, 48, 173-183.	0.6	13
153	Early Discontinuation and Related Treatment Costs After Initiation of Basal Insulin in Type 2 Diabetes Patients. <i>Journal of Diabetes Science and Technology</i> , 2015, 9, 644-650.	2.2	12
154	Psoriasis risk in patients with type 2 diabetes in German primary care practices. <i>Primary Care Diabetes</i> , 2017, 11, 52-56.	1.8	12
155	Prescription Patterns and the Cost of Antihyperglycemic Drugs in Patients With Type 2 Diabetes Mellitus in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 123-127.	2.2	12
156	Demographic data and seasonal variation in peripheral vestibular disorders in ENT practices in Germany. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2019, 29, 181-190.	2.0	12
157	Cancer risk in stroke survivors followed for up to 10 years in general practices in Germany. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 1013-1020.	2.5	12
158	Factors associated with fracture after stroke and TIA: a long-term follow-up. <i>Osteoporosis International</i> , 2020, 31, 2395-2402.	3.1	12
159	Association between antiseizure medication use and risk of urinary tract infection: A caseâ€“control study. <i>Epilepsy and Behavior</i> , 2021, 115, 107502.	1.7	12
160	Association between sleep disorders and osteoarthritis: A caseâ€“control study of 351,932 adults in the UK. <i>Journal of Sleep Research</i> , 2021, 30, e13367.	3.2	12
161	Physical and mental health comorbidities of patients with epilepsy in Germany â€“ A retrospective cohort study. <i>Epilepsy and Behavior</i> , 2021, 117, 107857.	1.7	12
162	Risk factors for epistaxis in patients followed in general practices in Germany. <i>Rhinology</i> , 2017, 55, 312-318.	1.3	12

#	ARTICLE	IF	CITATIONS
163	Glycemic Durability of Monotherapy for Diabetes. <i>New England Journal of Medicine</i> , 2007, 356, 1378-1380.	27.0	11
164	Prevalence and treatment profile of patients with grass pollen and house dust mite allergy. <i>JDDG - Journal of the German Society of Dermatology</i> , 2013, 11, 653-661.	0.8	11
165	Emergency peripartum hysterectomy and risk factors by mode of delivery and obstetric history: a 10-year review from Helsinki University Central Hospital. <i>Journal of Perinatal Medicine</i> , 2015, 43, 721-8.	1.4	11
166	Increased treatment persistence and its determinants in women with osteoporosis with prior fracture compared to those without fracture. <i>Osteoporosis International</i> , 2016, 27, 963-969.	3.1	11
167	Risk Factors for Mild Cognitive Impairment in German Primary Care Practices. <i>Journal of Alzheimer's Disease</i> , 2017, 56, 379-384.	2.6	11
168	Increased depression symptom score in newly diagnosed type 2 diabetes patients. <i>Psychiatry Research</i> , 2018, 261, 259-263.	3.3	11
169	Association of characteristics of people with type 2 diabetes mellitus with discordant values of fasting glucose and HbA1c. <i>Journal of Diabetes</i> , 2018, 10, 934-941.	1.8	11
170	Risk Factors for Tinnitus in 37,692 Patients Followed in General Practices in Germany. <i>Otology and Neurotology</i> , 2019, 40, 436-440.	1.3	11
171	Achalasia is associated with a higher incidence of depression in outpatients in Germany. <i>PLoS ONE</i> , 2021, 16, e0250503.	2.5	11
172	Hepatobiliary Adverse Drug Reactions Associated With Remdesivir: The WHO International Pharmacovigilance Study. <i>Clinical Gastroenterology and Hepatology</i> , 2021, 19, 1970-1972.e3.	4.4	11
173	Inflammatory Bowel Disease as a Precondition for Stroke or TIA: A Matter of Crohn's Disease Rather than Ulcerative Colitis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021, 30, 105787.	1.6	11
174	Factors Associated with Non-Severe Adverse Reactions after Vaccination against SARS-CoV-2: A Cohort Study of 908,869 Outpatient Vaccinations in Germany. <i>Vaccines</i> , 2022, 10, 566.	4.4	11
175	Prevalence of diabetes mellitus among 80,193 gastrointestinal cancer patients in five European and three Asian countries. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 1057-1062.	2.5	11
176	Post-COVID-19 conditions in children and adolescents diagnosed with COVID-19. <i>Pediatric Research</i> , 2024, 95, 182-187.	2.3	11
177	Effects of selected antidiabetics on weight loss – A retrospective database analysis. <i>Primary Care Diabetes</i> , 2015, 9, 74-77.	1.8	10
178	Second-line therapy with biological drugs in rheumatoid arthritis patients in German rheumatologist practices: a retrospective database analysis. <i>Rheumatology International</i> , 2016, 36, 1113-1118.	3.0	10
179	Risk of stillbirth in pregnant women with obesity in the United Kingdom. <i>Obesity Research and Clinical Practice</i> , 2016, 10, 574-579.	1.8	10
180	Association between Anti-Dementia Treatment Persistence and Daily Dosage of the First Prescription: A Retrospective Analysis in Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 58, 37-44.	2.6	10

#	ARTICLE	IF	CITATIONS
181	Risk of depression, anxiety, and adjustment disorders in women with a suspected but unconfirmed diagnosis of breast or genital organ cancer in Germany. <i>Cancer Causes and Control</i> , 2017, 28, 1021-1026.	1.8	10
182	Impact of attention deficit hyperactivity disorder therapy on fracture risk in children treated in German pediatric practices. <i>Osteoporosis International</i> , 2017, 28, 1265-1269.	3.1	10
183	The impact of physician-level drug budgets on prescribing behavior. <i>European Journal of Health Economics</i> , 2018, 19, 213-222.	2.8	10
184	Low annual frequency of HbA _{1c} testing in people with Type 2 diabetes in primary care practices in Germany. <i>Diabetic Medicine</i> , 2018, 35, 249-254.	2.3	10
185	Proton pump inhibitor use is associated with a variety of infections in patients with liver cirrhosis. <i>Medicine (United States)</i> , 2020, 99, e23436.	1.0	10
186	Urinary and fecal incontinence in stroke survivors followed in general practice: A retrospective cohort study. <i>Annals of Physical and Rehabilitation Medicine</i> , 2020, 63, 488-494.	2.3	10
187	Osteoarthritis and the incidence of fracture in the United Kingdom: a retrospective cohort study of 258,696 patients. <i>Osteoarthritis and Cartilage</i> , 2021, 29, 215-221.	1.3	10
188	Cancer Patients Have an Increased Incidence of Dementia: A Retrospective Cohort Study of 185,736 Outpatients in Germany. <i>Cancers</i> , 2021, 13, 2027.	3.7	10
189	Prevalence of and factors associated with COVID-19 diagnosis in symptomatic patients followed in general practices in Germany between March 2020 and March 2021. <i>International Journal of Infectious Diseases</i> , 2021, 111, 37-42.	3.3	10
190	Treatment persistence evaluation of tamoxifen and aromatase inhibitors in breast cancer patients in early and late stage disease. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2014, 52, 933-939.	0.6	10
191	Risk of dementia in German patients treated with antidepressants in general or psychiatric practices. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 322-328.	0.6	10
192	Persistence and adherence to dutasteride/tamsulosin fixed-dose versus free-combination alpha blocker/5ARI therapy in patients with benign prostate hyperplasia in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 37-49.	0.6	10
193	Impact of caesarean section on mode of delivery, pregnancy-induced and pregnancy-associated disorders, and complications in the subsequent pregnancy in Germany. <i>GMS German Medical Science</i> , 2016, 14, Doc06.	2.7	10
194	Pre-diagnostic presentations of Multiple System Atrophy case control study in a primary care dataset. <i>Parkinsonism and Related Disorders</i> , 2022, 99, 101-104.	2.2	10
195	Prevalence of and Factors Associated With Post-Coronavirus Disease 2019 (COVID-19) Condition in the 12 Months After the Diagnosis of COVID-19 in Adults Followed in General Practices in Germany. <i>Open Forum Infectious Diseases</i> , 2022, 9, .	0.9	10
196	Prescription of Insulin Glargine in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007, 115, 252-256.	1.2	9
197	Risk of breast cancer in patients on long-acting insulin analogues in comparison with those on human insulin. <i>Diabetologia</i> , 2012, 55, 1554-1555.	6.3	9
198	Time to pregnancy in subfertile women in German gynecological practices: analysis of a representative cohort of more than 60,000 patients. <i>Archives of Gynecology and Obstetrics</i> , 2015, 291, 657-662.	1.7	9

#	ARTICLE	IF	CITATIONS
199	Depression risk in patients with heart failure in primary care practices in Germany. <i>International Psychogeriatrics</i> , 2016, 28, 1889-1894.	1.0	9
200	Caesarean section and its impact on fertility and time to a subsequent pregnancy in Germany: a database analysis in gynecological practices. <i>Archives of Gynecology and Obstetrics</i> , 2016, 294, 1005-1010.	1.7	9
201	Diabetes treatment in people with type 2 diabetes and schizophrenia: Retrospective primary care database analyses. <i>Primary Care Diabetes</i> , 2016, 10, 36-40.	1.8	9
202	Association between depression and persistence with oral antihyperglycemic drugs in type 2 diabetes mellitus patients in Germany. <i>Psychiatry Research</i> , 2018, 261, 90-93.	3.3	9
203	Changes in patient characteristics, glucose lowering treatment, glycemic control and complications in type 2 diabetes in general practices (Disease Analyzer, Germany: 2008-2016). <i>Postgraduate Medicine</i> , 2018, 130, 244-250.	2.0	9
204	20-year risk of breast cancer recurrence. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 765-766.	2.5	9
205	The Influence of Polypharmacy on the Initiation of Anti-Dementia Therapy in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 64, 827-833.	2.6	9
206	Real-world treatment patterns in patients with nontuberculous mycobacterial lung disease in general and pneumologist practices in Germany. <i>Journal of Clinical Tuberculosis and Other Mycobacterial Diseases</i> , 2020, 20, 100178.	1.3	9
207	Increased prevalence of depression, anxiety, and adjustment and somatoform disorders in transsexual individuals. <i>Journal of Affective Disorders</i> , 2020, 274, 482-485.	4.1	9
208	Osteoarthritis and incidence of stroke and transient ischemic attack in 320,136 adults followed in general practices in the United Kingdom. <i>Joint Bone Spine</i> , 2021, 88, 105104.	1.6	9
209	Anti-seizure medication is not associated with an increased risk to develop cancer in epilepsy patients. <i>Journal of Neurology</i> , 2021, 268, 2185-2191.	3.6	9
210	Trends in Dementia and Mild Cognitive Impairment Prevalence and Incidence in German General and Specialist Practices Between 2015 and 2019. <i>Journal of Alzheimer's Disease</i> , 2021, 79, 1683-1690.	2.6	9
211	Prevalence of and Factors Associated with Antibiotic Prescriptions in Patients with Acute Lower and Upper Respiratory Tract Infections – A Case-Control Study. <i>Antibiotics</i> , 2021, 10, 455.	3.7	9
212	Variables associated with increased incidence of non-alcoholic fatty liver disease (NAFLD) in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002243.	2.8	9
213	Impact of thyroid disorders on the incidence of non-alcoholic fatty liver disease in Germany. <i>United European Gastroenterology Journal</i> , 2021, 9, 829-836.	3.8	9
214	Clinical and Patient-Related Variables Associated with Initiating GLP-1 Receptor Agonist Therapy in Type 2 Diabetes Patients in Primary Care in Germany. <i>PLoS ONE</i> , 2016, 11, e0152281.	2.5	9
215	Patterns of benzodiazepine prescribing by neuropsychiatrists and general practitioners for elderly patients in Germany in 2014. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 466-471.	0.6	9
216	No significant association between COVID-19 diagnosis and the incidence of depression and anxiety disorder? A retrospective cohort study conducted in Germany. <i>Journal of Psychiatric Research</i> , 2022, 147, 79-84.	3.1	9

#	ARTICLE	IF	CITATIONS
217	Longitudinal change in HbA1c after insulin initiation in primary care patients with type 2 diabetes: A database analysis in UK and Germany. <i>Primary Care Diabetes</i> , 2012, 6, 47-52.	1.8	8
218	Unlicensed use of metformin in children and adolescents in Germany and France. <i>British Journal of Clinical Pharmacology</i> , 2012, 73, 307-308.	2.4	8
219	Topical therapy of hand eczema – analysis of the prescription profile from dermatologists in private practice. <i>JDDG - Journal of the German Society of Dermatology</i> , 2012, 10, 180-184.	0.8	8
220	The Prescribing of Contraceptives for Adolescents in German Gynecologic Practices in 2007 and 2011: A Retrospective Database Analysis. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2013, 26, 261-264.	0.7	8
221	Risk factors for discontinuation of insulin pump therapy in pediatric and young adult patients. <i>Primary Care Diabetes</i> , 2014, 8, 346-351.	1.8	8
222	Persistence with opioid treatment in Germany in patients suffering from chronic non-malignant or cancer pain. <i>Current Medical Research and Opinion</i> , 2015, 31, 1157-1163.	1.9	8
223	Persistence with antiepileptic drugs in epilepsy patients treated in neurological practices in Germany. <i>Epilepsy and Behavior</i> , 2017, 73, 204-207.	1.7	8
224	Prescription Patterns and the Cost of Migraine Treatments in German General and Neurological Practices. <i>Pain Practice</i> , 2017, 17, 747-752.	1.9	8
225	Increased risk for cancer after stroke at a young age: etiological relevance or incidental finding?. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 3047-3054.	2.5	8
226	Persistence with biological drugs in patients treated in rheumatology practices in Germany. <i>Rheumatology International</i> , 2019, 39, 525-531.	3.0	8
227	Long-term risk of stroke and its predictors in transient ischaemic attack patients in Germany. <i>European Journal of Neurology</i> , 2020, 27, 723-728.	3.3	8
228	Comparative Analysis of the 3-Year Persistence Rate with Second-Line Vedolizumab and Tumor Necrosis Factor- α Inhibitors in Patients with Inflammatory Bowel Disease Followed in Gastroenterology Practices in Germany. <i>Digestive Diseases</i> , 2020, 38, 466-473.	1.9	8
229	Sarcopenia and Mild Cognitive Impairment in Older Adults from Six Low- and Middle-Income Countries. <i>Journal of Alzheimer's Disease</i> , 2021, 82, 1745-1754.	2.6	8
230	Effects of the COVID-19 lockdown on primary health care for persons with type 2 diabetes – Results from the German Disease Analyzer database. <i>Diabetes Research and Clinical Practice</i> , 2021, 179, 109002.	2.8	8
231	Increased Incidence of Stroke and Transient Ischemic Attack in Patients with Rheumatoid Arthritis and Ankylosing Spondylitis in Germany. <i>Neuroepidemiology</i> , 2021, 55, 162-170.	2.3	8
232	A retrospective database analysis on persistence with inhaled corticosteroid therapy: comparison of two dry powder inhalers during asthma treatment in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 257-264.	0.6	8
233	Epilepsy is associated with an increased incidence of heart failure diagnoses. <i>Epilepsy and Behavior</i> , 2021, 125, 108393.	1.7	8
234	Risk factors for ectopic pregnancy in Germany: a retrospective study of 100,197 patients. <i>GMS German Medical Science</i> , 2017, 15, Doc19.	2.7	8

#	ARTICLE	IF	CITATIONS
235	The spectrum of comorbidities at the initial diagnosis of heart failure a case control study. Scientific Reports, 2022, 12, 2670.	3.3	8
236	Predictors for the initiation of a basal supported oral therapy (BOT) in type 2 diabetic patients under real-life conditions in Germany. Primary Care Diabetes, 2012, 6, 329-335.	1.8	7
237	Different Injection Frequencies of Basal Insulins in Type 2 Diabetes Patients under Real-Life Conditions: A Retrospective Database Analysis. Journal of Diabetes Science and Technology, 2013, 7, 1354-1358.	2.2	7
238	Prevalence of female subfertility in German gynecological practices. Gynecological Endocrinology, 2013, 29, 767-770.	1.7	7
239	Use of antidiabetic agents in the treatment of gestational diabetes mellitus in Germany, 2008-2012. Journal of Obstetrics and Gynaecology Research, 2014, 40, 1592-1597.	1.3	7
240	Persistency with estrogen replacement therapy among hysterectomized women after the Women's Health Initiative study. Climacteric, 2015, 18, 826-834.	2.4	7
241	Age-related differences in persistence with bisphosphonates in women with metastatic breast cancer. Journal of Bone Oncology, 2016, 5, 63-66.	2.4	7
242	Predictors of early discontinuation of basal insulin therapy in type 2 diabetes in primary care. Primary Care Diabetes, 2016, 10, 142-147.	1.8	7
243	The impact of non-persistence on the direct and indirect costs in patients treated with subcutaneous tumour necrosis factor-alpha inhibitors in Germany. Rheumatology, 2018, 57, 1276-1281.	1.9	7
244	Analysis of the Effects of Selective Serotonin (and Noradrenaline) Reuptake Inhibitors on the Risk of Dementia in Patients with Depression. Journal of Alzheimer's Disease, 2019, 69, 577-583.	2.6	7
245	Time to Insulin Initiation in Type 2 Diabetes Patients in 2010/2011 and 2016/2017 in Germany. Journal of Diabetes Science and Technology, 2019, 13, 1129-1134.	2.2	7
246	Influence of chemotherapy and endocrine treatment on fractures in postmenopausal women with breast cancer - a retrospective cohort study. Journal of Bone Oncology, 2020, 22, 100292.	2.4	7
247	Proton pump inhibitors increase risk of bone fractures in men with cirrhosis: a population-based study. Alimentary Pharmacology and Therapeutics, 2020, 52, 1042-1050.	3.7	7
248	Prescription of hormone replacement therapy prior to and after the diagnosis of gynecological cancers in German patients. Journal of Cancer Research and Clinical Oncology, 2020, 146, 1567-1573.	2.5	7
249	Burn injury and incidence of psychiatric disorders: A retrospective cohort study of 18,198 patients from Germany. Burns, 2021, 47, 1110-1117.	1.9	7
250	Impact of the COVID-19 Pandemic on Consultations and Diagnoses in Gastroenterology Practices in Germany. Frontiers in Medicine, 2021, 8, 684032.	2.6	7
251	Multimorbidity and Subjective Cognitive Complaints: Findings from 48 Low- and Middle-Income Countries of the World Health Survey 2002-2004. Journal of Alzheimer's Disease, 2021, 81, 1737-1747.	2.6	7
252	Association between type 2 diabetes and chronic low back pain in general practices in Germany. BMJ Open Diabetes Research and Care, 2021, 9, e002426.	2.8	7

#	ARTICLE	IF	CITATIONS
253	Pulmonary hypertension is associated with an increased incidence of NAFLD: A retrospective cohort study of 18,910 patients. <i>Journal of Internal Medicine</i> , 2021, 290, 886-893.	6.0	7
254	Comparative Analysis of 3-Year Persistence With Vedolizumab Compared With Antibodies Against Tumor Necrosis Factor-Alpha in Patients With Inflammatory Bowel Disease in Germany. <i>Journal of Clinical Gastroenterology</i> , 2021, 55, e1-e7.	2.2	7
255	Physicians' influence on breast cancer patient compliance. <i>GMS German Medical Science</i> , 2014, 12, Doc03.	2.7	7
256	Prevalence of depression, anxiety, adjustment disorders, and somatoform disorders in patients with age-related macular degeneration in Germany. <i>GMS German Medical Science</i> , 2017, 15, Doc04.	2.7	7
257	Regional differences in insulin therapy regimens in five European countries. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 403-408.	0.6	7
258	Prevalence and predictors of prescription of antibiotics in pregnant women treated by gynecologists in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 643-649.	0.6	7
259	Diagnosing and treating dementia in German primary and specialized care between 2011 and 2015. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 301-309.	0.6	7
260	Patterns of insulin therapy and insulin daily doses in children and adolescents with type 1 diabetes in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 296-301.	4.4	7
261	Overlap between irritable bowel syndrome and common gastrointestinal diagnoses: a retrospective cohort study of 29,553 outpatients in Germany. <i>BMC Gastroenterology</i> , 2022, 22, 48.	2.0	7
262	Influence of macro- and microvascular comorbidity on time to insulin initiation in type 2 diabetes patients: A retrospective database analysis in Germany, France, and UK. <i>Primary Care Diabetes</i> , 2013, 7, 167-171.	1.8	6
263	Predictors of Insulin Initiation in Metformin and Sulfonylurea Users in Primary Care Practices. <i>Journal of Diabetes Science and Technology</i> , 2014, 8, 1023-1028.	2.2	6
264	Macro- and Microvascular Outcomes in Patients with Type 2 Diabetes Treated with Rapid-Acting Insulin Analogues or Human Regular Insulin: a Retrospective Database Analysis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, 92-99.	1.2	6
265	Prevalence of high-risk cardiovascular patients with therapy-resistant hypercholesterolemia. <i>Cardiovascular Endocrinology</i> , 2017, 6, 81-85.	0.8	6
266	Treatment Outcomes and Tolerability Following Initiation of GLP-1 Receptor Agonists Among Type 2 Diabetes Patients in Primary Care Practices in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 272-277.	2.2	6
267	Seasonal Variation of Epistaxis in Germany. <i>Journal of Craniofacial Surgery</i> , 2018, 29, e365-e367.	0.7	6
268	Frequency of blood pressure and estimated glomerular filtration rate monitoring in patients affected by hypertension. <i>Blood Pressure Monitoring</i> , 2018, 23, 85-90.	0.8	6
269	Coded Prevalence of Mild Cognitive Impairment in General and Neuropsychiatrists Practices in Germany Between 2007 and 2017. <i>Journal of Alzheimer's Disease</i> , 2019, 67, 1313-1318.	2.6	6
270	Is There an Association Between Antiepileptic Drug Use and Dementia Risk? A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2019, 68, 97-103.	2.6	6

#	ARTICLE	IF	CITATIONS
271	Relationship between physical conditions and attempted or completed suicide in more than 9,300 individuals from the United Kingdom: a case-control study. <i>Journal of Affective Disorders</i> , 2020, 274, 457-463.	4.1	6
272	Prevalence of insulin as a first-line therapy and associated factors in people with type 2 diabetes in German primary care practices. <i>Diabetic Medicine</i> , 2020, 37, 1333-1339.	2.3	6
273	Impact of Non-Alcoholic Fatty Liver Disease on Metabolic Comorbidities in Type 2 Diabetes Mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, , .	1.2	6
274	Incidences of hypothyroidism and autoimmune thyroiditis are increased in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2021, 33, e1008-e1012.	1.6	6
275	Heart failure and depression: a comparative analysis with different types of cancer. <i>European Journal of Preventive Cardiology</i> , 2022, 29, e112-e114.	1.8	6
276	Multiple Sclerosis and Incidence of Urinary and Fecal Incontinence in Almost 9,000 Patients Followed Up for up to 10 Years in Germany. <i>Neuroepidemiology</i> , 2021, 55, 92-99.	2.3	6
277	Conflicts at work are associated with a higher risk of cardiovascular disease. <i>GMS German Medical Science</i> , 2017, 15, Doc08.	2.7	6
278	Impact of comorbidities on the treatment of atopic dermatitis in clinical practice. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2014, 52, 726-731.	0.6	6
279	Prevalence of switching from two anti-TNF biosimilars back to biologic reference products in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 323-328.	0.6	6
280	Use of a German longitudinal prescription database (LRx) in pharmacoepidemiology. <i>GMS German Medical Science</i> , 2015, 13, Doc14.	2.7	6
281	Effects of the COVID-19 pandemic on clinically diagnosed psychiatric disorders in persons with type 2 diabetes. <i>Diabetic Medicine</i> , 2022, 39, e14852.	2.3	6
282	Effects of a low-dose oral estrogen only treatment on bone mineral density and quantitative ultrasonometry in postmenopausal women. <i>Gynecological Endocrinology</i> , 2012, 28, 1002-1005.	1.7	5
283	Resource Consumption and Costs of Treatment in Patients with Type 1 Diabetes under Intensified Conventional Therapy under German Real-Life Conditions. <i>Journal of Diabetes Science and Technology</i> , 2013, 7, 736-742.	2.2	5
284	Which Adverse Effects Influence the Dropout Rate in Selective Serotonin Reuptake Inhibitor (Ssri) Treatment?. <i>Value in Health</i> , 2014, 17, A460.	0.3	5
285	Prevalence of Use of Cardiovascular Drugs in Dementia Patients Treated in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 56, 1519-1524.	2.6	5
286	Tramadol prescription patterns in patients followed by general practitioners and orthopedists in Germany in the year 2015. <i>Postgraduate Medicine</i> , 2018, 130, 37-41.	2.0	5
287	Prescription Patterns and Disease Control in Type 2 Diabetes Mellitus Patients in Nursing Home and Home Care Settings: A Retrospective Analysis in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018, 12, 136-139.	2.2	5
288	Persistence With Oral Antihyperglycemic Drugs in Type 2 Diabetes Mellitus Patients With Dementia in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018, 12, 140-144.	2.2	5

#	ARTICLE	IF	CITATIONS
289	Medication use in patients with chronic rhinosinusitis in Germany - a large retrospective patient-based study. <i>Rhinology</i> , 2018, 57, 94-100.	1.3	5
290	Diagnostic Behavior for Mild Cognitive Impairment in General and Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019, 68, 925-930.	2.6	5
291	Incidence and Risk of Various Disorders of the External Ear in Patients With Hearing Aids Treated in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2019, 40, 1076-1081.	1.3	5
292	Fecal and urinary incontinence are major problems associated with rectal cancer. <i>International Journal of Colorectal Disease</i> , 2020, 35, 35-40.	2.2	5
293	Association between skin disorders and depression in children and adolescents: A retrospective case-control study. <i>Journal of Affective Disorders</i> , 2021, 282, 939-944.	4.1	5
294	Nonalcoholic fatty liver disease is associated with a higher incidence of coeliac disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2022, 34, 328-331.	1.6	5
295	Clinical characteristics and sick leave associated with infectious mononucleosis in a real-world setting in Germany. <i>International Journal of Clinical Practice</i> , 2021, 75, e14690.	1.7	5
296	Predicting the risk of stroke in patients with late-onset epilepsy: A machine learning approach. <i>Epilepsy and Behavior</i> , 2021, 122, 108211.	1.7	5
297	Micro- and macrovascular outcomes in Type 2 diabetic patients treated with insulin glulisine or human regular insulin: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 821-829.	0.6	5
298	Prevalence and treatment of sexually transmitted infections in men followed by urologists in Germany - a cross sectional study with 347,090 men. <i>GMS German Medical Science</i> , 2018, 16, Doc03.	2.7	5
299	Prevalence of and factors associated with long-term sick leave in working-age adults with chronic low back pain in Germany. <i>International Archives of Occupational and Environmental Health</i> , 2022, 95, 1549-1556.	2.3	5
300	No increased incidence of venous thrombosis or pulmonary embolism after SARS-CoV-2 vaccination in Germany. <i>Public Health</i> , 2022, 207, 14-18.	2.9	5
301	Mild Cognitive Disorder in Post-COVID-19 Syndrome: A Retrospective Cohort Study of 67,000 Primary Care Post-COVID Patients. <i>Journal of Alzheimer's Disease Reports</i> , 2022, 6, 297-305.	2.2	5
302	PCV108 RISK AND COSTS OF THE FIRST HYPERTENSION-ASSOCIATED EVENT, COMPLIANCE AND PERSISTENCE IN NAÏVE HYPERTENSIVE PATIENTS AFTER INITIATING MONOTHERAPY. <i>Value in Health</i> , 2008, 11, A415.	0.3	4
303	Different persistence on initial basal supported oral therapy in Type 2 diabetics is associated with unequal distributions of insulin treatment regimens under real-life conditions in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2010, 48, 761-766.	0.6	4
304	Basal Supported Oral Therapy with Insulin Glargine Results in Longer Persistence and Lower Costs Compared with Insulin Detemir in Type 2 Diabetics in Germany. <i>Health Outcomes Research in Medicine</i> , 2011, 2, e39-e50.	0.6	4
305	Risk of diabetic foot ulceration during treatment with insulin glargine and NPH insulin. <i>Journal of Wound Care</i> , 2012, 21, 483-489.	1.2	4
306	Cancer is associated with intraoperative and postprocedural complications and disorders. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 777-781.	2.5	4

#	ARTICLE	IF	CITATIONS
307	Seasonality of Insulin Use in German Outpatients With Diabetes: A Retrospective Analysis. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 996-1000.	2.2	4
308	Changes in Glycemic Control and Body Weight After Initiation of Dapagliflozin or Basal Insulin Supported Oral Therapy in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 590-596.	2.2	4
309	Risk of Nonfatal Stroke in Type 2 Diabetes Mellitus Patients: A Retrospective Comparison Between Disease Management Programs and Standard Care. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 808-813.	2.2	4
310	Persistence with Antihypertensive Drugs in Patients with Hypertension and Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 60, 505-510.	2.6	4
311	Long-term use of basal insulin and the risk of breast cancer. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 763-764.	2.5	4
312	Current healthcare pathways in the treatment of rhinosinusitis in Germany. <i>Acta Oto-Laryngologica</i> , 2018, 138, 1086-1091.	0.9	4
313	Prescription-based prevalence of biological therapy in patients with psoriasis, rheumatoid arthritis, and inflammatory bowel diseases. <i>Biologicals</i> , 2019, 61, 52-54.	1.4	4
314	Duration of sick leave in patients with depression treated by general practitioners and psychiatrists. <i>Psychiatry Research</i> , 2019, 279, 382-383.	3.3	4
315	Prescription patterns of antiepileptic drugs in Kazakhstan in 2018: A retrospective study of 57,959 patients. <i>Epilepsy and Behavior</i> , 2019, 99, 106445.	1.7	4
316	Frequency of Blood Pressure and Estimated Glomerular Filtration Rate Testing in Type 2 Diabetes Mellitus: A Retrospective Study with 43,509 Patients. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2019, 127, 455-460.	1.2	4
317	Adherence to neuroleptic treatment in psychiatric practices: A retrospective study of 55 practices with more than 5000 bipolar and schizophrenic patients in Germany. <i>Psychiatry Research</i> , 2020, 284, 112758.	3.3	4
318	Association Between Migraine Headaches and Dry Eye Disease in Patients Studied in General Practices in Germany. <i>JAMA Ophthalmology</i> , 2020, 138, 223.	2.5	4
319	Incidence of and factors associated with manic episodes and bipolar disorder in the decade following depression onset in Germany. <i>Journal of Affective Disorders</i> , 2020, 266, 534-539.	4.1	4
320	Multimorbidity Among Adult Outpatients With Type 1 Diabetes in Germany. <i>Journal of Diabetes Science and Technology</i> , 2020, , 193229682096526.	2.2	4
321	Treatment Patterns of Patients with All-Cause Dementia in Russia. <i>Journal of Alzheimer's Disease Reports</i> , 2020, 4, 9-14.	2.2	4
322	Usage of antiepileptic drugs in different diseases in Germany â€“ A retrospective study. <i>Epilepsy and Behavior</i> , 2020, 111, 107210.	1.7	4
323	What have we learned in the past year? A study on pharmacy purchases of psychiatric drugs from wholesalers in the days prior to the first and second COVID-19 lockdowns in Germany. <i>Journal of Psychiatric Research</i> , 2021, 140, 346-349.	3.1	4
324	Suspected cancer diagnoses made by general practitioners in a population with subsequently confirmed cancer diagnoses in Germany: a retrospective study of 31,628 patients. <i>Oncotarget</i> , 2017, 8, 84540-84545.	1.8	4

#	ARTICLE	IF	CITATIONS
325	Treatment persistence in the use of basal insulins in Poland and Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 119-125.	0.6	4
326	Compliance with vaccination against hepatitis A virus in Germany: a retrospective analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 740-745.	0.6	4
327	Persistence with antihypertensive drugs in patients with depression in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 162-168.	0.6	4
328	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 147916412110338.	2.0	4
329	Association between ivy leaves dry extract EA 575 prescriptions and antibiotic use, sick leave duration, and repeated infections in adult patients. <i>Postgraduate Medicine</i> , 2022, 134, 333-340.	2.0	4
330	Atrial fibrillation is associated with a subsequent epilepsy diagnosis independent of stroke: A retrospective matched administrative cohort study on 149,632 patients. <i>Epilepsy and Behavior</i> , 2022, 132, 108721.	1.7	4
331	The role of private prescriptions in benzodiazepine and Z-drug use—a secondary analysis of office-based prescription data. <i>Deutsches Ärzteblatt International</i> , 0, , .	0.9	4
332	Sex-Differential Associations Between Body Mass Index and the Incidence of Dementia. <i>Journal of Alzheimer's Disease</i> , 2022, 88, 631-639.	2.6	4
333	Calculated Daily Insulin Dosages Overestimate Prescribed Insulin Doses in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 597-601.	2.2	3
334	Relevance of Coded Prodromal Mild Cognitive Impairment in the Routine Treatment of Patients with Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 393-399.	2.6	3
335	Diagnoses of suspected cancer in otolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018, 9, 459-463.	1.0	3
336	One-Year Persistence with Donepezil, Memantine, and Rivastigmine in More than 66,000 Elderly Patients Followed in Poland. <i>Journal of Alzheimer's Disease</i> , 2019, 70, 899-905.	2.6	3
337	Prevalence of valproate prescriptions in women of childbearing age in certain regions of Russia. <i>Epilepsy and Behavior</i> , 2019, 101, 106584.	1.7	3
338	Persistence with selective serotonin (norepinephrine) reuptake inhibitors in Germany—A retrospective database analysis. <i>Journal of Affective Disorders</i> , 2019, 247, 156-160.	4.1	3
339	Real-world prevalence and incidence of lower back pain in Germany. <i>Rheumatology International</i> , 2019, 39, 1113-1113.	3.0	3
340	Changes in the utilization of blood glucose test strips among patients using intermittent scanning continuous glucose monitoring in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 922-928.	4.4	3
341	Effect of progestogen-only contraception on premenopausal fracture risk: a case-control study. <i>Osteoporosis International</i> , 2020, 31, 1801-1806.	3.1	3
342	Incidence of Inner Ear Disorders in Various Forms of Acute Otitis Media in ENT Practices in Germany. <i>Ear, Nose and Throat Journal</i> , 2021, 100, 325S-332S.	0.8	3

#	ARTICLE	IF	CITATIONS
343	Heart failure and dementia: a comparative analysis with different types of cancer. <i>European Heart Journal Open</i> , 2021, 1, .	2.3	3
344	Risk of hypoglycaemia in type 2 diabetes patients under different insulin regimens: a primary care database analysis. <i>GMS German Medical Science</i> , 2015, 13, Doc01.	2.7	3
345	The negative impact of rebate contracts on the health care of patients with depression in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2011, 49, 397-402.	0.6	3
346	Frequency of hospitalizations prior to and after conversion to a rebate pharmaceutical in depression patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013, 51, 416-422.	0.6	3
347	Differences in baseline characteristics between type 2 diabetes mellitus patients treated with dipeptidyl peptidase-4 inhibitors in randomized controlled trials and those receiving the same treatment in real-world settings. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 411-416.	0.6	3
348	Multimorbidity and Anxiety Symptoms among Adults Aged 50 Years and Over from Six Low- and Middle-Income Countries. <i>Journal of Ageing and Longevity</i> , 2021, 1, 36-47.	0.6	3
349	Age-Related Changes in the Association Between Traumatic Brain Injury and Dementia in Older Men and Women. <i>Journal of Head Trauma Rehabilitation</i> , 2021, 36, E139-E146.	1.7	3
350	Prevalence of Lung Metastases among 19,321 Metastatic Colorectal Cancer Patients in Eight Countries of Europe and Asia. <i>Current Oncology</i> , 2021, 28, 5035-5040.	2.2	3
351	Association Between Ginkgo Biloba Extract Prescriptions and Dementia Incidence in Outpatients with Mild Cognitive Impairment in Germany: A Retrospective Cohort Study. <i>Journal of Alzheimer's Disease</i> , 2022, , 1-7.	2.6	3
352	An Elevated FIB-4 Score Is Associated with an Increased Incidence of Depression among Outpatients in Germany. <i>Journal of Clinical Medicine</i> , 2022, 11, 2214.	2.4	3
353	Antibiotic therapy is associated with an increased incidence of cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2023, 149, 1285-1293.	2.5	3
354	No effects of the COVID-19 pandemic on the prescription of insulin in Germany. <i>Primary Care Diabetes</i> , 2022, 16, 588-590.	1.8	3
355	Longitudinal Change in HbA1c in Patients with Type 2 Diabetes: Comparison of Short-acting, Biphasic and Basal Insulin. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2009, 117, 230-233.	1.2	2
356	PCV104 Persistence in Hypertension Treatment with Olmesartan Medoxomil Versus Valsartan - Analysis of Real-Life Prescription Data in Germany. <i>Value in Health</i> , 2011, 14, A383.	0.3	2
357	Risk of Psychiatric and Neurological Diseases in Patients with Workspace Mobbing Experience in Germany: A Retrospective Database Analysis. <i>Value in Health</i> , 2013, 16, A541.	0.3	2
358	Reduction In C-Reactive Protein With Biologic Drugs In Rheumatoid Arthritis and Spondylitis Patients In German Rheumatologist Practices. <i>Value in Health</i> , 2015, 18, A635.	0.3	2
359	Prevalence Of Depression And Anxiety And Their Risk Factors In Women With Breast Cancer In German Primary Care. <i>Value in Health</i> , 2015, 18, A431.	0.3	2
360	Impact of comorbidities on the cost of depression drug therapy in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2016, 83, 130-136.	3.1	2

#	ARTICLE	IF	CITATIONS
361	Fracture risk reduction with use of dipeptidyl peptidase-4 inhibitors: response to Driessen et al.. Osteoporosis International, 2017, 28, 2431-2431.	3.1	2
362	Association between gout and osteoporosis risk in the United Kingdom. Osteoporosis International, 2018, 29, 1685-1685.	3.1	2
363	Late Detection of Atrial Fibrillation after Stroke: Implications for the Secondary Prevention. European Neurology, 2019, 81, 262-269.	1.4	2
364	Longitudinal changes in HbA1c following therapy intensification after inadequate response to two oral antidiabetic agents in patients with type 2 diabetes in Germany. Diabetes, Obesity and Metabolism, 2019, 21, 1757-1757.	4.4	2
365	Sex Differences in People Aging With HIV in Germany. Journal of Acquired Immune Deficiency Syndromes (1999), 2020, 84, e11-e11.	2.1	2
366	Prevalence of urinary system, pelvic organ, and genital tract disorders among women with type 1 diabetes in Germany. Primary Care Diabetes, 2021, 15, 257-261.	1.8	2
367	Is there an association between multiple sclerosis and osteoarthritis in Germany? A retrospective cohort study of 8,600 patients from Germany. Multiple Sclerosis Journal - Experimental, Translational and Clinical, 2021, 7, 205521732110227.	1.0	2
368	Association between inflammatory central nervous system diseases and epilepsy: A retrospective cohort study of 4252 patients in Germany. Epilepsy and Behavior, 2021, 117, 107879.	1.7	2
369	Improved Balance and Gait Ability and Basic Activities of Daily Living after Comprehensive Geriatric Care in Frail Older Patients with Fractures. Healthcare (Switzerland), 2021, 9, 560.	2.0	2
370	Diverticular disease is associated with an increased incidence rate of depression and anxiety disorders. International Journal of Colorectal Disease, 2021, 36, 2437-2443.	2.2	2
371	Incidence of antiepileptic drug therapy and factors associated with their prescribing in outpatients with diabetic polyneuropathy. Primary Care Diabetes, 2021, 15, 535-540.	1.8	2
372	Association of objective and subjective far vision impairment with perceived stress among older adults in six low- and middle-income countries. Eye, 2022, 36, 1274-1280.	2.1	2
373	Relationship between chronic gingivitis and subsequent depression in 13,088 patients followed in general practices. Journal of Psychiatric Research, 2021, 138, 103-106.	3.1	2
374	Factors associated with the diagnosis of depression in women followed in gynecological practices in Germany. Journal of Psychiatric Research, 2021, 141, 358-363.	3.1	2
375	Bipolar Disorder and Outcomes of Monotherapy with Lithium, Valproate, Quetiapine, Olanzapine, Venlafaxine, and Citalopram. Pharmacopsychiatry, 2021, 54, 126-130.	3.3	2
376	Economic prescribing of corticosteroid nasal sprays in Germany. Comparison of mometasone and budesonide nasal sprays on the basis of the DDD, the PDD and reference prices. International Journal of Clinical Pharmacology and Therapeutics, 2013, 51, 12-18.	0.6	2
377	Gender differences in persistency to bisphosphonates in patients with metastatic breast and prostate cancer. International Journal of Clinical Pharmacology and Therapeutics, 2014, 52, 352-359.	0.6	2
378	Association of time-to-levodopa with initial Parkinsonian medication: a retrospective cohort study. International Journal of Clinical Pharmacology and Therapeutics, 2015, 53, 107-114.	0.6	2

#	ARTICLE	IF	CITATIONS
379	Age effects on treatment patterns in 138,097 patients with unipolar depression followed in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021, 144, 208-216.	3.1	2
380	Long-term use of proton pump inhibitors (PPIs) is associated with an increased risk of type 2 diabetes. <i>Gut</i> , 2021, , gutjnl-2021-326297.	12.1	2
381	Gender-based differences in the antidepressant treatment of patients with depression in German psychiatric practices. <i>GMS German Medical Science</i> , 2016, 14, Doc02.	2.7	2
382	Pulmonary hypertension is associated with an increased incidence of cancer diagnoses. <i>Pulmonary Circulation</i> , 0, , .	1.7	2
383	Incidence of Cancer in Patients with Irritable Bowl Syndrome. <i>Journal of Clinical Medicine</i> , 2021, 10, 5911.	2.4	2
384	An elevated FIB-4 score is not associated with cardiovascular events: a longitudinal analysis from 137 842 patients with and without chronic liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2022, Publish Ahead of Print, .	1.6	2
385	Incidence of other cancer diagnoses in women with breast cancer: a retrospective cohort study with 42,248 women. <i>Breast Cancer Research and Treatment</i> , 2022, 195, 75-82.	2.5	2
386	PHS10 HbA1c Testing Frequency in Primary Care Diabetes Patients in Germany and in the UK. <i>Value in Health</i> , 2012, 15, A519.	0.3	1
387	PCN106 Persistence in Patients With Breast Cancer Treated With Tamoxifen or Aromatase Inhibitors: Results of a Retrospective Cohort Study. <i>Value in Health</i> , 2012, 15, A428.	0.3	1
388	Secondary analysis of electronic databases: potentials and limitations. Reply to Asghari S, Mahdavian M [letter]. <i>Diabetologia</i> , 2013, 56, 2098-2099.	6.3	1
389	Persistence in Patients with Breast Cancer Treated with Tamoxifen or Aromatase Inhibitors- Analysis Based on Oncology Analyzer Database. <i>Value in Health</i> , 2013, 16, A418-A419.	0.3	1
390	Comparing Quality Effects Of Patient Care In Integrated And Regular Care For Patients With Hypertension. <i>Value in Health</i> , 2014, 17, A498.	0.3	1
391	Frequency of Thyroid-Stimulating Hormone Monitoring in Patients After Thyroidectomy: A Retrospective Study in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2018, 126, 640-644.	1.2	1
392	Time between suspected and confirmed diagnoses of Crohn's disease and ulcerative colitis in patients followed in gastroenterological practices in Germany. <i>International Journal of Colorectal Disease</i> , 2018, 33, 967-971.	2.2	1
393	P1930Dronedarone is associated with a lower risk of myocardial infarction in Germany - a retrospective database analysis. <i>European Heart Journal</i> , 2018, 39, .	2.2	1
394	Are data from general practices suitable for survival analyses in the field of breast cancer? A retrospective study conducted in the United Kingdom, France and Germany. <i>Molecular and Clinical Oncology</i> , 2019, 11, 177-180.	1.0	1
395	Persistence and treatment-free interval in patients being prescribed biological drugs in rheumatology practices in Germany. <i>European Journal of Clinical Pharmacology</i> , 2019, 75, 717-722.	1.9	1
396	Comment on Mizokami-Stout et al. The Contemporary Prevalence of Diabetic Neuropathy in Type 1 Diabetes: Findings From the T1D Exchange. <i>Diabetes Care</i> 2020;43:806-812. <i>Diabetes Care</i> , 2020, 43, e113-e113.	8.6	1

#	ARTICLE	IF	CITATIONS
397	Associations of physical and psychiatric conditions with chronic fatigue syndrome in Germany: an exploratory case-control study. <i>Psychological Medicine</i> , 2022, 52, 780-786.	4.5	1
398	Manufacturer switch of anti-seizure drugs may not increase the risk of seizure recurrence in Children: A nationwide study of prescription data in Germany. <i>Epilepsy and Behavior</i> , 2021, 115, 107705.	1.7	1
399	Prevalence of and reasons for referral of primary care outpatients to dentists in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 182-188.	0.6	1
400	Association between sexual orientation and subjective cognitive complaints in the general population in England. <i>Journal of Psychiatric Research</i> , 2021, 136, 1-6.	3.1	1
401	(Absence of) Association Between Non-Vitamin K Antagonist Oral Anticoagulant Therapy and Urinary Tract Infection in Patients With Atrial Fibrillation. <i>Journal of Cardiovascular Pharmacology</i> , 2021, 77, 830-834.	1.9	1
402	Prevalence of Hospitalization and Readmission in Patients with Inflammatory Bowel Diseases. <i>Digestive Diseases and Sciences</i> , 2022, 67, 3471-3471.	2.3	1
403	Increased pharmacy purchases of cardiovascular drugs from wholesalers prior to the first and second COVID-19 lockdowns. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 572-577.	0.6	1
404	Predicting the Risk of Ischemic Stroke in Patients Treated with Novel Oral Anticoagulants: A Machine Learning Approach. <i>Neuroepidemiology</i> , 2021, 55, 387-392.	2.3	1
405	Lines of therapy with biological drugs in dermatology, gastroenterology, and rheumatology practices in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 182-187.	0.6	1
406	Direct Oral Anticoagulants Versus Preprohormone in Patients With Cirrhosis and the Risk of Gastrointestinal Bleeding. <i>Annals of Pharmacotherapy</i> , 2021, , 106002802110480.	1.9	1
407	Versorgungsprofil von Patienten mit schweren allergischen Reaktionen. <i>Allergologie</i> , 2011, 34, 285-293.	0.1	1
408	Association of diabetes and cancer-diagnoses in primary care practices in Germany.. <i>Journal of Clinical Oncology</i> , 2014, 32, 1604-1604.	1.6	1
409	Treatment of depression in patients with cardiovascular diseases by German psychiatrists. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 557-563.	0.6	1
410	Prevalence and type of antidepressant therapy used by German general practitioners to treat female patients with osteoporosis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 743-749.	0.6	1
411	Prevalence of bisphosphonate therapy in women with breast cancer treated with aromatase inhibitors in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 394-396.	0.6	1
412	The role of general practitioners and psychiatrists in issuing initiation and follow-up prescriptions for selective serotonin (norepinephrine) reuptake inhibitors in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 329-333.	0.6	1
413	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 147916412111033819.	2.0	1
414	No association between antiviral treatment and risk of Alzheimer's disease in German outpatients. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2021, 7, e12216.	3.7	1

#	ARTICLE	IF	CITATIONS
415	Burden of depression and anxiety disorders per disease codes in patients with lymphoma in Germany. Supportive Care in Cancer, 2022, 30, 2387-2395.	2.2	1
416	Association between osteoarthritis and the incidence of Parkinson's disease in the United Kingdom. Clinical Parkinsonism & Related Disorders, 2021, 5, 100120.	0.9	1
417	Demographic and clinical characteristics of primary care patients vaccinated against COVID-19 in spring 2021. International Journal of Clinical Pharmacology and Therapeutics, 2022, , .	0.6	1
418	Persistence with insulin pump therapy among children and young adults with type 1 diabetes in Germany: An update. Diabetes, Obesity and Metabolism, 2022, 24, 948-950.	4.4	1
419	Frequency of over- and under-treatment with levothyroxine in primary care in Germany. European Journal of Endocrinology, 2022, 186, L5.	3.7	1
420	Portal Vein Thrombosis Is Associated with an Increased Incidence of Depression and Anxiety Disorders. Journal of Clinical Medicine, 2021, 10, 5689.	2.4	1
421	Portal vein thrombosis is associated with an increased risk of bone fractures. PLoS ONE, 2022, 17, e0267535.	2.5	1
422	Omega-3 fatty acids supplementation on major cardiovascular outcomes: an umbrella review of meta-analyses of observational studies and randomized controlled trials. European Review for Medical and Pharmacological Sciences, 2021, 25, 2079-2092.	0.7	1
423	Electroencephalography Findings in Older Adults Undergoing Geriatric Treatment: A Surrogate for the Outcome?. Brain Sciences, 2022, 12, 839.	2.3	1
424	PDB43 ADDING INSULIN GLARGINE TO ORAL THERAPY IN PATIENTS WITH TYPE 2 DIABETES RESULTS IN LONGER PERSISTENCE WITH TREATMENT COMPARED TO NPH INSULIN. Value in Health, 2008, 11, A509-A510.	0.3	0
425	PDB11 DATABASE ANALYSIS ON PREVALENCE AND RISK FACTORS OF DIABETIC FOOT SYNDROME (DFS) IN GERMANY. Value in Health, 2009, 12, A403.	0.3	0
426	PMS66 GRAND: THE GERMAN RETROSPECTIVE COHORT ANALYSIS ON NON-ADHERENCE AND ASSOCIATED RISK OF FRACTURES IN OSTEOPOROSIS PATIENTS TREATED WITH ORAL BIPHOSPHONATES. Value in Health, 2009, 12, A445.	0.3	0
427	PMS68 GRAND: THE GERMAN RETROSPECTIVE COHORT ANALYSIS ON NON-ADHERENCE IN OSTEOPOROSIS PATIENTS TREATED WITH ORAL BIPHOSPHONATES. Value in Health, 2009, 12, A446.	0.3	0
428	PDB10 PREVALENCE AND OUTCOME OF INSULIN ANALOGUE TREATMENT COMPARING GERMANY, UK, AND FRANCE. Value in Health, 2010, 13, A285.	0.3	0
429	PDB16 INSULIN GLARGINE IS ASSOCIATED WITH A LOWER INCIDENCE OF DIABETIC FOOT SYNDROME AND MACROVASCULAR COMPLICATIONS COMPARED TO NPH INSULIN IN TYPE 2 DIABETICS UNDER GERMAN REAL-LIFE CONDITIONS. Value in Health, 2010, 13, A286.	0.3	0
430	PDB55 COMPARISON OF RESOURCE USE AND COSTS IN TYPE 1 DIABETES PATIENTS TREATED WITH DIFFERENT LONG ACTING INSULINS IN A BASAL-BOLUS REGIMEN IN GERMANY. Value in Health, 2010, 13, A294.	0.3	0
431	PDB5 Micro- and Macrovascular Outcomes in Primary Care Patients with Type 2 Diabetes Treated with Insulin Glulisine or Human Regular Insulin: A Retrospective German Database Analysis. Value in Health, 2011, 14, A472.	0.3	0
432	Duration of first prescribed long acting insulin therapy in type 2 diabetes. Diabetes Research and Clinical Practice, 2012, 97, e3.	2.8	0

#	ARTICLE	IF	CITATIONS
433	PCN23 Cancer Related Bisphosphonate Therapy in Different Treatment Facilities. Value in Health, 2012, 15, A412.	0.3	0
434	Risk Factors for Discontinuation of Insulin Pump Therapy in Pediatric and Young Adult Patient Groups. Value in Health, 2013, 16, A444.	0.3	0
435	Pregnancy After Breast Cancer in 2000-2002 and 2010-2012: A Retrospective Database Analysis. Value in Health, 2013, 16, A393.	0.3	0
436	Prevalence of Chronic Diseases Among Older Patients (>65 Years) in German General Practitioner Practices. Value in Health, 2014, 17, A505.	0.3	0
437	Treatment Discontinuation In Patients With Urinary Incontinence Suffering From Glaucoma. Value in Health, 2014, 17, A471.	0.3	0
438	Therapy Of Depression In Cancer And Non-Cancer Patients In German Neuropsychiatric Practices. Value in Health, 2015, 18, A431.	0.3	0
439	High Risk of Depression In Patients with Heart Failure in Germany. Value in Health, 2016, 19, A636.	0.3	0
440	Treatment Outcomes Following Initiation of Dapagliflozin or Basal Supported Oral Therapy Among Type 2 Diabetes Patients In Primary Care Practices In Germany. Value in Health, 2016, 19, A666-A667.	0.3	0
441	Rheumatoid Arthritis is Associated with an Increased Risk of Depression in Elderly Patients in Germany. Value in Health, 2016, 19, A521.	0.3	0
442	Prevalence of Therapy-Resistant Hypercholesterolemia in High-Risk Cardiovascular Patients in Germany. Value in Health, 2016, 19, A659.	0.3	0
443	Cost of Antihyperglycemic Therapy In Type 2 Diabetes Patients Depending on HBA1C And Bmi Values. Value in Health, 2016, 19, A665.	0.3	0
444	Prescription Patterns And Costs Of Migraine Therapy In Germany. Value in Health, 2016, 19, A430.	0.3	0
445	Prescription Patterns and Drugs Costs in Patients with Dementia - A Retrospective Database Analysis. Value in Health, 2016, 19, A439.	0.3	0
446	Referral for occupational therapy after diagnosis of developmental disorder by German child psychiatrists. Early Child Development and Care, 2016, 186, 959-970.	1.3	0
447	Incidence of Psychiatric Disorders in Women WITH a Suspected But Unconfirmed Diagnosis of Breast or Genital Organ Cancer. Value in Health, 2017, 20, A709-A710.	0.3	0
448	Factors Associated with an Adherence with Antiepileptic Drugs in Children Treated in Pediatric Practices in Germany. Value in Health, 2017, 20, A402.	0.3	0
449	The Presentation Of Patients With Epistaxis Shows A Marked Seasonal Variation In Germany. Value in Health, 2017, 20, A602.	0.3	0
450	Persistence in Hypertension Therapy in Patients with and Without Depression in Germany. Value in Health, 2017, 20, A620.	0.3	0

#	ARTICLE	IF	CITATIONS
451	Spontaneous Abortion is Associated with an Increased Risk of Developing Psychiatric Diseases. Value in Health, 2017, 20, A716.	0.3	0
452	PIH35 - HTTPS://ISPOR.CONFEX.COM/ISPOR/EURO21/RESEARCH/PAPERS/INDEX.CGI?USERNAME=85228&PASSWORD=173196 Value in Health, 2018, 21, S219.	0.3	0
453	PMH16 - THE PRESCRIPTION PATTERNS OF SELECTIVE SEROTONIN (NOREPINEPHRINE) REUPTAKE INHIBITORS IN GERMANY. Value in Health, 2018, 21, S278.	0.3	0
454	PMH65 - PERSISTENCE WITH SELECTIVE SEROTONIN (NOREPINEPHRINE) REUPTAKE INHIBITOR THERAPY IN GERMANY. Value in Health, 2018, 21, S285.	0.3	0
455	PHP337 - OFF-LABEL PRESCRIPTION OF DRUGS COMPARED TO PRESCRIPTION AFTER LABEL EXTENSION IN CHILDREN FOR GERMANY. Value in Health, 2018, 21, S207-S208.	0.3	0
456	Discontinuation of Antihypertensive Therapy and Dementia Incidence in the United Kingdom. Journal of the American Medical Directors Association, 2019, 20, 790-791.	2.5	0
457	Epidemiology of Gynaecological and Breast Cancers. Advances in Medical Diagnosis, Treatment, and Care, 2021, , 1-21.	0.1	0
458	Correspondence. Deutsches Ärzteblatt International, 2021, 118, 271-272.	0.9	0
459	Prevalence and associated factors of referrals to hospitals in Germany: A retrospective study including 2,050,453 patients. International Journal of Clinical Pharmacology and Therapeutics, 2021, 59, 315-321.	0.6	0
460	Comment on Eilam et al. Incidence of acute ischemic cerebrovascular events in a cohort of HIV-infected patients. Acta Neurologica Belgica, 2021, , 1.	1.1	0
461	Effect of SARS-CoV-2 vaccination among health care workers in a geriatric care unit after a B.1.1.7-variant outbreak. Public Health, 2021, 198, e20-e22.	2.9	0
462	Prevalence of and factors associated with the prescription of phytopharmaceuticals in outpatients with urinary tract infections. International Journal of Clinical Pharmacology and Therapeutics, 2021, 59, 654-661.	0.6	0
463	Prevalence of and factors associated with the prescription of fibrates among patients receiving lipid-lowering drugs in Germany. Journal of Cardiovascular Pharmacology, 2021, Publish Ahead of Print, 885-890.	1.9	0
464	Importance of durational hormone therapy in breast cancer patients in Germany: Retrospective database analysis.. Journal of Clinical Oncology, 2014, 32, 1603-1603.	1.6	0
465	Benzodiazepine prescribing in the elderly in Germany and Korea: A comparison of two observational studies. International Journal of Clinical Pharmacology and Therapeutics, 2017, 55, 480-482.	0.6	0
466	Prevalence and treatment of sexually transmitted diseases in gynecological practices in Germany: A retrospective study with more than 1,000,000 patients. International Journal of Clinical Pharmacology and Therapeutics, 2018, 56, 212-216.	0.6	0
467	Persistence with biological drugs in psoriasis patients followed in dermatology practices in Germany: A retrospective cohort study of 1,201 patients. International Journal of Clinical Pharmacology and Therapeutics, 2020, 58, 74-81.	0.6	0
468	The role of the treating practice in persistence among dementia patients in Germany and the UK. International Journal of Clinical Pharmacology and Therapeutics, 2020, 58, 247-253.	0.6	0

#	ARTICLE	IF	CITATIONS
469	Retrospektive Analyse zum Einsatz kurzwirksamer Beta-2-Sympathomimetika (SABA) bei Patienten mit Asthma in Deutschland. , 2020, 74, .		0
470	Risk factors for epistaxis in patients followed in general practices in Germany. Rhinology, 2017, 55, 312-318.	1.3	0
471	Prescription rates of common medications in patients with decompensated cirrhosis in Germany. Zeitschrift Fur Gastroenterologie, 2021, , .	0.5	0
472	Factor V Leiden and the 10-year incidence of depression: A retrospective cohort study conducted in Germany. Journal of Psychiatric Research, 2022, 146, 87-91.	3.1	0
473	Overuse of short-acting beta-agonists (SABA) and risk factors among patients with asthma. , 2020, , .		0
474	Authors' reply: Pulmonary hypertension is associated with an increased incidence of NAFLD. Journal of Internal Medicine, 2022, 291, 527-527.	6.0	0
475	Autoimmune disorders are positively associated with achalasia. Neurogastroenterology and Motility, 2022, 34, e14327.	3.0	0
476	An elevated FIB-4 score is associated with an increased incidence of liver cancer: A longitudinal analysis among 248,224 outpatients in Germany. European Journal of Cancer, 2022, 168, 41-50.	2.8	0
477	Epidemiology of Gynaecological and Breast Cancers. , 2022, , 187-212.		0