

# Karel Kostev

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3412860/publications.pdf>

Version: 2024-02-01

477  
papers

6,652  
citations

109321

35  
h-index

149698

56  
g-index

506  
all docs

506  
docs citations

506  
times ranked

7779  
citing authors

#	ARTICLE	IF	CITATIONS
1	Post-COVID-19 conditions in children and adolescents diagnosed with COVID-19. <i>Pediatric Research</i> , 2024, 95, 182-187.	2.3	11
2	Increase in depression and anxiety disorder diagnoses during the COVID-19 pandemic in children and adolescents followed in pediatric practices in Germany. <i>European Child and Adolescent Psychiatry</i> , 2023, 32, 873-879.	4.7	16
3	Antibiotic therapy is associated with an increased incidence of cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2023, 149, 1285-1293.	2.5	3
4	Associations of physical and psychiatric conditions with chronic fatigue syndrome in Germany: an exploratory case-control study. <i>Psychological Medicine</i> , 2022, 52, 780-786.	4.5	1
5	Heart failure and depression: a comparative analysis with different types of cancer. <i>European Journal of Preventive Cardiology</i> , 2022, 29, e112-e114.	1.8	6
6	Association of objective and subjective far vision impairment with perceived stress among older adults in six low- and middle-income countries. <i>Eye</i> , 2022, 36, 1274-1280.	2.1	2
7	Nonalcoholic fatty liver disease is associated with a higher incidence of coeliac disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2022, 34, 328-331.	1.6	5
8	Prevalence of Hospitalization and Readmission in Patients with Inflammatory Bowel Diseases. <i>Digestive Diseases and Sciences</i> , 2022, 67, 3471-3471.	2.3	1
9	An elevated FIB-4 score predicts liver cancer development: A longitudinal analysis from 29,999 patients with NAFLD. <i>Journal of Hepatology</i> , 2022, 76, 247-248.	3.7	25
10	Fewer non- $\text{COVID-19}$ respiratory tract infections and gastrointestinal infections during the $\text{COVID-19}$ pandemic. <i>Journal of Medical Virology</i> , 2022, 94, 298-302.	5.0	32
11	Patterns of insulin therapy and insulin daily doses in children and adolescents with type 1 diabetes in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 296-301.	4.4	7
12	Burden of depression and anxiety disorders per disease codes in patients with lymphoma in Germany. <i>Supportive Care in Cancer</i> , 2022, 30, 2387-2395.	2.2	1
13	Demographic and clinical characteristics of primary care patients vaccinated against COVID-19 in spring 2021. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2022, , .	0.6	1
14	Factor V Leiden and the 10-year incidence of depression: A retrospective cohort study conducted in Germany. <i>Journal of Psychiatric Research</i> , 2022, 146, 87-91.	3.1	0
15	No significant association between COVID-19 diagnosis and the incidence of depression and anxiety disorder? A retrospective cohort study conducted in Germany. <i>Journal of Psychiatric Research</i> , 2022, 147, 79-84.	3.1	9
16	Decrease in the number of patients diagnosed with cancer during the COVID-19 pandemic in Germany. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 3117-3123.	2.5	15
17	Authors' reply: Pulmonary hypertension is associated with an increased incidence of NAFLD. <i>Journal of Internal Medicine</i> , 2022, 291, 527-527.	6.0	0
18	Association between ivy leaves dry extract EA 575 prescriptions and antibiotic use, sick leave duration, and repeated infections in adult patients. <i>Postgraduate Medicine</i> , 2022, 134, 333-340.	2.0	4

#	ARTICLE	IF	CITATIONS
19	Persistence with insulin pump therapy among children and young adults with type 1 diabetes in Germany: An update. <i>Diabetes, Obesity and Metabolism</i> , 2022, 24, 948-950.	4.4	1
20	Investigation of the prevalence of non-COVID-19 infectious diseases during the COVID-19 pandemic. <i>Public Health</i> , 2022, 203, 53-57.	2.9	17
21	Overlap between irritable bowel syndrome and common gastrointestinal diagnoses: a retrospective cohort study of 29,553 outpatients in Germany. <i>BMC Gastroenterology</i> , 2022, 22, 48.	2.0	7
22	Autoimmune disorders are positively associated with achalasia. <i>Neurogastroenterology and Motility</i> , 2022, 34, e14327.	3.0	0
23	Association Between Ginkgo Biloba Extract Prescriptions and Dementia Incidence in Outpatients with Mild Cognitive Impairment in Germany: A Retrospective Cohort Study. <i>Journal of Alzheimer's Disease</i> , 2022, , 1-7.	2.6	3
24	Frequency of over- and under-treatment with levothyroxine in primary care in Germany. <i>European Journal of Endocrinology</i> , 2022, 186, L5.	3.7	1
25	Overweight and Obesity Determine the Risk for Gastrointestinal Cancer in a Sex-Dependent Manner: A Retrospective Cohort Study of 287,357 Outpatients in Germany. <i>Cancers</i> , 2022, 14, 931.	3.7	14
26	Prevalence of and factors associated with long-term sick leave in working-age adults with chronic low back pain in Germany. <i>International Archives of Occupational and Environmental Health</i> , 2022, 95, 1549-1556.	2.3	5
27	The spectrum of comorbidities at the initial diagnosis of heart failure a case control study. <i>Scientific Reports</i> , 2022, 12, 2670.	3.3	8
28	Pre-diagnostic presentations of Multiple System Atrophy case control study in a primary care dataset. <i>Parkinsonism and Related Disorders</i> , 2022, 99, 101-104.	2.2	10
29	No increased incidence of venous thrombosis or pulmonary embolism after SARS-CoV-2 vaccination in Germany. <i>Public Health</i> , 2022, 207, 14-18.	2.9	5
30	Incidence of newly diagnosed diabetes after Covid-19. <i>Diabetologia</i> , 2022, 65, 949-954.	6.3	92
31	Obesity and lipid metabolism disorders determine the risk for development of long COVID syndrome: a cross-sectional study from 50,402 COVID-19 patients. <i>Infection</i> , 2022, 50, 1165-1170.	4.7	31
32	Factors Associated with Non-Severe Adverse Reactions after Vaccination against SARS-CoV-2: A Cohort Study of 908,869 Outpatient Vaccinations in Germany. <i>Vaccines</i> , 2022, 10, 566.	4.4	11
33	Prevalence of diabetes mellitus among 80,193 gastrointestinal cancer patients in five European and three Asian countries. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 1057-1062.	2.5	11
34	Low blood levels of high-density lipoprotein (HDL) cholesterol are positively associated with cancer. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022, 148, 3039-3046.	2.5	16
35	An Elevated FIB-4 Score Is Associated with an Increased Incidence of Depression among Outpatients in Germany. <i>Journal of Clinical Medicine</i> , 2022, 11, 2214.	2.4	3
36	An elevated FIB-4 score is not associated with cardiovascular events: a longitudinal analysis from 137 842 patients with and without chronic liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2022, Publish Ahead of Print, .	1.6	2

#	ARTICLE	IF	CITATIONS
37	Effects of the COVID-19 pandemic on clinically diagnosed psychiatric disorders in persons with type 2 diabetes. <i>Diabetic Medicine</i> , 2022, 39, e14852.	2.3	6
38	An elevated FIB-4 score is associated with an increased incidence of liver cancer: A longitudinal analysis among 248,224 outpatients in Germany. <i>European Journal of Cancer</i> , 2022, 168, 41-50.	2.8	0
39	Portal vein thrombosis is associated with an increased risk of bone fractures. <i>PLoS ONE</i> , 2022, 17, e0267535.	2.5	1
40	No effects of the COVID-19 pandemic on the prescription of insulin in Germany. <i>Primary Care Diabetes</i> , 2022, 16, 588-590.	1.8	3
41	Atrial fibrillation is associated with a subsequent epilepsy diagnosis independent of stroke: A retrospective matched administrative cohort study on 149,632 patients. <i>Epilepsy and Behavior</i> , 2022, 132, 108721.	1.7	4
42	Epidemiology of Gynaecological and Breast Cancers. , 2022, , 187-212.		0
43	Sex-Differential Associations Between Body Mass Index and the Incidence of Dementia. <i>Journal of Alzheimer's Disease</i> , 2022, 88, 631-639.	2.6	4
44	Mild Cognitive Disorder in Post-COVID-19 Syndrome: A Retrospective Cohort Study of 67,000 Primary Care Post-COVID Patients. <i>Journal of Alzheimer's Disease Reports</i> , 2022, 6, 297-305.	2.2	5
45	Electroencephalography Findings in Older Adults Undergoing Geriatric Treatment: A Surrogate for the Outcome?. <i>Brain Sciences</i> , 2022, 12, 839.	2.3	1
46	Incidence of other cancer diagnoses in women with breast cancer: a retrospective cohort study with 42,248 women. <i>Breast Cancer Research and Treatment</i> , 2022, 195, 75-82.	2.5	2
47	Prevalence of and Factors Associated With Post-Coronavirus Disease 2019 (COVID-19) Condition in the 12 Months After the Diagnosis of COVID-19 in Adults Followed in General Practices in Germany. <i>Open Forum Infectious Diseases</i> , 2022, 9, .	0.9	10
48	Incident Dementia in Elderly Patients with Nonalcoholic Fatty Liver Disease in Germany. <i>Digestive Diseases and Sciences</i> , 2021, 66, 3179-3185.	2.3	32
49	Depression and sleep disorders are associated with early mortality in women with breast cancer in the United Kingdom. <i>Journal of Psychiatric Research</i> , 2021, 143, 481-484.	3.1	18
50	Impact of the coronavirus 2019 (COVID-19) pandemic on anxiety diagnosis in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021, 143, 528-533.	3.1	32
51	Burn injury and incidence of psychiatric disorders: A retrospective cohort study of 18,198 patients from Germany. <i>Burns</i> , 2021, 47, 1110-1117.	1.9	7
52	Incidence of Inner Ear Disorders in Various Forms of Acute Otitis Media in ENT Practices in Germany. <i>Ear, Nose and Throat Journal</i> , 2021, 100, 325S-332S.	0.8	3
53	Prevalence of urinary system, pelvic organ, and genital tract disorders among women with type 1 diabetes in Germany. <i>Primary Care Diabetes</i> , 2021, 15, 257-261.	1.8	2
54	Association Between Low-Density Lipoprotein Cholesterol Levels, Statin Use, and Dementia in Patients followed in German General Practices. <i>Journal of Alzheimer's Disease</i> , 2021, 79, 37-46.	2.6	19

#	ARTICLE	IF	CITATIONS
55	Effect of the COVID-19 lockdown on disease recognition and utilisation of healthcare services in the older population in Germany: a cross-sectional study. <i>Age and Ageing</i> , 2021, 50, 317-325.	1.6	71
56	Association between antiseizure medication use and risk of urinary tract infection: A caseâ€“control study. <i>Epilepsy and Behavior</i> , 2021, 115, 107502.	1.7	12
57	Osteoarthritis and incidence of stroke and transient ischemic attack in 320,136 adults followed in general practices in the United Kingdom. <i>Joint Bone Spine</i> , 2021, 88, 105104.	1.6	9
58	Change in glucoseâ€“lowering medication regimens in individuals with type 2 diabetes mellitus during the COVIDâ€“19 pandemic in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 910-915.	4.4	15
59	Anti-seizure medication is not associated with an increased risk to develop cancer in epilepsy patients. <i>Journal of Neurology</i> , 2021, 268, 2185-2191.	3.6	9
60	Epidemiology of Gynaecological and Breast Cancers. <i>Advances in Medical Diagnosis, Treatment, and Care</i> , 2021, , 1-21.	0.1	0
61	Manufacturer switch of anti-seizure drugs may not increase the risk of seizure recurrence in Children: A nationwide study of prescription data in Germany. <i>Epilepsy and Behavior</i> , 2021, 115, 107705.	1.7	1
62	Consultations Decline for Stroke, Transient Ischemic Attack, and Myocardial Infarction during the COVID-19 Pandemic in Germany. <i>Neuroepidemiology</i> , 2021, 55, 70-78.	2.3	18
63	Impact of Non-Alcoholic Fatty Liver Disease on Metabolic Comorbidities in Type 2 Diabetes Mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021, , .	1.2	6
64	Trends in Dementia and Mild Cognitive Impairment Prevalence and Incidence in German General and Specialist Practices Between 2015 and 2019. <i>Journal of Alzheimer's Disease</i> , 2021, 79, 1683-1690.	2.6	9
65	Osteoarthritis and the incidence of fracture in the United Kingdom: a retrospective cohort study of 258,696 patients. <i>Osteoarthritis and Cartilage</i> , 2021, 29, 215-221.	1.3	10
66	Prevalence of and reasons for referral of primary care outpatients to dentists in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 182-188.	0.6	1
67	Is loneliness associated with mild cognitive impairment in lowâ€“and middleâ€“income countries?. <i>International Journal of Geriatric Psychiatry</i> , 2021, 36, 1345-1353.	2.7	17
68	Association between skin disorders and depression in children and adolescents: A retrospective case-control study. <i>Journal of Affective Disorders</i> , 2021, 282, 939-944.	4.1	5
69	Antihypertensive medication and incident dementia: Observational cohort studies in four countries. <i>European Journal of Neurology</i> , 2021, 28, 1840-1848.	3.3	26
70	Association between sexual orientation and subjective cognitive complaints in the general population in England. <i>Journal of Psychiatric Research</i> , 2021, 136, 1-6.	3.1	1
71	Is there an association between multiple sclerosis and osteoarthritis in Germany? A retrospective cohort study of 8,600 patients from Germany. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2021, 7, 205521732110227.	1.0	2
72	Association Between Inflammatory Bowel Disease and Dementia: A Retrospective Cohort Study. <i>Journal of Alzheimer's Disease</i> , 2021, 80, 1471-1478.	2.6	26

#	ARTICLE	IF	CITATIONS
73	Association between inflammatory central nervous system diseases and epilepsy: A retrospective cohort study of 4252 patients in Germany. <i>Epilepsy and Behavior</i> , 2021, 117, 107879.	1.7	2
74	Prevalence of overuse of short-acting beta-2 agonists (SABA) and associated factors among patients with asthma in Germany. <i>Respiratory Research</i> , 2021, 22, 108.	3.6	22
75	Achalasia is associated with a higher incidence of depression in outpatients in Germany. <i>PLoS ONE</i> , 2021, 16, e0250503.	2.5	11
76	Incidences of hypothyroidism and autoimmune thyroiditis are increased in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2021, 33, e1008-e1012.	1.6	6
77	Mild cognitive impairment is associated with fall-related injury among adults aged ≥65 years in low- and middle-income countries. <i>Experimental Gerontology</i> , 2021, 146, 111222.	2.8	13
78	Prevalence of and Factors Associated with Antibiotic Prescriptions in Patients with Acute Lower and Upper Respiratory Tract Infections—A Case-Control Study. <i>Antibiotics</i> , 2021, 10, 455.	3.7	9
79	The impact of the coronavirus disease (COVID-19) pandemic on outpatient epilepsy care: An analysis of physician practices in Germany. <i>Epilepsy and Behavior</i> , 2021, 117, 107833.	1.7	16
80	Correspondence. <i>Deutsches Ärzteblatt International</i> , 2021, 118, 271-272.	0.9	0
81	Cancer Patients Have an Increased Incidence of Dementia: A Retrospective Cohort Study of 185,736 Outpatients in Germany. <i>Cancers</i> , 2021, 13, 2027.	3.7	10
82	Association between sleep disorders and osteoarthritis: A case-control study of 351,932 adults in the UK. <i>Journal of Sleep Research</i> , 2021, 30, e13367.	3.2	12
83	Prevalence and associated factors of referrals to hospitals in Germany: A retrospective study including 2,050,453 patients. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 315-321.	0.6	0
84	Hepatobiliary Adverse Drug Reactions Associated With Remdesivir: The WHO International Pharmacovigilance Study. <i>Clinical Gastroenterology and Hepatology</i> , 2021, 19, 1970-1972.e3.	4.4	11
85	Physical and mental health comorbidities of patients with epilepsy in Germany—A retrospective cohort study. <i>Epilepsy and Behavior</i> , 2021, 117, 107857.	1.7	12
86	Improved Balance and Gait Ability and Basic Activities of Daily Living after Comprehensive Geriatric Care in Frail Older Patients with Fractures. <i>Healthcare (Switzerland)</i> , 2021, 9, 560.	2.0	2
87	Variables associated with increased incidence of non-alcoholic fatty liver disease (NAFLD) in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002243.	2.8	9
88	Impact of the COVID-19 Pandemic on Consultations and Diagnoses in Gastroenterology Practices in Germany. <i>Frontiers in Medicine</i> , 2021, 8, 684032.	2.6	7
89	Diverticular disease is associated with an increased incidence rate of depression and anxiety disorders. <i>International Journal of Colorectal Disease</i> , 2021, 36, 2437-2443.	2.2	2
90	Incidence of antiepileptic drug therapy and factors associated with their prescribing in outpatients with diabetic polyneuropathy. <i>Primary Care Diabetes</i> , 2021, 15, 535-540.	1.8	2

#	ARTICLE	IF	CITATIONS
91	Comment on Eilam et al. Incidence of acute ischemic cerebrovascular events in a cohort of HIV-infected patients. <i>Acta Neurologica Belgica</i> , 2021, , 1.	1.1	0
92	(Absence of) Association Between Non-Vitamin K Antagonist Oral Anticoagulant Therapy and Urinary Tract Infection in Patients With Atrial Fibrillation. <i>Journal of Cardiovascular Pharmacology</i> , 2021, 77, 830-834.	1.9	1
93	Multimorbidity and Subjective Cognitive Complaints: Findings from 48 Low- and Middle-Income Countries of the World Health Survey 2002-2004. <i>Journal of Alzheimer's Disease</i> , 2021, 81, 1737-1747.	2.6	7
94	Heart failure is associated with an increased incidence of cancer diagnoses. <i>ESC Heart Failure</i> , 2021, 8, 3628-3633.	3.1	31
95	Treatment of Urinary Tract Infections with Canephron® in Germany: A Retrospective Database Analysis. <i>Antibiotics</i> , 2021, 10, 685.	3.7	16
96	Relationship between chronic gingivitis and subsequent depression in 13,088 patients followed in general practices. <i>Journal of Psychiatric Research</i> , 2021, 138, 103-106.	3.1	2
97	Inflammatory Bowel Disease as a Precondition for Stroke or TIA: A Matter of Crohn's Disease Rather than Ulcerative Colitis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021, 30, 105787.	1.6	11
98	Impact of thyroid disorders on the incidence of non-alcoholic fatty liver disease in Germany. <i>United European Gastroenterology Journal</i> , 2021, 9, 829-836.	3.8	9
99	Heart failure and dementia: a comparative analysis with different types of cancer. <i>European Heart Journal Open</i> , 2021, 1, .	2.3	3
100	Association between type 2 diabetes and chronic low back pain in general practices in Germany. <i>BMJ Open Diabetes Research and Care</i> , 2021, 9, e002426.	2.8	7
101	Increased pharmacy purchases of cardiovascular drugs from wholesalers prior to the first and second COVID-19 lockdowns. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 572-577.	0.6	1
102	Prevalence of, and factors associated with, long-term COVID-19 sick leave in working-age patients followed in general practices in Germany. <i>International Journal of Infectious Diseases</i> , 2021, 109, 203-208.	3.3	21
103	What have we learned in the past year? A study on pharmacy purchases of psychiatric drugs from wholesalers in the days prior to the first and second COVID-19 lockdowns in Germany. <i>Journal of Psychiatric Research</i> , 2021, 140, 346-349.	3.1	4
104	Clinical characteristics and sick leave associated with infectious mononucleosis in a real-world setting in Germany. <i>International Journal of Clinical Practice</i> , 2021, 75, e14690.	1.7	5
105	Sarcopenia and Mild Cognitive Impairment in Older Adults from Six Low- and Middle-Income Countries. <i>Journal of Alzheimer's Disease</i> , 2021, 82, 1745-1754.	2.6	8
106	Effects of the COVID-19 lockdown on primary health care for persons with type 2 diabetes - Results from the German Disease Analyzer database. <i>Diabetes Research and Clinical Practice</i> , 2021, 179, 109002.	2.8	8
107	Pulmonary hypertension is associated with an increased incidence of NAFLD: A retrospective cohort study of 18,910 patients. <i>Journal of Internal Medicine</i> , 2021, 290, 886-893.	6.0	7
108	Effect of SARS-CoV-2 vaccination among health care workers in a geriatric care unit after a B.1.1.7-variant outbreak. <i>Public Health</i> , 2021, 198, e20-e22.	2.9	0

#	ARTICLE	IF	CITATIONS
109	Predicting the risk of stroke in patients with late-onset epilepsy: A machine learning approach. <i>Epilepsy and Behavior</i> , 2021, 122, 108211.	1.7	5
110	Factors associated with the diagnosis of depression in women followed in gynecological practices in Germany. <i>Journal of Psychiatric Research</i> , 2021, 141, 358-363.	3.1	2
111	Prevalence of and factors associated with the prescription of phytopharmaceuticals in outpatients with urinary tract infections. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2021, 59, 654-661.	0.6	0
112	Prevalence of and factors associated with COVID-19 diagnosis in symptomatic patients followed in general practices in Germany between March 2020 and March 2021. <i>International Journal of Infectious Diseases</i> , 2021, 111, 37-42.	3.3	10
113	Effect of the COVID-19 pandemic on stress, anxiety, and depressive disorders in German primary care: A cross-sectional study. <i>Journal of Psychiatric Research</i> , 2021, 143, 43-49.	3.1	21
114	Bipolar Disorder and Outcomes of Monotherapy with Lithium, Valproate, Quetiapine, Olanzapine, Venlafaxine, and Citalopram. <i>Pharmacopsychiatry</i> , 2021, 54, 126-130.	3.3	2
115	Increased Incidence of Stroke and Transient Ischemic Attack in Patients with Rheumatoid Arthritis and Ankylosing Spondylitis in Germany. <i>Neuroepidemiology</i> , 2021, 55, 162-170.	2.3	8
116	Multiple Sclerosis and Incidence of Urinary and Fecal Incontinence in Almost 9,000 Patients Followed Up for up to 10 Years in Germany. <i>Neuroepidemiology</i> , 2021, 55, 92-99.	2.3	6
117	Predicting the Risk of Ischemic Stroke in Patients Treated with Novel Oral Anticoagulants: A Machine Learning Approach. <i>Neuroepidemiology</i> , 2021, 55, 387-392.	2.3	1
118	Impact of the COVID-19 Pandemic on Cancer Diagnoses in General and Specialized Practices in Germany. <i>Cancers</i> , 2021, 13, 408.	3.7	90
119	Comparative Analysis of 3-Year Persistence With Vedolizumab Compared With Antibodies Against Tumor Necrosis Factor-Alpha in Patients With Inflammatory Bowel Disease in Germany. <i>Journal of Clinical Gastroenterology</i> , 2021, 55, e1-e7.	2.2	7
120	Multimorbidity and Anxiety Symptoms among Adults Aged 50 Years and Over from Six Low- and Middle-Income Countries. <i>Journal of Ageing and Longevity</i> , 2021, 1, 36-47.	0.6	3
121	Direct Oral Anticoagulants Versus Preprohormone in Patients With Cirrhosis and the Risk of Gastrointestinal Bleeding. <i>Annals of Pharmacotherapy</i> , 2021, , 106002802110480.	1.9	1
122	Prevalence of and factors associated with the prescription of fibrates among patients receiving lipid-lowering drugs in Germany. <i>Journal of Cardiovascular Pharmacology</i> , 2021, Publish Ahead of Print, 885-890.	1.9	0
123	Age effects on treatment patterns in 138,097 patients with unipolar depression followed in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021, 144, 208-216.	3.1	2
124	Non-alcoholic fatty liver disease (NAFLD) is associated with an increased incidence of osteoporosis and bone fractures. <i>Zeitschrift Fur Gastroenterologie</i> , 2021, , .	0.5	14
125	Long-term use of proton pump inhibitors (PPIs) is associated with an increased risk of type 2 diabetes. <i>Gut</i> , 2021, , gutjnl-2021-326297.	12.1	2
126	Epilepsy is associated with an increased incidence of heart failure diagnoses. <i>Epilepsy and Behavior</i> , 2021, 125, 108393.	1.7	8



#	ARTICLE	IF	CITATIONS
127	Age-Related Changes in the Association Between Traumatic Brain Injury and Dementia in Older Men and Women. <i>Journal of Head Trauma Rehabilitation</i> , 2021, 36, E139-E146.	1.7	3
128	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 14791641211033819.	2.0	1
129	No association between antiviral treatment and risk of Alzheimer's disease in German outpatients. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2021, 7, e12216.	3.7	1
130	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021, 18, 147916412110338.	2.0	4
131	Prevalence of Lung Metastases among 19,321 Metastatic Colorectal Cancer Patients in Eight Countries of Europe and Asia. <i>Current Oncology</i> , 2021, 28, 5035-5040.	2.2	3
132	Prescription rates of common medications in patients with decompensated cirrhosis in Germany. <i>Zeitschrift Fur Gastroenterologie</i> , 2021, , .	0.5	0
133	Association between osteoarthritis and the incidence of Parkinson's disease in the United Kingdom. <i>Clinical Parkinsonism &amp; Related Disorders</i> , 2021, 5, 100120.	0.9	1
134	Portal Vein Thrombosis Is Associated with an Increased Incidence of Depression and Anxiety Disorders. <i>Journal of Clinical Medicine</i> , 2021, 10, 5689.	2.4	1
135	Incidence of Cancer in Patients with Irritable Bowl Syndrome. <i>Journal of Clinical Medicine</i> , 2021, 10, 5911.	2.4	2
136	Omega-3 fatty acids supplementation on major cardiovascular outcomes: an umbrella review of meta-analyses of observational studies and randomized controlled trials. <i>European Review for Medical and Pharmacological Sciences</i> , 2021, 25, 2079-2092.	0.7	1
137	Adherence to neuroleptic treatment in psychiatric practices: A retrospective study of 55 practices with more than 5000 bipolar and schizophrenic patients in Germany. <i>Psychiatry Research</i> , 2020, 284, 112758.	3.3	4
138	Impact of NAFLD on the Incidence of Cardiovascular Diseases in a Primary Care Population in Germany. <i>Digestive Diseases and Sciences</i> , 2020, 65, 2112-2119.	2.3	68
139	Association Between Migraine Headaches and Dry Eye Disease in Patients Studied in General Practices in Germany. <i>JAMA Ophthalmology</i> , 2020, 138, 223.	2.5	4
140	Long-term risk of stroke and its predictors in transient ischaemic attack patients in Germany. <i>European Journal of Neurology</i> , 2020, 27, 723-728.	3.3	8
141	Fecal and urinary incontinence are major problems associated with rectal cancer. <i>International Journal of Colorectal Disease</i> , 2020, 35, 35-40.	2.2	5
142	Comment on Mizokami-Stout et al. The Contemporary Prevalence of Diabetic Neuropathy in Type 1 Diabetes: Findings From the T1D Exchange. <i>Diabetes Care</i> 2020;43:806-812. <i>Diabetes Care</i> , 2020, 43, e113-e113.	8.6	1
143	Real-world treatment patterns in patients with nontuberculous mycobacterial lung disease in general and pneumologist practices in Germany. <i>Journal of Clinical Tuberculosis and Other Mycobacterial Diseases</i> , 2020, 20, 100178.	1.3	9
144	Changes in the utilization of blood glucose test strips among patients using intermittent scanning continuous glucose monitoring in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 922-928.	4.4	3

#	ARTICLE	IF	CITATIONS
145	Influence of chemotherapy and endocrine treatment on fractures in postmenopausal women with breast cancer – a retrospective cohort study. <i>Journal of Bone Oncology</i> , 2020, 22, 100292.	2.4	7
146	Factors associated with fracture after stroke and TIA: a long-term follow-up. <i>Osteoporosis International</i> , 2020, 31, 2395-2402.	3.1	12
147	Incidence of and factors associated with manic episodes and bipolar disorder in the decade following depression onset in Germany. <i>Journal of Affective Disorders</i> , 2020, 266, 534-539.	4.1	4
148	Non-alcoholic fatty liver disease increases the risk of incident chronic kidney disease. <i>United European Gastroenterology Journal</i> , 2020, 8, 942-948.	3.8	43
149	Panic buying or good adherence? Increased pharmacy purchases of drugs from wholesalers in the last week prior to Covid-19 lockdown. <i>Journal of Psychiatric Research</i> , 2020, 130, 19-21.	3.1	43
150	Proton pump inhibitors increase risk of bone fractures in men with cirrhosis: a population-based study. <i>Alimentary Pharmacology and Therapeutics</i> , 2020, 52, 1042-1050.	3.7	7
151	Multimorbidity Among Adult Outpatients With Type 1 Diabetes in Germany. <i>Journal of Diabetes Science and Technology</i> , 2020, , 193229682096526.	2.2	4
152	Proton pump inhibitor use is associated with a variety of infections in patients with liver cirrhosis. <i>Medicine (United States)</i> , 2020, 99, e23436.	1.0	10
153	Treatment Patterns of Patients with All-Cause Dementia in Russia. <i>Journal of Alzheimer's Disease Reports</i> , 2020, 4, 9-14.	2.2	4
154	Association between thyroid gland diseases and breast cancer: a case-control study. <i>Breast Cancer Research and Treatment</i> , 2020, 182, 207-213.	2.5	24
155	Effect of progestogen-only contraception on premenopausal fracture risk: a case-control study. <i>Osteoporosis International</i> , 2020, 31, 1801-1806.	3.1	3
156	Relationship between physical conditions and attempted or completed suicide in more than 9,300 individuals from the United Kingdom: a case-control study. <i>Journal of Affective Disorders</i> , 2020, 274, 457-463.	4.1	6
157	Increased prevalence of depression, anxiety, and adjustment and somatoform disorders in transsexual individuals. <i>Journal of Affective Disorders</i> , 2020, 274, 482-485.	4.1	9
158	Nonalcoholic Fatty Liver Disease Increases the Risk of Anxiety and Depression. <i>Hepatology Communications</i> , 2020, 4, 1293-1301.	4.3	70
159	Usage of antiepileptic drugs in different diseases in Germany – A retrospective study. <i>Epilepsy and Behavior</i> , 2020, 111, 107210.	1.7	4
160	Prevalence of insulin as a first-line therapy and associated factors in people with type 2 diabetes in German primary care practices. <i>Diabetic Medicine</i> , 2020, 37, 1333-1339.	2.3	6
161	Association between various thyroid gland diseases, TSH values and thyroid cancer: a case-control study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 2989-2994.	2.5	13
162	Prescription of hormone replacement therapy prior to and after the diagnosis of gynecological cancers in German patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020, 146, 1567-1573.	2.5	7

#	ARTICLE	IF	CITATIONS
163	Sex Differences in People Aging With HIV in Germany. <i>Journal of Acquired Immune Deficiency Syndromes</i> (1999), 2020, 84, e11-e11.	2.1	2
164	Comparative Analysis of the 3-Year Persistence Rate with Second-Line Vedolizumab and Tumor Necrosis Factor- $\alpha$ Inhibitors in Patients with Inflammatory Bowel Disease Followed in Gastroenterology Practices in Germany. <i>Digestive Diseases</i> , 2020, 38, 466-473.	1.9	8
165	Incidence of fractures in patients with type 1 diabetes mellitus—a retrospective study with 4420 patients. <i>Osteoporosis International</i> , 2020, 31, 1315-1322.	3.1	13
166	Urinary and fecal incontinence in stroke survivors followed in general practice: A retrospective cohort study. <i>Annals of Physical and Rehabilitation Medicine</i> , 2020, 63, 488-494.	2.3	10
167	Drug treatment for patients with bipolar disorders in psychiatric practices in Germany in 2009 and 2018. <i>Psychiatry Research</i> , 2020, 289, 112965.	3.3	19
168	Reduced antibiotic use after initial treatment of acute respiratory infections with phytopharmaceuticals- a retrospective cohort study. <i>Postgraduate Medicine</i> , 2020, 132, 412-418.	2.0	22
169	Tumor Incidence in Patients with Non-Alcoholic Fatty Liver Disease. <i>Deutsches Arzteblatt International</i> , 2020, 117, 719-724.	0.9	21
170	To Be Continued? Long-Term Treatment Effects of Antidepressant Drug Classes and Individual Antidepressants on the Risk of Developing Dementia. <i>Journal of Clinical Psychiatry</i> , 2020, 81, .	2.2	15
171	Persistence and adherence to dutasteride/tamsulosin fixed-dose versus free-combination alpha blocker/5ARI therapy in patients with benign prostate hyperplasia in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 37-49.	0.6	10
172	Prescription rates of cardiovascular and diabetes therapies prior to and during the COVID-19 lockdown in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 475-481.	0.6	16
173	Persistence with biological drugs in psoriasis patients followed in dermatology practices in Germany: A retrospective cohort study of 1,201 patients. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 74-81.	0.6	0
174	The role of the treating practice in persistence among dementia patients in Germany and the UK. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020, 58, 247-253.	0.6	0
175	Retrospektive Analyse zum Einsatz kurzwirksamer Beta-2-Sympathomimetika (SABA) bei Patienten mit Asthma in Deutschland. , 2020, 74, .		0
176	Overuse of short-acting beta-agonists (SABA) and risk factors among patients with asthma. , 2020, , .		0
177	Incidence of fractures in young women with breast cancer - a retrospective cohort study. <i>Journal of Bone Oncology</i> , 2019, 18, 100254.	2.4	13
178	Association Between Migraine Headaches and Dementia in More than 7,400 Patients Followed in General Practices in the United Kingdom. <i>Journal of Alzheimer's Disease</i> , 2019, 71, 353-360.	2.6	30
179	Are data from general practices suitable for survival analyses in the field of breast cancer? A retrospective study conducted in the United Kingdom, France and Germany. <i>Molecular and Clinical Oncology</i> , 2019, 11, 177-180.	1.0	1
180	Prescription-based prevalence of biological therapy in patients with psoriasis, rheumatoid arthritis, and inflammatory bowel diseases. <i>Biologics</i> , 2019, 61, 52-54.	1.4	4

#	ARTICLE	IF	CITATIONS
181	Demographic data and seasonal variation in peripheral vestibular disorders in ENT practices in Germany. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2019, 29, 181-190.	2.0	12
182	One-Year Persistence with Donepezil, Memantine, and Rivastigmine in More than 66,000 Elderly Patients Followed in Poland. <i>Journal of Alzheimer's Disease</i> , 2019, 70, 899-905.	2.6	3
183	Prevalence of valproate prescriptions in women of childbearing age in certain regions of Russia. <i>Epilepsy and Behavior</i> , 2019, 101, 106584.	1.7	3
184	Late Detection of Atrial Fibrillation after Stroke: Implications for the Secondary Prevention. <i>European Neurology</i> , 2019, 81, 262-269.	1.4	2
185	Increased risk for cancer after stroke at a young age: etiological relevance or incidental finding?. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 3047-3054.	2.5	8
186	Persistence with selective serotonin (norepinephrine) reuptake inhibitors in Germany – A retrospective database analysis. <i>Journal of Affective Disorders</i> , 2019, 247, 156-160.	4.1	3
187	Coded Prevalence of Mild Cognitive Impairment in General and Neuropsychiatrists Practices in Germany Between 2007 and 2017. <i>Journal of Alzheimer's Disease</i> , 2019, 67, 1313-1318.	2.6	6
188	Is There an Association Between Antiepileptic Drug Use and Dementia Risk? A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2019, 68, 97-103.	2.6	6
189	Persistence and treatment-free interval in patients being prescribed biological drugs in rheumatology practices in Germany. <i>European Journal of Clinical Pharmacology</i> , 2019, 75, 717-722.	1.9	1
190	The Relationship Between the Use of Antihypertensive Drugs and the Incidence of Dementia in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019, 70, 91-97.	2.6	13
191	Analysis of the Effects of Selective Serotonin (and Noradrenaline) Reuptake Inhibitors on the Risk of Dementia in Patients with Depression. <i>Journal of Alzheimer's Disease</i> , 2019, 69, 577-583.	2.6	7
192	Discontinuation of Antihypertensive Therapy and Dementia Incidence in the United Kingdom. <i>Journal of the American Medical Directors Association</i> , 2019, 20, 790-791.	2.5	0
193	Relationship between induced abortion and the incidence of depression, anxiety disorder, adjustment disorder, and somatoform disorder in Germany. <i>Journal of Psychiatric Research</i> , 2019, 114, 75-79.	3.1	14
194	Longitudinal changes in HbA1c following therapy intensification after inadequate response to two oral antidiabetic agents in patients with type 2 diabetes in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2019, 21, 1757-1757.	4.4	2
195	Diagnostic Behavior for Mild Cognitive Impairment in General and Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019, 68, 925-930.	2.6	5
196	Association between induced abortion, spontaneous abortion, and infertility respectively and the risk of psychiatric disorders in 57,770 women followed in gynecological practices in Germany. <i>Journal of Affective Disorders</i> , 2019, 251, 107-113.	4.1	19
197	Time to Insulin Initiation in Type 2 Diabetes Patients in 2010/2011 and 2016/2017 in Germany. <i>Journal of Diabetes Science and Technology</i> , 2019, 13, 1129-1134.	2.2	7
198	Real-world treatment patterns of sacubitril/valsartan: a longitudinal cohort study in Germany. <i>European Journal of Heart Failure</i> , 2019, 21, 588-597.	7.1	49

#	ARTICLE	IF	CITATIONS
199	Real-world prevalence and incidence of lower back pain in Germany. <i>Rheumatology International</i> , 2019, 39, 1113-1113.	3.0	3
200	Duration of sick leave in patients with depression treated by general practitioners and psychiatrists. <i>Psychiatry Research</i> , 2019, 279, 382-383.	3.3	4
201	Cancer risk in stroke survivors followed for up to 10 years in general practices in Germany. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019, 145, 1013-1020.	2.5	12
202	Incidence and Risk of Various Disorders of the External Ear in Patients With Hearing Aids Treated in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2019, 40, 1076-1081.	1.3	5
203	Prescription patterns of antiepileptic drugs in Kazakhstan in 2018: A retrospective study of 57,959 patients. <i>Epilepsy and Behavior</i> , 2019, 99, 106445.	1.7	4
204	Risk Factors for Tinnitus in 37,692 Patients Followed in General Practices in Germany. <i>Otology and Neurotology</i> , 2019, 40, 436-440.	1.3	11
205	Association Between Hearing and Vision Impairment and Risk of Dementia: Results of a Case-Control Study Based on Secondary Data. <i>Frontiers in Aging Neuroscience</i> , 2019, 11, 363.	3.4	32
206	Incidence of epilepsy and associated factors in elderly patients in Germany. <i>Epilepsy and Behavior</i> , 2019, 90, 107-111.	1.7	14
207	Trends in valproate use in patients in Germany between 2009 and 2017. <i>Epilepsy and Behavior</i> , 2019, 92, 26-30.	1.7	14
208	Persistence with biological drugs in patients treated in rheumatology practices in Germany. <i>Rheumatology International</i> , 2019, 39, 525-531.	3.0	8
209	Association between chronic somatic conditions and depression in children and adolescents: A retrospective study of 13,326 patients. <i>Journal of Affective Disorders</i> , 2019, 245, 697-701.	4.1	20
210	Impact of dronedarone on the risk of myocardial infarction and stroke in atrial fibrillation patients followed in general practices in Germany. <i>International Journal of Cardiology</i> , 2019, 278, 126-132.	1.7	17
211	Frequency of Blood Pressure and Estimated Glomerular Filtration Rate Testing in Type 2 Diabetes Mellitus: A Retrospective Study with 43,509 Patients. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2019, 127, 455-460.	1.2	4
212	Progression of mild cognitive impairment to dementia in German specialist practices. <i>Dementia</i> , 2019, 18, 380-390.	2.0	19
213	Lines of therapy with biological drugs in dermatology, gastroenterology, and rheumatology practices in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 182-187.	0.6	1
214	Prevalence of switching from two anti-TNF biosimilars back to biologic reference products in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 323-328.	0.6	6
215	The association between antihypertensive therapy and the incidence of Parkinson's disease in patients followed in general practices in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 483-488.	0.6	15
216	The role of general practitioners and psychiatrists in issuing initiation and follow-up prescriptions for selective serotonin (norepinephrine) reuptake inhibitors in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019, 57, 329-333.	0.6	1

#	ARTICLE	IF	CITATIONS
217	Association between gout and osteoporosis risk in the United Kingdom. <i>Osteoporosis International</i> , 2018, 29, 1685-1685.	3.1	2
218	Seasonal Variation of Epistaxis in Germany. <i>Journal of Craniofacial Surgery</i> , 2018, 29, e365-e367.	0.7	6
219	Frequency of Thyroid-Stimulating Hormone Monitoring in Patients After Thyroidectomy: A Retrospective Study in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2018, 126, 640-644.	1.2	1
220	Frequency of blood pressure and estimated glomerular filtration rate monitoring in patients affected by hypertension. <i>Blood Pressure Monitoring</i> , 2018, 23, 85-90.	0.8	6
221	Treatment patterns and low-density lipoprotein cholesterol (LDL-C) goal attainment among patients receiving high- or moderate-intensity statins. <i>Clinical Research in Cardiology</i> , 2018, 107, 380-388.	3.3	59
222	Long-term use of basal insulin and the risk of breast cancer. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 763-764.	2.5	4
223	Tramadol prescription patterns in patients followed by general practitioners and orthopedists in Germany in the year 2015. <i>Postgraduate Medicine</i> , 2018, 130, 37-41.	2.0	5
224	Association between depression and persistence with oral antihyperglycemic drugs in type 2 diabetes mellitus patients in Germany. <i>Psychiatry Research</i> , 2018, 261, 90-93.	3.3	9
225	Increased depression symptom score in newly diagnosed type 2 diabetes patients. <i>Psychiatry Research</i> , 2018, 261, 259-263.	3.3	11
226	Changes in patient characteristics, glucose lowering treatment, glycemic control and complications in type 2 diabetes in general practices (Disease Analyzer, Germany: 2008-2016). <i>Postgraduate Medicine</i> , 2018, 130, 244-250.	2.0	9
227	20-year risk of breast cancer recurrence. <i>Breast Cancer Research and Treatment</i> , 2018, 168, 765-766.	2.5	9
228	Time between suspected and confirmed diagnoses of Crohn's disease and ulcerative colitis in patients followed in gastroenterological practices in Germany. <i>International Journal of Colorectal Disease</i> , 2018, 33, 967-971.	2.2	1
229	Diagnoses of Dizziness- and Vertigo-related Disorders in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2018, 39, 474-480.	1.3	16
230	The impact of physician-level drug budgets on prescribing behavior. <i>European Journal of Health Economics</i> , 2018, 19, 213-222.	2.8	10
231	Prescription Patterns and Disease Control in Type 2 Diabetes Mellitus Patients in Nursing Home and Home Care Settings: A Retrospective Analysis in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018, 12, 136-139.	2.2	5
232	Persistence With Oral Antihyperglycemic Drugs in Type 2 Diabetes Mellitus Patients With Dementia in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018, 12, 140-144.	2.2	5
233	Prevalence of pain medication prescriptions in France, Germany, and the UK - a cross-sectional study including 4,270,142 patients. <i>Postgraduate Medicine</i> , 2018, 130, 32-36.	2.0	17
234	Low annual frequency of HbA <sub>1c</sub> testing in people with Type 2 diabetes in primary care practices in Germany. <i>Diabetic Medicine</i> , 2018, 35, 249-254.	2.3	10

#	ARTICLE	IF	CITATIONS
235	Prevalence and risk factors for delirium diagnosis in patients followed in general practices in Germany. <i>International Psychogeriatrics</i> , 2018, 30, 511-518.	1.0	15
236	PIH35 - <a href="https://ispor.confex.com/ispor/euro21/research/papers/index.cgi?username=85228&amp;password=173196">HTTPS://ISPOR.CONFEX.COM/ISPOR/EURO21/RESEARCH/PAPERS/INDEX.CGI?USERNAME=85228&amp;PASSWORD=173196</a> Value in Health, 2018, 21, S219.	0.1	0
237	PMH16 - THE PRESCRIPTION PATTERNS OF SELECTIVE SEROTONIN (NOREPINEPHRINE) REUPTAKE INHIBITORS IN GERMANY. <i>Value in Health</i> , 2018, 21, S278.	0.3	0
238	Impact of tobacco smoking on the risk of developing 25 different cancers in the UK: a retrospective study of 422,010 patients followed for up to 30 years. <i>Oncotarget</i> , 2018, 9, 17420-17429.	1.8	67
239	PMH65 - PERSISTENCE WITH SELECTIVE SEROTONIN (NOREPINEPHRINE) REUPTAKE INHIBITOR THERAPY IN GERMANY. <i>Value in Health</i> , 2018, 21, S285.	0.3	0
240	Prevalence of vulvovaginal candidiasis in gynecological practices in Germany: A retrospective study of 954,186 patients. <i>Current Medical Mycology</i> , 2018, 4, 6-11.	0.8	26
241	Current healthcare pathways in the treatment of rhinosinusitis in Germany. <i>Acta Oto-Laryngologica</i> , 2018, 138, 1086-1091.	0.9	4
242	PHP337 - OFF-LABEL PRESCRIPTION OF DRUGS COMPARED TO PRESCRIPTION AFTER LABEL EXTENSION IN CHILDREN FOR GERMANY. <i>Value in Health</i> , 2018, 21, S207-S208.	0.3	0
243	P1930Dronedarone is associated with a lower risk of myocardial infarction in Germany - a retrospective database analysis. <i>European Heart Journal</i> , 2018, 39, .	2.2	1
244	Medication use in patients with chronic rhinosinusitis in Germany - a large retrospective patient-based study. <i>Rhinology</i> , 2018, 57, 94-100.	1.3	5
245	Switching the manufacturer of antiepileptic drugs is associated with higher risk of seizures: A nationwide study of prescription data in Germany. <i>Annals of Neurology</i> , 2018, 84, 918-925.	5.3	38
246	Relevance of Coded Prodromal Mild Cognitive Impairment in the Routine Treatment of Patients with Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 393-399.	2.6	3
247	Diagnoses of suspected cancer in ENT otolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018, 9, 459-463.	1.0	3
248	Association Between the Use of Antihyperglycemic Drugs and Dementia Risk: A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2018, 66, 725-732.	2.6	40
249	Association between sleep disorders and the presence of breast cancer metastases in gynecological practices in Germany: a case-control study of 11,412 women. <i>Breast Cancer Research and Treatment</i> , 2018, 171, 443-448.	2.5	16
250	The tamoxifen paradox: influence of adjuvant tamoxifen on fracture risk in pre- and postmenopausal women with breast cancer. <i>Osteoporosis International</i> , 2018, 29, 2557-2564.	3.1	27
251	The impact of non-persistence on the direct and indirect costs in patients treated with subcutaneous tumour necrosis factor-alpha inhibitors in Germany. <i>Rheumatology</i> , 2018, 57, 1276-1281.	1.9	7
252	Association of characteristics of people with type 2 diabetes mellitus with discordant values of fasting glucose and HbA1c. <i>Journal of Diabetes</i> , 2018, 10, 934-941.	1.8	11

#	ARTICLE	IF	CITATIONS
253	Multimorbidity and polypharmacy among elderly people followed in general practices in Germany. <i>European Journal of Internal Medicine</i> , 2018, 55, 66-68.	2.2	20
254	Association of laryngeal cancer with vocal cord leukoplakia and associated risk factors in 1,184 patients diagnosed in otorhinolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018, 8, 689-693.	1.0	17
255	Impact of Osteoporosis on the Risk of Dementia in Almost 60,000 Patients Followed in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 65, 401-407.	2.6	22
256	The Influence of Polypharmacy on the Initiation of Anti-Dementia Therapy in Germany. <i>Journal of Alzheimer's Disease</i> , 2018, 64, 827-833.	2.6	9
257	Potential delay in the diagnosis of vulvar cancer and associated risk factors in women treated in German gynecological practices. <i>Oncotarget</i> , 2018, 9, 8725-8730.	1.8	16
258	Diagnosing and treating dementia in German primary and specialized care between 2011 and 2015. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 301-309.	0.6	7
259	Differences in baseline characteristics between type 2 diabetes mellitus patients treated with dipeptidyl peptidase-4 inhibitors in randomized controlled trials and those receiving the same treatment in real-world settings. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 411-416.	0.6	3
260	Basic characteristics and representativeness of the German Disease Analyzer database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 459-466.	0.6	261
261	Persistence with antihypertensive drugs in patients with depression in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 162-168.	0.6	4
262	Prevalence and treatment of sexually transmitted diseases in gynecological practices in Germany: A retrospective study with more than 1,000,000 patients. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 212-216.	0.6	0
263	Prevalence and treatment of sexually transmitted infections in men followed by urologists in Germany - a cross sectional study with 347,090 men. <i>GMS German Medical Science</i> , 2018, 16, Doc03.	2.7	5
264	Basic characteristics and representativeness of the German Disease Analyzer database©. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018, 56, 459-466.	0.6	73
265	Impact of Disease Management Programs on HbA1c Values in Type 2 Diabetes Patients in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 117-122.	2.2	20
266	Hip fracture risk in patients with dementia in German primary care practices. <i>Dementia</i> , 2017, 16, 853-864.	2.0	17
267	Risk factors and outcomes in well-selected vaginal breech deliveries: a retrospective observational study. <i>Journal of Perinatal Medicine</i> , 2017, 45, 291-297.	1.4	17
268	Compliance with vaccination against tick-borne encephalitis virus in Germany. <i>Clinical Microbiology and Infection</i> , 2017, 23, 460-463.	6.0	14
269	Calculated Daily Insulin Dosages Overestimate Prescribed Insulin Doses in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 597-601.	2.2	3
270	Impact of prior breast cancer on mode of delivery and pregnancy-associated disorders: a retrospective analysis of subsequent pregnancy outcomes. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 1069-1074.	2.5	22



#	ARTICLE	IF	CITATIONS
271	Association of dipeptidyl peptidase 4 inhibitors with risk of metastases in patients with type 2 diabetes and breast, prostate or digestive system cancer. <i>Journal of Diabetes and Its Complications</i> , 2017, 31, 687-692.	2.3	19
272	Oral contraceptive use and fracture risk—a retrospective study of 12,970 women in the UK. <i>Osteoporosis International</i> , 2017, 28, 2349-2355.	3.1	18
273	Seasonality of Insulin Use in German Outpatients With Diabetes: A Retrospective Analysis. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 996-1000.	2.2	4
274	Use of dipeptidyl peptidase-4 inhibitors and risk of bone fracture in patients with type 2 diabetes in Germany—a retrospective analysis of real-world data. <i>Osteoporosis International</i> , 2017, 28, 2421-2428.	3.1	52
275	Incidence of depression and anxiety among women newly diagnosed with breast or genital organ cancer in Germany. <i>Psycho-Oncology</i> , 2017, 26, 1535-1540.	2.3	28
276	Persistence with antiepileptic drugs in epilepsy patients treated in neurological practices in Germany. <i>Epilepsy and Behavior</i> , 2017, 73, 204-207.	1.7	8
277	Prevalence of and risk factors associated with cesarean section in Lebanon — A retrospective study based on a sample of 29,270 women. <i>Women and Birth</i> , 2017, 30, e265-e271.	2.0	26
278	Depression Risk in Patients with Rheumatoid Arthritis in the United Kingdom. <i>Rheumatology and Therapy</i> , 2017, 4, 195-200.	2.3	41
279	Association between Anti-Dementia Treatment Persistence and Daily Dosage of the First Prescription: A Retrospective Analysis in Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 58, 37-44.	2.6	10
280	Changes in Glycemic Control and Body Weight After Initiation of Dapagliflozin or Basal Insulin Supported Oral Therapy in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 590-596.	2.2	4
281	Long-term use of benzodiazepines in older patients in Germany: a retrospective analysis. <i>Therapeutic Advances in Psychopharmacology</i> , 2017, 7, 191-200.	2.7	27
282	Risk of Nonfatal Stroke in Type 2 Diabetes Mellitus Patients: A Retrospective Comparison Between Disease Management Programs and Standard Care. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 808-813.	2.2	4
283	Risk Factors for Mild Cognitive Impairment in German Primary Care Practices. <i>Journal of Alzheimer's Disease</i> , 2017, 56, 379-384.	2.6	11
284	Incidence of Psychiatric Disorders in Women WITH a Suspected But Unconfirmed Diagnosis of Breast or Genital Organ Cancer. <i>Value in Health</i> , 2017, 20, A709-A710.	0.3	0
285	Risk of depression, anxiety, and adjustment disorders in women with a suspected but unconfirmed diagnosis of breast or genital organ cancer in Germany. <i>Cancer Causes and Control</i> , 2017, 28, 1021-1026.	1.8	10
286	Fracture risk reduction with use of dipeptidyl peptidase-4 inhibitors: response to Driessen et al.. <i>Osteoporosis International</i> , 2017, 28, 2431-2431.	3.1	2
287	Prevalence of high-risk cardiovascular patients with therapy-resistant hypercholesterolemia. <i>Cardiovascular Endocrinology</i> , 2017, 6, 81-85.	0.8	6
288	Adherence to antiepileptic drugs in children and adolescents: A retrospective study in primary care settings in Germany. <i>Epilepsy and Behavior</i> , 2017, 75, 36-41.	1.7	21

#	ARTICLE	IF	CITATIONS
289	The impact of depot medroxyprogesterone acetate on fracture risk: a case-control study from the UK. <i>Osteoporosis International</i> , 2017, 28, 291-297.	3.1	35
290	Depression risk in patients with late-onset rheumatoid arthritis in Germany. <i>Quality of Life Research</i> , 2017, 26, 437-443.	3.1	20
291	Psoriasis risk in patients with type 2 diabetes in German primary care practices. <i>Primary Care Diabetes</i> , 2017, 11, 52-56.	1.8	12
292	Prescription Patterns and the Cost of Antihyperglycemic Drugs in Patients With Type 2 Diabetes Mellitus in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 123-127.	2.2	12
293	Treatment Outcomes and Tolerability Following Initiation of GLP-1 Receptor Agonists Among Type 2 Diabetes Patients in Primary Care Practices in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017, 11, 272-277.	2.2	6
294	Impact of attention deficit hyperactivity disorder therapy on fracture risk in children treated in German pediatric practices. <i>Osteoporosis International</i> , 2017, 28, 1265-1269.	3.1	10
295	Prescription Patterns and the Cost of Migraine Treatments in German General and Neurological Practices. <i>Pain Practice</i> , 2017, 17, 747-752.	1.9	8
296	Factors Associated with an Adherence with Antiepileptic Drugs in Children Treated in Pediatric Practices in Germany. <i>Value in Health</i> , 2017, 20, A402.	0.3	0
297	The Presentation Of Patients With Epistaxis Shows A Marked Seasonal Variation In Germany. <i>Value in Health</i> , 2017, 20, A602.	0.3	0
298	Persistence in Hypertension Therapy in Patients with and Without Depression in Germany. <i>Value in Health</i> , 2017, 20, A620.	0.3	0
299	Spontaneous Abortion is Associated with an Increased Risk of Developing Psychiatric Diseases. <i>Value in Health</i> , 2017, 20, A716.	0.3	0
300	Persistence with Antihypertensive Drugs in Patients with Hypertension and Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 60, 505-510.	2.6	4
301	Prevalence of depression, anxiety, and adjustment disorders in women with spontaneous abortion in Germany – A retrospective cohort study. <i>Psychiatry Research</i> , 2017, 258, 382-386.	3.3	22
302	Prevalence of Use of Cardiovascular Drugs in Dementia Patients Treated in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017, 56, 1519-1524.	2.6	5
303	Suspected cancer diagnoses made by general practitioners in a population with subsequently confirmed cancer diagnoses in Germany: a retrospective study of 31,628 patients. <i>Oncotarget</i> , 2017, 8, 84540-84545.	1.8	4
304	Prevalence of depression, anxiety, adjustment disorders, and somatoform disorders in patients with age-related macular degeneration in Germany. <i>GMS German Medical Science</i> , 2017, 15, Doc04.	2.7	7
305	Conflicts at work are associated with a higher risk of cardiovascular disease. <i>GMS German Medical Science</i> , 2017, 15, Doc08.	2.7	6
306	Risk factors for epistaxis in patients followed in general practices in Germany. <i>Rhinology</i> , 2017, 55, 312-318.	1.3	12

#	ARTICLE	IF	CITATIONS
307	Prescription patterns and drug costs in German patients with dementia in nursing homes and home-care settings. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 9-15.	0.6	24
308	Risk of dementia in German patients treated with antidepressants in general or psychiatric practices. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 322-328.	0.6	10
309	Treatment persistence in the use of basal insulins in Poland and Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 119-125.	0.6	4
310	Patterns of benzodiazepine prescribing by neuropsychiatrists and general practitioners for elderly patients in Germany in 2014. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 466-471.	0.6	9
311	Regional differences in insulin therapy regimens in five European countries. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 403-408.	0.6	7
312	Prevalence and predictors of prescription of antibiotics in pregnant women treated by gynecologists in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 643-649.	0.6	7
313	Prevalence of bisphosphonate therapy in women with breast cancer treated with aromatase inhibitors in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 394-396.	0.6	1
314	Benzodiazepine prescribing in the elderly in Germany and Korea: A comparison of two observational studies. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 480-482.	0.6	0
315	Compliance with vaccination against hepatitis A virus in Germany: a retrospective analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017, 55, 740-745.	0.6	4
316	Risk factors for epistaxis in patients followed in general practices in Germany. <i>Rhinology</i> , 2017, 55, 312-318.	1.3	0
317	Risk factors for ectopic pregnancy in Germany: a retrospective study of 100,197 patients. <i>GMS German Medical Science</i> , 2017, 15, Doc19.	2.7	8
318	Adherence to GLP-1 receptor agonist therapy administered by once-daily or once-weekly injection in patients with type 2 diabetes in Germany. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, Volume 9, 201-205.	2.4	65
319	Gender- and age-related treatment compliance in patients with osteoporosis in Germany. <i>Patient Preference and Adherence</i> , 2016, Volume 10, 2379-2385.	1.8	21
320	Do patients with lactose intolerance exhibit more frequent comorbidities than patients without lactose intolerance? An analysis of routine data from German medical practices. <i>Annals of Gastroenterology</i> , 2016, 29, 174-9.	0.6	18
321	Changes in HbA1c, body weight, and systolic blood pressure in type 2 diabetes patients initiating dapagliflozin therapy: a primary care database study. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, Volume 9, 337-345.	2.4	24
322	Fracture risk in patients with type 2 diabetes under different antidiabetic treatment regimens: a retrospective database analysis in primary care. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016, 9, 17.	2.4	23
323	Depression risk in patients with coronary heart disease in Germany. <i>World Journal of Cardiology</i> , 2016, 8, 547.	1.5	19
324	Treatment of depression in cancer and non-cancer patients in German neuropsychiatric practices. <i>Psycho-Oncology</i> , 2016, 25, 1324-1328.	2.3	16

#	ARTICLE	IF	CITATIONS
325	Change in glycated haemoglobin levels after initiating second-line therapy in type 2 diabetes: a primary care database study. <i>Diabetes, Obesity and Metabolism</i> , 2016, 18, 840-843.	4.4	14
326	High Risk of Depression In Patients with Heart Failure in Germany. <i>Value in Health</i> , 2016, 19, A636.	0.3	0
327	Depression risk in patients with heart failure in primary care practices in Germany. <i>International Psychogeriatrics</i> , 2016, 28, 1889-1894.	1.0	9
328	Risk factors for dementia diagnosis in German primary care practices. <i>International Psychogeriatrics</i> , 2016, 28, 1059-1065.	1.0	79
329	Treatment Outcomes Following Initiation of Dapagliflozin or Basal Supported Oral Therapy Among Type 2 Diabetes Patients In Primary Care Practices In Germany. <i>Value in Health</i> , 2016, 19, A666-A667.	0.3	0
330	Rheumatoid Arthritis is Associated with an Increased Risk of Depression in Elderly Patients in Germany. <i>Value in Health</i> , 2016, 19, A521.	0.3	0
331	Prevalence of Therapy-Resistant Hypercholesterolemia in High-Risk Cardiovascular Patients in Germany. <i>Value in Health</i> , 2016, 19, A659.	0.3	0
332	Cost of Antihyperglycemic Therapy In Type 2 Diabetes Patients Depending on HBA1C And Bmi Values. <i>Value in Health</i> , 2016, 19, A665.	0.3	0
333	Prescription Patterns And Costs Of Migraine Therapy In Germany. <i>Value in Health</i> , 2016, 19, A430.	0.3	0
334	Prescription Patterns and Drugs Costs in Patients with Dementia - A Retrospective Database Analysis. <i>Value in Health</i> , 2016, 19, A439.	0.3	0
335	Age-related differences in persistence with bisphosphonates in women with metastatic breast cancer. <i>Journal of Bone Oncology</i> , 2016, 5, 63-66.	2.4	7
336	Depression risk in female patients with osteoporosis in primary care practices in Germany. <i>Osteoporosis International</i> , 2016, 27, 2739-2744.	3.1	16
337	Impact of metformin on metastases in patients with breast cancer and type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 1056-1059.	2.3	27
338	Impact of comorbidities on the cost of depression drug therapy in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2016, 83, 130-136.	3.1	2
339	Comorbid disorders associated with asthma in children in Germany – National analysis of pediatric primary care data. <i>Pediatric Allergy and Immunology</i> , 2016, 27, 861-866.	2.6	14
340	Caesarean section and its impact on fertility and time to a subsequent pregnancy in Germany: a database analysis in gynecological practices. <i>Archives of Gynecology and Obstetrics</i> , 2016, 294, 1005-1010.	1.7	9
341	Nonadherence to antiepileptic drugs in Germany. <i>Neurology</i> , 2016, 87, 466-472.	1.1	62
342	Prevalence of medically treated children with ADHD and type 1 diabetes in Germany – Analysis of two representative databases. <i>Journal of Pediatric Endocrinology and Metabolism</i> , 2016, 29, 1293-1297.	0.9	21

#	ARTICLE	IF	CITATIONS
343	Age-related differences in persistence in women with breast cancer treated with tamoxifen or aromatase inhibitors in Germany. <i>Journal of Geriatric Oncology</i> , 2016, 7, 169-175.	1.0	13
344	Increased treatment persistence and its determinants in women with osteoporosis with prior fracture compared to those without fracture. <i>Osteoporosis International</i> , 2016, 27, 963-969.	3.1	11
345	The use of proton pump inhibitors is positively associated with osteoporosis in postmenopausal women in Germany. <i>Climacteric</i> , 2016, 19, 478-481.	2.4	17
346	Cancer is associated with intraoperative and postprocedural complications and disorders. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 777-781.	2.5	4
347	Changes in Type 2 Diabetes Mellitus Patients in German Primary Care Prior to (2006) and After (2010,) Tj ETQq1 1 0,784314 rgBT /Over 2.2 15	2.2	15
348	Second-line therapy with biological drugs in rheumatoid arthritis patients in German rheumatologist practices: a retrospective database analysis. <i>Rheumatology International</i> , 2016, 36, 1113-1118.	3.0	10
349	Referral for occupational therapy after diagnosis of developmental disorder by German child psychiatrists. <i>Early Child Development and Care</i> , 2016, 186, 959-970.	1.3	0
350	Prevalence of depression in type 2 diabetes patients in German primary care practices. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 432-437.	2.3	36
351	Risk of stillbirth in pregnant women with obesity in the United Kingdom. <i>Obesity Research and Clinical Practice</i> , 2016, 10, 574-579.	1.8	10
352	The impact of treatment compliance on fracture risk in women with breast cancer treated with aromatase inhibitors in the United Kingdom. <i>Breast Cancer Research and Treatment</i> , 2016, 155, 151-157.	2.5	32
353	Prevalence of depression, anxiety and their risk factors in German women with breast cancer in general and gynecological practices. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016, 142, 447-452.	2.5	42
354	Induction of labor in breech presentations at term: a retrospective observational study. <i>Archives of Gynecology and Obstetrics</i> , 2016, 293, 549-555.	1.7	22
355	Predictors of early discontinuation of basal insulin therapy in type 2 diabetes in primary care. <i>Primary Care Diabetes</i> , 2016, 10, 142-147.	1.8	7
356	Diabetes treatment in people with type 2 diabetes and schizophrenia: Retrospective primary care database analyses. <i>Primary Care Diabetes</i> , 2016, 10, 36-40.	1.8	9
357	Clinical and Patient-Related Variables Associated with Initiating GLP-1 Receptor Agonist Therapy in Type 2 Diabetes Patients in Primary Care in Germany. <i>PLoS ONE</i> , 2016, 11, e0152281.	2.5	9
358	Prevalence of chronic diseases among older patients in German general practices. <i>GMS German Medical Science</i> , 2016, 14, Doc03.	2.7	27
359	Persistence with antidepressant drugs in patients with dementia: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 323-329.	0.6	13
360	Persistence with antipsychotics in dementia patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 835-840.	0.6	13

#	ARTICLE	IF	CITATIONS
361	Treatment of depression in patients with cardiovascular diseases by German psychiatrists. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 557-563.	0.6	1
362	Prevalence and type of antidepressant therapy used by German general practitioners to treat female patients with osteoporosis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016, 54, 743-749.	0.6	1
363	Gender-based differences in the antidepressant treatment of patients with depression in German psychiatric practices. <i>GMS German Medical Science</i> , 2016, 14, Doc02.	2.7	2
364	Impact of caesarean section on mode of delivery, pregnancy-induced and pregnancy-associated disorders, and complications in the subsequent pregnancy in Germany. <i>GMS German Medical Science</i> , 2016, 14, Doc06.	2.7	10
365	Continuous treatment with antedementia drugs in Germany 2003–2013: a retrospective database analysis. <i>International Psychogeriatrics</i> , 2015, 27, 1335-1342.	1.0	29
366	Therapy Of Depression In Cancer And Non-Cancer Patients In German Neuropsychiatric Practices. <i>Value in Health</i> , 2015, 18, A431.	0.3	0
367	Reduction In C-Reactive Protein With Biologic Drugs In Rheumatoid Arthritis and Spondylitis Patients In German Rheumatologist Practices. <i>Value in Health</i> , 2015, 18, A635.	0.3	2
368	Prevalence Of Depression And Anxiety And Their Risk Factors In Women With Breast Cancer In German Primary Care. <i>Value in Health</i> , 2015, 18, A431.	0.3	2
369	No reduced risk of overall, colorectal, lung, breast, and prostate cancer with metformin therapy in diabetic patients: database analyses from Germany and the UK. <i>Pharmacoepidemiology and Drug Safety</i> , 2015, 24, 865-874.	1.9	81
370	Glycemic control after initiating basal insulin therapy in patients with type 2 diabetes: a primary care database analysis. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2015, 8, 45.	2.4	19
371	Are Sulfonylurea and Insulin Therapies Associated With a Larger Risk of Cancer Than Metformin Therapy? A Retrospective Database Analysis. <i>Diabetes Care</i> , 2015, 38, 59-65.	8.6	41
372	Trends and patterns of hormonal contraceptive prescribing for adolescents in primary care in the UK. <i>Journal of Family Planning and Reproductive Health Care</i> , 2015, 41, 216-222.	0.8	18
373	Time to pregnancy in subfertile women in German gynecological practices: analysis of a representative cohort of more than 60,000 patients. <i>Archives of Gynecology and Obstetrics</i> , 2015, 291, 657-662.	1.7	9
374	Fracture risk in patients with newly diagnosed type 2 diabetes: a retrospective database analysis in primary care. <i>Journal of Diabetes and Its Complications</i> , 2015, 29, 766-770.	2.3	56
375	Impact of disease management programs on women with breast cancer in Germany. <i>Breast Cancer Research and Treatment</i> , 2015, 153, 391-395.	2.5	19
376	Early Discontinuation and Related Treatment Costs After Initiation of Basal Insulin in Type 2 Diabetes Patients. <i>Journal of Diabetes Science and Technology</i> , 2015, 9, 644-650.	2.2	12
377	Discontinuation rates of menopausal hormone therapy among postmenopausal women in the post-WHI study era. <i>Climacteric</i> , 2015, 18, 737-742.	2.4	15
378	Persistence with opioid treatment in Germany in patients suffering from chronic non-malignant or cancer pain. <i>Current Medical Research and Opinion</i> , 2015, 31, 1157-1163.	1.9	8

#	ARTICLE	IF	CITATIONS
379	Treatment persistence after initiating basal insulin in type 2 diabetes patients: A primary care database analysis. <i>Primary Care Diabetes</i> , 2015, 9, 377-384.	1.8	17
380	Emergency peripartum hysterectomy and risk factors by mode of delivery and obstetric history: a 10-year review from Helsinki University Central Hospital. <i>Journal of Perinatal Medicine</i> , 2015, 43, 721-8.	1.4	11
381	Persistency with estrogen replacement therapy among hysterectomized women after the Women's Health Initiative study. <i>Climacteric</i> , 2015, 18, 826-834.	2.4	7
382	Effects of selected antidiabetics on weight loss – A retrospective database analysis. <i>Primary Care Diabetes</i> , 2015, 9, 74-77.	1.8	10
383	Risk of hypoglycaemia in type 2 diabetes patients under different insulin regimens: a primary care database analysis. <i>GMS German Medical Science</i> , 2015, 13, Doc01.	2.7	3
384	Association of time-to-levodopa with initial Parkinsonian medication: a retrospective cohort study. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2015, 53, 107-114.	0.6	2
385	Use of a German longitudinal prescription database (LRx) in pharmacoepidemiology. <i>GMS German Medical Science</i> , 2015, 13, Doc14.	2.7	6
386	Predictors of Insulin Initiation in Metformin and Sulfonylurea Users in Primary Care Practices. <i>Journal of Diabetes Science and Technology</i> , 2014, 8, 1023-1028.	2.2	6
387	Macro- and Microvascular Outcomes in Patients with Type 2 Diabetes Treated with Rapid-Acting Insulin Analogues or Human Regular Insulin: a Retrospective Database Analysis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014, 122, 92-99.	1.2	6
388	Use of antidiabetic agents in the treatment of gestational diabetes mellitus in Germany, 2008–2012. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014, 40, 1592-1597.	1.3	7
389	Discontinuation of Treatment Using Anticholinergic Medications in Patients With Urinary Incontinence. <i>Obstetrics and Gynecology</i> , 2014, 124, 794-800.	2.4	32
390	Impact of Rotavirus Vaccination in Germany. <i>Pediatric Infectious Disease Journal</i> , 2014, 33, e299-e304.	2.0	33
391	Predictors of hypoglycaemia in insulin-treated type 2 diabetes patients in primary care: A retrospective database analysis. <i>Primary Care Diabetes</i> , 2014, 8, 127-131.	1.8	31
392	Risk factors for discontinuation of insulin pump therapy in pediatric and young adult patients. <i>Primary Care Diabetes</i> , 2014, 8, 346-351.	1.8	8
393	Risk of venous thrombosis in users of hormonal contraceptives in German gynaecological practices: a patient database analysis. <i>Archives of Gynecology and Obstetrics</i> , 2014, 289, 413-419.	1.7	24
394	Prevalence and risk factors of neuropathy in newly diagnosed type 2 diabetes in primary care practices: A retrospective database analysis in Germany and UK. <i>Primary Care Diabetes</i> , 2014, 8, 250-255.	1.8	46
395	Sociodemographic Disparities in Administration of Antiepileptic Drugs to Adults with Epilepsy in Germany: A Retrospective, Database Study of Drug Prescriptions. <i>CNS Drugs</i> , 2014, 28, 753-759.	5.9	24
396	Differences in persistency with teriparatide in patients with osteoporosis according to gender and health care provider. <i>Osteoporosis International</i> , 2014, 25, 2721-2728.	3.1	24

#	ARTICLE	IF	CITATIONS
397	Prevalence of Chronic Diseases Among Older Patients (>65 Years) in German General Practitioner Practices. Value in Health, 2014, 17, A505.	0.3	0
398	Treatment Discontinuation In Patients With Urinary Incontinence Suffering From Glaucoma. Value in Health, 2014, 17, A471.	0.3	0
399	Which Adverse Effects Influence the Dropout Rate in Selective Serotonin Reuptake Inhibitor (Ssri) Treatment?. Value in Health, 2014, 17, A460.	0.3	5
400	Comparing Quality Effects Of Patient Care In Integrated And Regular Care For Patients With Hypertension. Value in Health, 2014, 17, A498.	0.3	1
401	Physicians' influence on breast cancer patient compliance. GMS German Medical Science, 2014, 12, Doc03.	2.7	7
402	Risk of psychiatric and neurological diseases in patients with workplace mobbing experience in Germany: a retrospective database analysis. GMS German Medical Science, 2014, 12, Doc10.	2.7	18
403	Which adverse effects influence the dropout rate in selective serotonin reuptake inhibitor (SSRI) treatment? Results for 50,824 patients. GMS German Medical Science, 2014, 12, Doc15.	2.7	13
404	Impact of comorbidities on the treatment of atopic dermatitis in clinical practice. International Journal of Clinical Pharmacology and Therapeutics, 2014, 52, 726-731.	0.6	6
405	Gender differences in persistency to bisphosphonates in patients with metastatic breast and prostate cancer. International Journal of Clinical Pharmacology and Therapeutics, 2014, 52, 352-359.	0.6	2
406	Treatment persistence evaluation of tamoxifen and aromatase inhibitors in breast cancer patients in early and late stage disease. International Journal of Clinical Pharmacology and Therapeutics, 2014, 52, 933-939.	0.6	10
407	Importance of durational hormone therapy in breast cancer patients in Germany: Retrospective database analysis.. Journal of Clinical Oncology, 2014, 32, 1603-1603.	1.6	0
408	Association of diabetes and cancer-diagnoses in primary care practices in Germany.. Journal of Clinical Oncology, 2014, 32, 1604-1604.	1.6	1
409	Secondary analysis of electronic databases: potentials and limitations. Reply to Asghari S, Mahdavian M [letter]. Diabetologia, 2013, 56, 2098-2099.	6.3	1
410	Diabetic retinopathy at diagnosis of type 2 diabetes in the UK: a database analysis. Diabetologia, 2013, 56, 109-111.	6.3	73
411	Persistence with bisphosphonates in patients with metastatic breast cancer: a retrospective database analysis. Journal of Cancer Research and Clinical Oncology, 2013, 139, 1149-1155.	2.5	13
412	Risk of Psychiatric and Neurological Diseases in Patients with Workspace Mobbing Experience in Germany: A Retrospective Database Analysis. Value in Health, 2013, 16, A541.	0.3	2
413	Influence of macro- and microvascular comorbidity on time to insulin initiation in type 2 diabetes patients: A retrospective database analysis in Germany, France, and UK. Primary Care Diabetes, 2013, 7, 167-171.	1.8	6
414	The Prescribing of Contraceptives for Adolescents in German Gynecologic Practices in 2007 and 2011: A Retrospective Database Analysis. Journal of Pediatric and Adolescent Gynecology, 2013, 26, 261-264.	0.7	8



#	ARTICLE	IF	CITATIONS
415	Prevalence and treatment profile of patients with grass pollen and house dust mite allergy. JDDG - Journal of the German Society of Dermatology, 2013, 11, 653-661.	0.8	11
416	Resource Consumption and Costs of Treatment in Patients with Type 1 Diabetes under Intensified Conventional Therapy under German Real-Life Conditions. Journal of Diabetes Science and Technology, 2013, 7, 736-742.	2.2	5
417	Treatment persistence, hypoglycaemia and clinical outcomes in type 2 diabetes patients with dipeptidyl peptidase-4 inhibitors and sulphonylureas: a primary care database analysis. Diabetes, Obesity and Metabolism, 2013, 15, 55-61.	4.4	78
418	Risk Factors for Discontinuation of Insulin Pump Therapy in Pediatric and Young Adult Patient Groups. Value in Health, 2013, 16, A444.	0.3	0
419	Persistence in Patients with Breast Cancer Treated with Tamoxifen or Aromatase Inhibitors- Analysis Based on Oncology Analyzer Database. Value in Health, 2013, 16, A418-A419.	0.3	1
420	Pregnancy After Breast Cancer in 2000-2002 and 2010-2012: A Retrospective Database Analysis. Value in Health, 2013, 16, A393.	0.3	0
421	Persistence in patients with breast cancer treated with tamoxifen or aromatase inhibitors: a retrospective database analysis. Breast Cancer Research and Treatment, 2013, 138, 185-191.	2.5	85
422	Changes in time to insulin initiation in type 2 diabetes patients: A retrospective database analysis in Germany and UK (2005-2010). Primary Care Diabetes, 2013, 7, 229-233.	1.8	20
423	Different Injection Frequencies of Basal Insulins in Type 2 Diabetes Patients under Real-Life Conditions: A Retrospective Database Analysis. Journal of Diabetes Science and Technology, 2013, 7, 1354-1358.	2.2	7
424	Prevalence of female subfertility in German gynecological practices. Gynecological Endocrinology, 2013, 29, 767-770.	1.7	7
425	Lower incidence of recorded cardiovascular outcomes in patients with type 2 diabetes using insulin aspart vs. those on human regular insulin: observational evidence from general practices. Diabetes, Obesity and Metabolism, 2013, 15, 358-363.	4.4	16
426	The risk of developing depression when suffering from neurological diseases. GMS German Medical Science, 2013, 11, Doc02.	2.7	17
427	Economic prescribing of corticosteroid nasal sprays in Germany. Comparison of mometasone and budesonide nasal sprays on the basis of the DDD, the PDD and reference prices. International Journal of Clinical Pharmacology and Therapeutics, 2013, 51, 12-18.	0.6	2
428	Frequency of hospitalizations prior to and after conversion to a rebate pharmaceutical in depression patients in Germany. International Journal of Clinical Pharmacology and Therapeutics, 2013, 51, 416-422.	0.6	3
429	Adherence in tamoxifen therapy after conversion to a rebate pharmaceutical in breast cancer patients in Germany. International Journal of Clinical Pharmacology and Therapeutics, 2013, 51, 969-975.	0.6	15
430	Cost comparison of outpatient treatment with granulocyte colony-stimulating factors (G-CSF) in Germany. International Journal of Clinical Pharmacology and Therapeutics, 2012, 50, 281-289.	0.6	24
431	Risk of diabetic foot ulceration during treatment with insulin glargine and NPH insulin. Journal of Wound Care, 2012, 21, 483-489.	1.2	4
432	Real-life treatment patterns, compliance, persistence, and medication costs in patients with hypertension in Germany. Journal of Medical Economics, 2012, 15, 155-165.	2.1	18

#	ARTICLE	IF	CITATIONS
433	Prevalence, utilization, and costs of antiepileptic drugs for epilepsy in Germany—a nationwide population-based study in children and adults. <i>Journal of Neurology</i> , 2012, 259, 2376-2384.	3.6	116
434	PHS10 HbA1c Testing Frequency in Primary Care Diabetes Patients in Germany and in the UK. <i>Value in Health</i> , 2012, 15, A519.	0.3	1
435	Effects of a low-dose oral estrogen only treatment on bone mineral density and quantitative ultrasonometry in postmenopausal women. <i>Gynecological Endocrinology</i> , 2012, 28, 1002-1005.	1.7	5
436	Duration of first prescribed long acting insulin therapy in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2012, 97, e3.	2.8	0
437	Longitudinal change in HbA1c after insulin initiation in primary care patients with type 2 diabetes: A database analysis in UK and Germany. <i>Primary Care Diabetes</i> , 2012, 6, 47-52.	1.8	8
438	Amputation rate and risk factors in type 2 patients with diabetic foot syndrome under real-life conditions in Germany. <i>Primary Care Diabetes</i> , 2012, 6, 241-246.	1.8	76
439	Predictors for the initiation of a basal supported oral therapy (BOT) in type 2 diabetic patients under real-life conditions in Germany. <i>Primary Care Diabetes</i> , 2012, 6, 329-335.	1.8	7
440	PCN23 Cancer Related Bisphosphonate Therapy in Different Treatment Facilities. <i>Value in Health</i> , 2012, 15, A412.	0.3	0
441	PCN106 Persistence in Patients With Breast Cancer Treated With Tamoxifen or Aromatase Inhibitors: Results of a Retrospective Cohort Study. <i>Value in Health</i> , 2012, 15, A428.	0.3	1
442	Risk of breast cancer in patients on long-acting insulin analogues in comparison with those on human insulin. <i>Diabetologia</i> , 2012, 55, 1554-1555.	6.3	9
443	Unlicensed use of metformin in children and adolescents in Germany and France. <i>British Journal of Clinical Pharmacology</i> , 2012, 73, 307-308.	2.4	8
444	Topical therapy of hand eczema — analysis of the prescription profile from dermatologists in private practice. <i>JDDG - Journal of the German Society of Dermatology</i> , 2012, 10, 180-184.	0.8	8
445	GRAND: the German retrospective cohort analysis on compliance and persistence and the associated risk of fractures in osteoporotic women treated with oral bisphosphonates. <i>Osteoporosis International</i> , 2012, 23, 223-231.	3.1	114
446	Antihypertensive treatment and risk of dementia: a retrospective database study. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 195-201.	0.6	28
447	Persistence and compliance of medications used in the treatment of osteoporosis — analysis using a large scale, representative, longitudinal German database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 315-322.	0.6	68
448	Micro- and macrovascular outcomes in Type 2 diabetic patients treated with insulin glulisine or human regular insulin: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 821-829.	0.6	5
449	A retrospective database analysis on persistence with inhaled corticosteroid therapy: comparison of two dry powder inhalers during asthma treatment in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012, 50, 257-264.	0.6	8
450	Basal Supported Oral Therapy with Insulin Glargine Results in Longer Persistence and Lower Costs Compared with Insulin Detemir in Type 2 Diabetics in Germany. <i>Health Outcomes Research in Medicine</i> , 2011, 2, e39-e50.	0.6	4

#	ARTICLE	IF	CITATIONS
451	PCV104 Persistence in Hypertension Treatment with Olmesartan Medoxomil Versus Valsartan - Analysis of Real-Life Prescription Data in Germany. Value in Health, 2011, 14, A383.	0.3	2
452	PDB5 Micro- and Macrovascular Outcomes in Primary Care Patients with Type 2 Diabetes Treated with Insulin Glulisine or Human Regular Insulin: A Retrospective German Database Analysis. Value in Health, 2011, 14, A472.	0.3	0
453	Rotavirus Vaccination in Germany. Pediatric Infectious Disease Journal, 2011, 30, e244-e247.	2.0	15
454	Drug utilization patterns and reported health status in ethnic German migrants (Aussiedler) in Germany: a cross-sectional study. BMC Public Health, 2011, 11, 509.	2.9	14
455	The negative impact of rebate contracts on the health care of patients with depression in Germany. International Journal of Clinical Pharmacology and Therapeutics, 2011, 49, 397-402.	0.6	3
456	Versorgungsprofil von Patienten mit schweren allergischen Reaktionen. Allergologie, 2011, 34, 285-293.	0.1	1
457	Characteristics of diabetic patients visiting a podiatry practice in Germany. Journal of Wound Care, 2010, 19, 140-148.	1.2	22
458	Different persistence on initial basal supported oral therapy in Type 2 diabetics is associated with unequal distributions of insulin treatment regimens under real-life conditions in Germany. International Journal of Clinical Pharmacology and Therapeutics, 2010, 48, 761-766.	0.6	4
459	Prevalence of diabetic foot syndrome and its risk factors in the UK. Journal of Wound Care, 2010, 19, 333-337.	1.2	74
460	PDB10 PREVALENCE AND OUTCOME OF INSULIN ANALOGUE TREATMENT COMPARING GERMANY, UK, AND FRANCE. Value in Health, 2010, 13, A285.	0.3	0
461	PDB16 INSULIN GLARGINE IS ASSOCIATED WITH A LOWER INCIDENCE OF DIABETIC FOOT SYNDROME AND MACROVASCULAR COMPLICATIONS COMPARED TO NPH INSULIN IN TYPE 2 DIABETICS UNDER GERMAN REAL-LIFE CONDITIONS. Value in Health, 2010, 13, A286.	0.3	0
462	PDB55 COMPARISON OF RESOURCE USE AND COSTS IN TYPE 1 DIABETES PATIENTS TREATED WITH DIFFERENT LONG ACTING INSULINS IN A BASAL-BOLUS REGIMEN IN GERMANY. Value in Health, 2010, 13, A294.	0.3	0
463	Relation of the first hypertension-associated event with medication, compliance and persistence in naïve hypertensive patients after initiating monotherapy. International Journal of Clinical Pharmacology and Therapeutics, 2010, 48, 173-183.	0.6	13
464	Longitudinal Change in HbA1c in Patients with Type 2 Diabetes: Comparison of Short-acting, Biphasic and Basal Insulin. Experimental and Clinical Endocrinology and Diabetes, 2009, 117, 230-233.	1.2	2
465	PDB11 DATABASE ANALYSIS ON PREVALENCE AND RISK FACTORS OF DIABETIC FOOT SYNDROME (DFS) IN GERMANY. Value in Health, 2009, 12, A403.	0.3	0
466	PMS66 GRAND: THE GERMAN RETROSPECTIVE COHORT ANALYSIS ON NON-ADHERENCE AND ASSOCIATED RISK OF FRACTURES IN OSTEOPOROSIS PATIENTS TREATED WITH ORAL BISPHOSPHONATES. Value in Health, 2009, 12, A445.	0.3	0
467	PMS68 GRAND: THE GERMAN RETROSPECTIVE COHORT ANALYSIS ON NON-ADHERENCE IN OSTEOPOROSIS PATIENTS TREATED WITH ORAL BISPHOSPHONATES. Value in Health, 2009, 12, A446.	0.3	0
468	Validity and representativeness of the "Disease Analyzer" patient database for use in pharmaco-epidemiological and pharmaco-economic studies. International Journal of Clinical Pharmacology and Therapeutics, 2009, 47, 617-626.	0.6	237

#	ARTICLE	IF	CITATIONS
469	Antiemetic Medications in Children with Presumed Infectious Gastroenteritisâ€”Pharmacoepidemiology in Europe and Northern America. <i>Journal of Pediatrics</i> , 2008, 153, 659-662.e3.	1.8	35
470	PCV108 RISK AND COSTS OF THE FIRST HYPERTENSION-ASSOCIATED EVENT, COMPLIANCE AND PERSISTENCE IN NAÏVE HYPERTENSIVE PATIENTS AFTER INITIATING MONOTHERAPY. <i>Value in Health</i> , 2008, 11, A415.	0.3	4
471	PDB43 ADDING INSULIN GLARGINE TO ORALTHERAPY IN PATIENTS WITHTYPE 2 DIABETES RESULTS IN LONGER PERSISTENCE WITH TREATMENT COMPARED TO NPH INSULIN. <i>Value in Health</i> , 2008, 11, A509-A510.	0.3	0
472	Prescription of Insulin Glargine in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007, 115, 252-256.	1.2	9
473	Glycemic Durability of Monotherapy for Diabetes. <i>New England Journal of Medicine</i> , 2007, 356, 1378-1380.	27.0	11
474	Persistence with antihypertensive treatments: results of a 3-year follow-up cohort study. <i>European Journal of Clinical Pharmacology</i> , 2007, 63, 1055-1061.	1.9	61
475	Isopropanolic black cohosh extract and recurrence-free survival after breast cancer. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2007, 45, 143-154.	0.6	60
476	Pulmonary hypertension is associated with an increased incidence of cancer diagnoses. <i>Pulmonary Circulation</i> , 0, , .	1.7	2
477	The role of private prescriptions in benzodiazepine and Z-drug useâ€”a secondary analysis of office-based prescription data. <i>Deutsches A&amp;#x0308;rztblatt International</i> , 0, , .	0.9	4