

Bukola Fawole

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3416092/publications.pdf>

Version: 2024-02-01

35
papers

2,541
citations

394421

19
h-index

377865

34
g-index

35
all docs

35
docs citations

35
times ranked

2591
citing authors

#	ARTICLE	IF	CITATIONS
1	Effect of early tranexamic acid administration on mortality, hysterectomy, and other morbidities in women with post-partum haemorrhage (WOMAN): an international, randomised, double-blind, placebo-controlled trial. <i>Lancet, The</i> , 2017, 389, 2105-2116.	13.7	974
2	How women are treated during facility-based childbirth in four countries: a cross-sectional study with labour observations and community-based surveys. <i>Lancet, The</i> , 2019, 394, 1750-1763.	13.7	279
3	Effect of treatment delay on the effectiveness and safety of antifibrinolytics in acute severe haemorrhage: a meta-analysis of individual patient-level data from 40â€™138 bleeding patients. <i>Lancet, The</i> , 2018, 391, 125-132.	13.7	261
4	Mistreatment of women during childbirth in Abuja, Nigeria: a qualitative study on perceptions and experiences of women and healthcare providers. <i>Reproductive Health</i> , 2017, 14, 9.	3.1	129
5	Biophysical profile for fetal assessment in high risk pregnancies. <i>The Cochrane Library</i> , 2008, , CD000038.	2.8	101
6	â€œ By slapping their laps, the patient will know that you truly care for her â€ A qualitative study on social norms and acceptability of the mistreatment of women during childbirth in Abuja, Nigeria. <i>SSM - Population Health</i> , 2016, 2, 640-655.	2.7	97
7	Methodological development of tools to measure how women are treated during facility-based childbirth in four countries: labor observation and community survey. <i>BMC Medical Research Methodology</i> , 2018, 18, 132.	3.1	70
8	The global maternal sepsis study and awareness campaign (GLOSS): study protocol. <i>Reproductive Health</i> , 2018, 15, 16.	3.1	62
9	Progression of the first stage of spontaneous labour: A prospective cohort study in two sub-Saharan African countries. <i>PLoS Medicine</i> , 2018, 15, e1002492.	8.4	57
10	Risk factors for peripartum hysterectomy among women with postpartum haemorrhage: analysis of data from the WOMAN trial. <i>BMC Pregnancy and Childbirth</i> , 2018, 18, 186.	2.4	51
11	Antenatal corticosteroids for women at risk of imminent preterm birth in low-resource countries: the case for equipoise and the need for efficacy trials. <i>BMJ Global Health</i> , 2017, 2, e000398.	4.7	44
12	Defining quality of care during childbirth from the perspectives of Nigerian and Ugandan women: A qualitative study. <i>International Journal of Gynecology and Obstetrics</i> , 2017, 139, 4-16.	2.3	42
13	Tranexamic acid for the prevention of postpartum bleeding in women with anaemia: study protocol for an international, randomised, double-blind, placebo-controlled trial. <i>Trials</i> , 2018, 19, 712.	1.6	39
14	Room temperature stable carbetocin for the prevention of postpartum haemorrhage during the third stage of labour in women delivering vaginally: study protocol for a randomized controlled trial. <i>Trials</i> , 2016, 17, 143.	1.6	31
15	The communication and emotional support needs to improve women's experience of childbirth care in health facilities in Southwest Nigeria: A qualitative study. <i>International Journal of Gynecology and Obstetrics</i> , 2017, 139, 27-37.	2.3	28
16	Haematological and fibrinolytic status of Nigerian women with post-partum haemorrhage. <i>BMC Pregnancy and Childbirth</i> , 2018, 18, 143.	2.4	28
17	Tranexamic acid for significant traumatic brain injury (The CRASH-3 trial): Statistical analysis plan for an international, randomised, double-blind, placebo-controlled trial. <i>Wellcome Open Research</i> , 2018, 3, 86.	1.8	28
18	Effect of tranexamic acid on coagulation and fibrinolysis in women with postpartum haemorrhage (WOMAN-ETAC): a single-centre, randomised, double-blind, placebo-controlled trial. <i>Wellcome Open Research</i> , 2018, 3, 100.	1.8	24

#	ARTICLE	IF	CITATIONS
19	WHO Better Outcomes in Labour Difficulty (BOLD) project: innovating to improve quality of care around the time of childbirth. <i>Reproductive Health</i> , 2015, 12, 48.	3.1	22
20	Understanding abortion-related complications in health facilities: results from WHO multicountry survey on abortion (MCS-A) across 11 sub-Saharan African countries. <i>BMJ Global Health</i> , 2021, 6, e003702.	4.7	22
21	Maternal oxygen administration for fetal distress. <i>The Cochrane Library</i> , 2012, 12, CD000136.	2.8	21
22	Formative research and development of innovative tools for "Better Outcomes in Labour Difficulty" (BOLD): study protocol. <i>Reproductive Health</i> , 2015, 12, 50.	3.1	20
23	Trends in misoprostol use and abortion complications: A cross-sectional study from nine referral hospitals in Nigeria. <i>PLoS ONE</i> , 2018, 13, e0209415.	2.5	14
24	Tranexamic acid for significant traumatic brain injury (The CRASH-3 trial): Statistical analysis plan for an international, randomised, double-blind, placebo-controlled trial. <i>Wellcome Open Research</i> , 2018, 3, 86.	1.8	14
25	Narratives of women presenting with abortion complications in Southwestern Nigeria: A qualitative study. <i>PLoS ONE</i> , 2019, 14, e0217616.	2.5	13
26	Using a service design model to develop the "Passport to Safer Birth" in Nigeria and Uganda. <i>International Journal of Gynecology and Obstetrics</i> , 2017, 139, 56-66.	2.3	12
27	Negotiating quality standards for effective delivery of labor and childbirth care in Nigeria and Uganda. <i>International Journal of Gynecology and Obstetrics</i> , 2017, 139, 47-55.	2.3	11
28	Healthcare providers' perspectives on labor monitoring in Nigeria and Uganda: A qualitative study on challenges and opportunities. <i>International Journal of Gynecology and Obstetrics</i> , 2017, 139, 17-26.	2.3	11
29	The impact of early outcome events on the effect of tranexamic acid in post-partum haemorrhage: an exploratory subgroup analysis of the WOMAN trial. <i>BMC Pregnancy and Childbirth</i> , 2018, 18, 215.	2.4	9
30	Secondary analysis of the WOMAN trial to explore the risk of sepsis after invasive treatments for postpartum hemorrhage. <i>International Journal of Gynecology and Obstetrics</i> , 2019, 146, 231-237.	2.3	8
31	Effect of tranexamic acid on coagulation and fibrinolysis in women with postpartum haemorrhage (WOMAN-ETAC): protocol and statistical analysis plan for a randomized controlled trial. <i>Wellcome Open Research</i> , 2016, 1, 31.	1.8	8
32	Early evaluation of the "STOP SEPSIS" WHO Global Maternal Sepsis Awareness Campaign implemented for healthcare providers in 46 low, middle and high-income countries. <i>BMJ Open</i> , 2020, 10, e036338.	1.9	6
33	A prospective study of severity of complications associated with use of misoprostol and other abortion methods in South West Nigeria. <i>International Journal of Gynecology and Obstetrics</i> , 2019, 146, 302-307.	2.3	3
34	Use of histograms to assess the efficacy of uterotonic treatment for post-partum haemorrhage: A feasibility study. <i>Best Practice and Research in Clinical Obstetrics and Gynaecology</i> , 2019, 61, 15-27.	2.8	2
35	The Log-Normal zero-inflated cure regression model for labor time in an African obstetric population. <i>Journal of Applied Statistics</i> , 0, , 1-14.	1.3	0