

Hirohito Sone

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3421735/publications.pdf>

Version: 2024-02-01

453
papers

17,350
citations

20817

60
h-index

20358

116
g-index

475
all docs

475
docs citations

475
times ranked

23734
citing authors

#	ARTICLE	IF	CITATIONS
1	Cardiorespiratory Fitness as a Quantitative Predictor of All-Cause Mortality and Cardiovascular Events in Healthy Men and Women. <i>JAMA - Journal of the American Medical Association</i> , 2009, 301, 2024.	7.4	2,357
2	Effect of Aerobic Exercise Training on Serum Levels of High-Density Lipoprotein Cholesterol. <i>Archives of Internal Medicine</i> , 2007, 167, 999.	3.8	471
3	Crucial role of a long-chain fatty acid elongase, Elovl6, in obesity-induced insulin resistance. <i>Nature Medicine</i> , 2007, 13, 1193-1202.	30.7	459
4	Association Between Serum Uric Acid and Development of Type 2 Diabetes. <i>Diabetes Care</i> , 2009, 32, 1737-1742.	8.6	415
5	Low-density lipoprotein receptor-related protein 5 (LRP5) is essential for normal cholesterol metabolism and glucose-induced insulin secretion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003, 100, 229-234.	7.1	382
6	Polyunsaturated Fatty Acids Suppress Sterol Regulatory Element-binding Protein 1c Promoter Activity by Inhibition of Liver X Receptor (LXR) Binding to LXR Response Elements. <i>Journal of Biological Chemistry</i> , 2002, 277, 1705-1711.	3.4	347
7	SREBPs suppress IRS-2-mediated insulin signalling in the liver. <i>Nature Cell Biology</i> , 2004, 6, 351-357.	10.3	305
8	Cross-Talk between Peroxisome Proliferator-Activated Receptor (PPAR) α and Liver X Receptor (LXR) in Nutritional Regulation of Fatty Acid Metabolism. I. PPARs Suppress Sterol Regulatory Element Binding Protein-1c Promoter through Inhibition of LXR Signaling. <i>Molecular Endocrinology</i> , 2003, 17, 1240-1254.	3.7	264
9	Alcohol Consumption and Risk of Atrial Fibrillation. <i>Journal of the American College of Cardiology</i> , 2011, 57, 427-436.	2.8	248
10	Dual regulation of mouse Delta(5)- and Delta(6)-desaturase gene expression by SREBP-1 and PPARalpha. <i>Journal of Lipid Research</i> , 2002, 43, 107-114.	4.2	220
11	HbA1c 5.7-6.4% and impaired fasting plasma glucose for diagnosis of prediabetes and risk of progression to diabetes in Japan (TOPICS 3): a longitudinal cohort study. <i>Lancet, The</i> , 2011, 378, 147-155.	13.7	212
12	Severe Hypercholesterolemia, Hypertriglyceridemia, and Atherosclerosis in Mice Lacking Both Leptin and the Low Density Lipoprotein Receptor. <i>Journal of Biological Chemistry</i> , 2001, 276, 37402-37408.	3.4	194
13	Diabetes and Risk of Hearing Impairment in Adults: A Meta-Analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 51-58.	3.6	194
14	Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: A meta-analysis. <i>Preventive Medicine</i> , 2011, 53, 260-267.	3.4	189
15	Sterol Regulatory Element-Binding Proteins Induce an Entire Pathway of Cholesterol Synthesis. <i>Biochemical and Biophysical Research Communications</i> , 2001, 286, 176-183.	2.1	187
16	Comparisons of the Strength of Associations With Future Type 2 Diabetes Risk Among Anthropometric Obesity Indicators, Including Waist-to-Height Ratio: A Meta-Analysis. <i>American Journal of Epidemiology</i> , 2012, 176, 959-969.	3.4	181
17	Cross-Talk between Peroxisome Proliferator-Activated Receptor (PPAR) α and Liver X Receptor (LXR) in Nutritional Regulation of Fatty Acid Metabolism. II. LXRs Suppress Lipid Degradation Gene Promoters through Inhibition of PPAR Signaling. <i>Molecular Endocrinology</i> , 2003, 17, 1255-1267.	3.7	177
18	Obesity and type 2 diabetes in Japanese patients. <i>Lancet, The</i> , 2003, 361, 85.	13.7	175

#	ARTICLE	IF	CITATIONS
19	Cloning and characterization of a mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. <i>Journal of Lipid Research</i> , 2002, 43, 911-920.	4.2	172
20	Serum Level of Triglycerides Is a Potent Risk Factor Comparable to LDL Cholesterol for Coronary Heart Disease in Japanese Patients with Type 2 Diabetes: Subanalysis of the Japan Diabetes Complications Study (JDCS). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, 3448-3456.	3.6	170
21	TFE3 transcriptionally activates hepatic IRS-2, participates in insulin signaling and ameliorates diabetes. <i>Nature Medicine</i> , 2006, 12, 107-113.	30.7	168
22	Insulin-Independent Induction of Sterol Regulatory Element-Binding Protein-1c Expression in the Livers of Streptozotocin-Treated Mice. <i>Diabetes</i> , 2004, 53, 560-569.	0.6	167
23	Prevalence of albuminuria and renal insufficiency and associated clinical factors in type 2 diabetes: the Japan Diabetes Clinical Data Management study (JDDM15). <i>Nephrology Dialysis Transplantation</i> , 2008, 24, 1212-1219.	0.7	159
24	Association Between Physical Activity and Risk of All-Cause Mortality and Cardiovascular Disease in Patients With Diabetes. <i>Diabetes Care</i> , 2013, 36, 471-479.	8.6	156
25	Influence of Fat and Carbohydrate Proportions on the Metabolic Profile in Patients With Type 2 Diabetes: A Meta-Analysis. <i>Diabetes Care</i> , 2009, 32, 959-965.	8.6	144
26	Elovl6 promotes nonalcoholic steatohepatitis. <i>Hepatology</i> , 2012, 56, 2199-2208.	7.3	144
27	Is the Diagnosis of Metabolic Syndrome Useful for Predicting Cardiovascular Disease in Asian Diabetic Patients?. <i>Diabetes Care</i> , 2005, 28, 1463-1471.	8.6	141
28	SREBP-1 Interacts with Hepatocyte Nuclear Factor-4 α and Interferes with PGC-1 Recruitment to Suppress Hepatic Gluconeogenic Genes. <i>Journal of Biological Chemistry</i> , 2004, 279, 12027-12035.	3.4	134
29	Cloning and characterization of a mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. <i>Journal of Lipid Research</i> , 2002, 43, 911-20.	4.2	133
30	Long-term lifestyle intervention lowers the incidence of stroke in Japanese patients with type 2 diabetes: a nationwide multicentre randomised controlled trial (the Japan Diabetes Complications) <i>Tj ETQq0 0 0 rgBT/Overlock2.0 Tf 50</i>		
31	Ocular vascular endothelial growth factor levels in diabetic rats are elevated before observable retinal proliferative changes. <i>Diabetologia</i> , 1997, 40, 726-730.	6.3	128
32	Effect of web-based lifestyle modification on weight control: a meta-analysis. <i>International Journal of Obesity</i> , 2012, 36, 675-685.	3.4	120
33	Impact of population aging on trends in diabetes prevalence: A meta-analysis of 160,000 Japanese adults. <i>Journal of Diabetes Investigation</i> , 2015, 6, 533-542.	2.4	111
34	Neutralization of Vascular Endothelial Growth Factor Prevents Collagen-Induced Arthritis and Ameliorates Established Disease in Mice. <i>Biochemical and Biophysical Research Communications</i> , 2001, 281, 562-568.	2.1	108
35	Vascular Endothelial Growth Factor Is Induced by Long-Term High Glucose Concentration and Up-Regulated by Acute Glucose Deprivation in Cultured Bovine Retinal Pigmented Epithelial Cells. <i>Biochemical and Biophysical Research Communications</i> , 1996, 221, 193-198.	2.1	104
36	Quantitative relationship between body weight gain in adulthood and incident type 2 diabetes: a meta-analysis. <i>Obesity Reviews</i> , 2014, 15, 202-214.	6.5	99

#	ARTICLE	IF	CITATIONS
37	Effects of Lifestyle Modifications on Patients with Type 2 Diabetes: The Japan Diabetes Complications Study (JDCS) Study Design, Baseline Analysis and Three Year-Interim Report. <i>Hormone and Metabolic Research</i> , 2002, 34, 509-515.	1.5	98
38	p57Kip2 Regulates Actin Dynamics by Binding and Translocating LIM-kinase 1 to the Nucleus. <i>Journal of Biological Chemistry</i> , 2003, 278, 52919-52923.	3.4	96
39	Predicting Macro- and Microvascular Complications in Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 1193-1199.	8.6	96
40	Protein Kinase A Suppresses Sterol Regulatory Element-binding Protein-1C Expression via Phosphorylation of Liver X Receptor in the Liver. <i>Journal of Biological Chemistry</i> , 2007, 282, 11687-11695.	3.4	93
41	Incidence and progression of diabetic retinopathy in Japanese adults with type 2 diabetes: 8-year follow-up study of the Japan Diabetes Complications Study (JDCS). <i>Diabetologia</i> , 2011, 54, 2288-2294.	6.3	90
42	HbA1c variability and the development of microalbuminuria in type 2 diabetes: Tsukuba Kawai Diabetes Registry 2. <i>Diabetologia</i> , 2012, 55, 2128-2131.	6.3	88
43	The hyperglycemia stimulated myocardial endoplasmic reticulum (ER) stress contributes to diabetic cardiomyopathy in the transgenic non-obese type 2 diabetic rats: A differential role of unfolded protein response (UPR) signaling proteins. <i>International Journal of Biochemistry and Cell Biology</i> , 2013, 45, 438-447.	2.8	88
44	Effect of Postmenopausal Status and Age at Menopause on Type 2 Diabetes and Prediabetes in Japanese Individuals: Toranomon Hospital Health Management Center Study 17 (TOPICS 17). <i>Diabetes Care</i> , 2013, 36, 4007-4014.	8.6	88
45	Astaxanthin suppresses scavenger receptor expression and matrix metalloproteinase activity in macrophages. <i>European Journal of Nutrition</i> , 2010, 49, 119-126.	3.9	86
46	Refractive changes in diabetic patients during intensive glycaemic control. <i>British Journal of Ophthalmology</i> , 2000, 84, 1097-1102.	3.9	85
47	Palmitate Impairs and Eicosapentaenoate Restores Insulin Secretion Through Regulation of SREBP-1c in Pancreatic Islets. <i>Diabetes</i> , 2008, 57, 2382-2392.	0.6	84
48	Diagnosis and Management of Type I and Type V Hyperlipoproteinemia. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 1-12.	2.0	81
49	Clinicopathological, Cytogenetic, and Prognostic Analysis of 131 Myeloid Sarcoma Patients. <i>American Journal of Surgical Pathology</i> , 2016, 40, 1473-1483.	3.7	81
50	Risk of Cardiovascular Diseases Is Increased Even with Mild Diabetic Retinopathy. <i>Ophthalmology</i> , 2013, 120, 574-582.	5.2	79
51	Transgenic Mice Overexpressing Nuclear SREBP-1c in Pancreatic β -Cells. <i>Diabetes</i> , 2005, 54, 492-499.	0.6	78
52	Dietary Sodium Intake and Incidence of Diabetes Complications in Japanese Patients with Type 2 Diabetes: Analysis of the Japan Diabetes Complications Study (JDCS). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 3635-3643.	3.6	76
53	Acetyl-coenzyme A synthetase is a lipogenic enzyme controlled by SREBP-1 and energy status. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2002, 282, E222-E230.	3.5	74
54	Genetic and physiological analysis of branched-chain alcohols and isoamyl acetate production in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2002, 59, 501-508.	3.6	73

#	ARTICLE	IF	CITATIONS
55	Fruit Intake and Incident Diabetic Retinopathy with Type 2 Diabetes. <i>Epidemiology</i> , 2013, 24, 204-211.	2.7	71
56	Insulin Inhibits Apoptosis of Macrophage Cell Line, THP-1 Cells, via Phosphatidylinositol-3-Kinase-Dependent Pathway. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2002, 22, 380-386.	2.4	67
57	Determinants of Decline in Glomerular Filtration Rate in Nonproteinuric Subjects with or without Diabetes and Hypertension. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009, 4, 1432-1440.	4.5	67
58	Metabolically Healthy Obesity, Presence or Absence of Fatty Liver, and Risk of Type 2 Diabetes in Japanese Individuals: Toranomon Hospital Health Management Center Study 20 (TOPICS 20). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, 2952-2960.	3.6	66
59	In Search of the Ideal Resistance Training Program to Improve Glycemic Control and its Indication for Patients with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. <i>Sports Medicine</i> , 2016, 46, 67-77.	6.5	66
60	Long-term multiple risk factor interventions in Japanese elderly diabetic patients: The Japanese Elderly Diabetes Intervention Trial study design, baseline characteristics and effects of intervention. <i>Geriatrics and Gerontology International</i> , 2012, 12, 7-17.	1.5	65
61	Increase in serum vascular endothelial growth factor levels during altitude training. <i>Acta Physiologica Scandinavica</i> , 1998, 162, 455-459.	2.2	61
62	Enhancement of glucose transport by vascular endothelial growth factor in retinal endothelial cells. <i>Investigative Ophthalmology and Visual Science</i> , 2000, 41, 1876-84.	3.3	61
63	Mouse MafA, homologue of zebrafish somite Maf 1, contributes to the specific transcriptional activity through the insulin promoter. <i>Biochemical and Biophysical Research Communications</i> , 2003, 312, 831-842.	2.1	60
64	Granuphilin is activated by SREBP-1c and involved in impaired insulin secretion in diabetic mice. <i>Cell Metabolism</i> , 2006, 4, 143-154.	16.2	60
65	Low transition rate from normo- and low microalbuminuria to proteinuria in Japanese type 2 diabetic individuals: the Japan Diabetes Complications Study (JDCS). <i>Diabetologia</i> , 2011, 54, 1025-1031.	6.3	60
66	Skeletal muscle-specific HMG-CoA reductase knockout mice exhibit rhabdomyolysis: A model for statin-induced myopathy. <i>Biochemical and Biophysical Research Communications</i> , 2015, 466, 536-540.	2.1	59
67	Statins downregulate ATP-binding-cassette transporter A1 gene expression in macrophages. <i>Biochemical and Biophysical Research Communications</i> , 2004, 316, 790-794.	2.1	57
68	Energy intake and obesity in Japanese patients with type 2 diabetes. <i>Lancet, The</i> , 2004, 363, 248-249.	13.7	57
69	Elevated levels of vascular endothelial growth factor in the sera of patients with rheumatoid arthritis Correlation with disease activity. <i>Life Sciences</i> , 2001, 69, 1861-1869.	4.3	56
70	Insulin Up-regulates Tumor Necrosis Factor- α Production in Macrophages through an Extracellular-regulated Kinase-dependent Pathway. <i>Journal of Biological Chemistry</i> , 2001, 276, 32531-32537.	3.4	56
71	Involvement of glomerular SREBP-1c in diabetic nephropathy. <i>Biochemical and Biophysical Research Communications</i> , 2007, 364, 502-508.	2.1	56
72	Long-term effects of eicosapentaenoic acid on diabetic peripheral neuropathy and serum lipids in patients with type II diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 1996, 10, 280-287.	2.3	55

#	ARTICLE	IF	CITATIONS
73	Hypoxia and endothelin-1 induce VEGF production in human vascular smooth muscle cells. <i>Life Sciences</i> , 1998, 63, 477-484.	4.3	55
74	Lipid Synthetic Transcription Factor SREBP-1a Activates p21WAF1/CIP1, a Universal Cyclin-Dependent Kinase Inhibitor. <i>Molecular and Cellular Biology</i> , 2005, 25, 8938-8947.	2.3	55
75	Effect of thiazolidinediones and metformin on LDL oxidation and aortic endothelium relaxation in diabetic GK rats. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2003, 284, E1125-E1130.	3.5	54
76	Screening for pre-diabetes to predict future diabetes using various cutoff points for HbA _{1c} and impaired fasting glucose: the Toranomon Hospital Health Management Center Study 4 (TOPICS 4). <i>Diabetic Medicine</i> , 2012, 29, e279-85.	2.3	54
77	The effect of exercise training on adiponectin receptor expression in KK ^Y obese/diabetic mice. <i>Journal of Endocrinology</i> , 2006, 189, 643-653.	2.6	52
78	Leisure-time physical activity is a significant predictor of stroke and total mortality in Japanese patients with type 2 diabetes: analysis from the Japan Diabetes Complications Study (JDCS). <i>Diabetologia</i> , 2013, 56, 1021-1030.	6.3	51
79	Different Effects of Eicosapentaenoic and Docosahexaenoic Acids on Atherogenic High-Fat Diet-Induced Non-Alcoholic Fatty Liver Disease in Mice. <i>PLoS ONE</i> , 2016, 11, e0157580.	2.5	50
80	Diabetic Retinopathy and Microalbuminuria Can Predict Macroalbuminuria and Renal Function Decline in Japanese Type 2 Diabetic Patients. <i>Diabetes Care</i> , 2013, 36, 2803-2809.	8.6	49
81	Hepatic CREB3L3 Controls Whole-Body Energy Homeostasis and Improves Obesity and Diabetes. <i>Endocrinology</i> , 2014, 155, 4706-4719.	2.8	49
82	Potential protective effect of lactation against incidence of type 2 diabetes mellitus in women with previous gestational diabetes mellitus: A systematic review and meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2017, 33, e2875.	4.0	49
83	Antioxidants and an inhibitor of advanced glycation ameliorate death of retinal microvascular cells in diabetic retinopathy. <i>Diabetes/Metabolism Research and Reviews</i> , 2006, 22, 38-45.	4.0	48
84	Sterol Regulatory Element-Binding Protein-1 Determines Plasma Remnant Lipoproteins and Accelerates Atherosclerosis in Low-Density Lipoprotein Receptor-Deficient Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 1788-1795.	2.4	46
85	A possible link between BDNF and mTOR in control of food intake. <i>Frontiers in Psychology</i> , 2014, 5, 1093.	2.1	46
86	Vascular Endothelial Growth Factor Level in Aqueous Humor of Diabetic Patients With Rubeotic Glaucoma Is Markedly Elevated. <i>Diabetes Care</i> , 1996, 19, 1306-1307.	8.6	45
87	Comparison of Various Lipid Variables as Predictors of Coronary Heart Disease in Japanese Men and Women With Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 1150-1157.	8.6	45
88	Longitudinal Trajectories of HbA _{1c} and Fasting Plasma Glucose Levels During the Development of Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 1050-1052.	8.6	45
89	CREB3L3 controls fatty acid oxidation and ketogenesis in synergy with PPAR α . <i>Scientific Reports</i> , 2016, 6, 39182.	3.3	45
90	Physiological changes in circulating mannose levels in normal, glucose-intolerant, and diabetic subjects. <i>Metabolism: Clinical and Experimental</i> , 2003, 52, 1019-1027.	3.4	44

#	ARTICLE	IF	CITATIONS
91	Meta-Analysis of the Quantitative Relation Between Pulse Pressure and Mean Arterial Pressure and Cardiovascular Risk in Patients With Diabetes Mellitus. <i>American Journal of Cardiology</i> , 2014, 113, 1058-1065.	1.6	44
92	A distinct subtype of Epstein-Barr virus-positive T/NK-cell lymphoproliferative disorder: adult patients with chronic active Epstein-Barr virus infection-like features. <i>Haematologica</i> , 2018, 103, 1018-1028.	3.5	44
93	Hepatocyte ELOVL Fatty Acid Elongase 6 Determines Ceramide Acyl-Chain Length and Hepatic Insulin Sensitivity in Mice. <i>Hepatology</i> , 2020, 71, 1609-1625.	7.3	44
94	Isolation and characterization of the ATF2 gene encoding alcohol acetyltransferase II in the bottom fermenting yeast <i>Saccharomyces pastorianus</i> . <i>Yeast</i> , 1999, 15, 409-417.	1.7	43
95	HMG-CoA reductase inhibitor decreases small dense low-density lipoprotein and remnant-like particle cholesterol in patients with type-2 diabetes. <i>Life Sciences</i> , 2002, 71, 2403-2412.	4.3	43
96	Development of a new scoring system for predicting the 5-year incidence of type 2 diabetes in Japan: the Toranomon Hospital Health Management Center Study 6 (TOPICS 6). <i>Diabetologia</i> , 2012, 55, 3213-3223.	6.3	43
97	Molecular understanding of curcumin in diabetic nephropathy. <i>Drug Discovery Today</i> , 2013, 18, 756-763.	6.4	43
98	Internal deletion of BCOR reveals a tumor suppressor function for BCOR in T lymphocyte malignancies. <i>Journal of Experimental Medicine</i> , 2017, 214, 2901-2913.	8.5	43
99	Chronic kidney disease categories and renal cardiovascular outcomes in type 2 diabetes without prevalent cardiovascular disease: a prospective cohort study (JDDM25). <i>Diabetologia</i> , 2012, 55, 1911-1918.	6.3	42
100	Cohort Profile: The Japan Diabetes Complications Study: a long-term follow-up of a randomised lifestyle intervention study of type 2 diabetes. <i>International Journal of Epidemiology</i> , 2014, 43, 1054-1062.	1.9	42
101	Stability and changes in metabolically healthy overweight or obesity and risk of future diabetes: Niigata wellness study. <i>Obesity</i> , 2014, 22, 2420-2425.	3.0	41
102	Accuracy of Japanese claims data in identifying diabetes-related complications. <i>Pharmacoepidemiology and Drug Safety</i> , 2021, 30, 594-601.	1.9	41
103	Effects of intraocular or systemic administration of neutralizing antibody against vascular endothelial growth factor on the murine experimental model of retinopathy. <i>Life Sciences</i> , 1999, 65, 2573-2580.	4.3	40
104	Non-high-density lipoprotein cholesterol: An important predictor of stroke and diabetes-related mortality in Japanese elderly diabetic patients. <i>Geriatrics and Gerontology International</i> , 2012, 12, 18-28.	1.5	40
105	Association Between Remission of Macroalbuminuria and Preservation of Renal Function in Patients With Type 2 Diabetes With Overt Proteinuria. <i>Diabetes Care</i> , 2013, 36, 3227-3233.	8.6	40
106	Effects of magnesium on postprandial serum lipid responses in healthy human subjects. <i>British Journal of Nutrition</i> , 2010, 103, 469-472.	2.3	39
107	Intakes of Dietary Fiber, Vegetables, and Fruits and Incidence of Cardiovascular Disease in Japanese Patients With Type 2 Diabetes. <i>Diabetes Care</i> , 2013, 36, 3916-3922.	8.6	39
108	Carotid Artery Plaque and LDL-to-HDL Cholesterol Ratio Predict Atherosclerotic Status in Coronary Arteries in Asymptomatic Patients with Type 2 Diabetes Mellitus. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 452-464.	2.0	39

#	ARTICLE	IF	CITATIONS
109	Type 2 Diabetes Prevalence in Asian Subjects: Response to McNeely and Boyko. <i>Diabetes Care</i> , 2004, 27, 1251-1252.	8.6	38
110	Low incidence of cardiovascular events in Japanese patients with Type 2 diabetes in primary care settings: a prospective cohort study (JDDM 20). <i>Diabetic Medicine</i> , 2011, 28, 1221-1228.	2.3	38
111	Role of alcohol drinking pattern in type 2 diabetes in Japanese men: the Toranomon Hospital Health Management Center Study 11 (TOPICS 11). <i>American Journal of Clinical Nutrition</i> , 2013, 97, 561-568.	4.7	37
112	High risk of failing eradication of <i>Helicobacter pylori</i> in patients with diabetes: A meta-analysis. <i>Diabetes Research and Clinical Practice</i> , 2014, 106, 81-87.	2.8	37
113	Physical Fitness Tests and Type 2 Diabetes Among Japanese: A Longitudinal Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2019, 29, 139-146.	2.4	37
114	The New Worldwide Definition of Metabolic Syndrome Is Not a Better Diagnostic Predictor of Cardiovascular Disease in Japanese Diabetic Patients Than the Existing Definitions: Additional analysis from the Japan Diabetes Complications Study. <i>Diabetes Care</i> , 2006, 29, 145-147.	8.6	37
115	Low HDL Cholesterol Is Associated With the Risk of Stroke in Elderly Diabetic Individuals. <i>Diabetes Care</i> , 2009, 32, 1221-1223.	8.6	36
116	Low serum potassium levels and risk of type 2 diabetes: the Toranomon Hospital Health Management Center Study 1 (TOPICS 1). <i>Diabetologia</i> , 2011, 54, 762-766.	6.3	36
117	Impact of Psychological Stress caused by the Great East Japan Earthquake on Glycemic Control in Patients with Diabetes. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2012, 120, 560-563.	1.2	36
118	Dietary intake in Japanese patients with type 2 diabetes: Analysis from Japanese Diabetes Complications Study. <i>Journal of Diabetes Investigation</i> , 2014, 5, 176-187.	2.4	36
119	Glucose Uptake and Adenoviral Mediated GLUT1 Infection Decrease Hypoxia-induced HIF-1 α Levels in Cardiac Myocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2002, 34, 1063-1073.	1.9	35
120	MYC translocation and/or BCL 2 protein expression are associated with poor prognosis in diffuse large B-cell lymphoma. <i>Cancer Science</i> , 2016, 107, 853-861.	3.9	35
121	Reduced GFR and microalbuminuria are independently associated with prevalent cardiovascular disease in Type 2 diabetes: JDDM study 16. <i>Diabetic Medicine</i> , 2008, 25, 1426-1432.	2.3	34
122	Quality of Internet information related to the Mediterranean diet. <i>Public Health Nutrition</i> , 2012, 15, 885-893.	2.2	34
123	Even Low-Intensity and Low-Volume Exercise Training May Improve Insulin Resistance in the Elderly. <i>Internal Medicine</i> , 2007, 46, 1071-1077.	0.7	33
124	Risks for glomerular filtration rate decline in association with progression of albuminuria in type 2 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 2924-2930.	0.7	33
125	Progesterone induces vascular endothelial growth factor on retinal pigment epithelial cells in culture. <i>Life Sciences</i> , 1996, 59, 21-25.	4.3	32
126	Macrophage Elovl6 Deficiency Ameliorates Foam Cell Formation and Reduces Atherosclerosis in Low-Density Lipoprotein Receptor-Deficient Mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011, 31, 1973-1979.	2.4	32

#	ARTICLE	IF	CITATIONS
127	Cholesterol sulfate induces expression of the skin barrier protein filaggrin in normal human epidermal keratinocytes through induction of ROR γ . <i>Biochemical and Biophysical Research Communications</i> , 2012, 428, 99-104.	2.1	32
128	Intestinal CREBH overexpression prevents high-cholesterol diet-induced hypercholesterolemia by reducing Npc1l1 expression. <i>Molecular Metabolism</i> , 2016, 5, 1092-1102.	6.5	32
129	TFE3 regulates muscle metabolic gene expression, increases glycogen stores, and enhances insulin sensitivity in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012, 302, E896-E902.	3.5	31
130	Low Lung Function and Risk of Type 2 Diabetes in Japanese Men: The Toranomon Hospital Health Management Center Study 9 (TOPICS 9). <i>Mayo Clinic Proceedings</i> , 2012, 87, 853-861.	3.0	31
131	TFE3 Controls Lipid Metabolism in Adipose Tissue of Male Mice by Suppressing Lipolysis and Thermogenesis. <i>Endocrinology</i> , 2013, 154, 3577-3588.	2.8	31
132	Body Mass Index and Mortality Among Japanese Patients With Type 2 Diabetes: Pooled Analysis of the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014, 99, E2692-E2696.	3.6	31
133	Hyperlipidemia and hepatitis in liver-specific CREB3L3 knockout mice generated using a one-step CRISPR/Cas9 system. <i>Scientific Reports</i> , 2016, 6, 27857.	3.3	31
134	Components of Metabolic Syndrome and their Combinations as Predictors of Cardiovascular Disease in Japanese Patients with Type 2 Diabetes. Implications for Improved Definition. Analysis from Japan Diabetes Complications Study (JDCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2009, 16, 380-387.	2.0	31
135	Efficacy and safety in sitagliptin therapy for diabetes complicated by nonalcoholic fatty liver disease. <i>Hepatology Research</i> , 2013, 43, 1163-1168.	3.4	30
136	Role of sleep duration as a risk factor for Type 2 diabetes among adults of different ages in Japan: the Niigata Wellness Study. <i>Diabetic Medicine</i> , 2014, 31, 1363-1367.	2.3	30
137	Physical Fitness During Adolescence and Adult Mortality. <i>Epidemiology</i> , 2009, 20, 463-464.	2.7	29
138	Effect of Endurance Exercise Training on Oxidative Stress in Spontaneously Hypertensive Rats (SHR) After Emergence of Hypertension. <i>Clinical and Experimental Hypertension</i> , 2010, 32, 407-415.	1.3	29
139	High cholesterol intake is associated with elevated risk of type 2 diabetes mellitus â€“ A meta-analysis. <i>Clinical Nutrition</i> , 2014, 33, 946-950.	5.0	29
140	Elovl6 Deficiency Improves Glycemic Control in Diabetic <i>db/db</i> Mice by Expanding β -Cell Mass and Increasing Insulin Secretory Capacity. <i>Diabetes</i> , 2017, 66, 1833-1846.	0.6	29
141	Disease model: hyperinsulinemia and insulin resistance. <i>Trends in Molecular Medicine</i> , 2001, 7, 320-322.	6.7	28
142	High normal HbA _{1c} levels were associated with impaired insulin secretion without escalating insulin resistance in Japanese individuals: the Toranomon Hospital Health Management Center Study 8 (TOPICS 8). <i>Diabetic Medicine</i> , 2012, 29, 1285-1290.	2.3	28
143	Fasting glucose and HbA _{1c} levels as risk factors for the development of hypertension in Japanese individuals: Toranomon hospital health management center study 16 (TOPICS 16). <i>Journal of Human Hypertension</i> , 2015, 29, 254-259.	2.2	28
144	Risk of the development of Type 2 diabetes in relation to overall obesity, abdominal obesity and the clustering of metabolic abnormalities in Japanese individuals: does metabolically healthy overweight really exist? The Niigata Wellness Study. <i>Diabetic Medicine</i> , 2015, 32, 665-672.	2.3	28

#	ARTICLE	IF	CITATIONS
145	Radioiodinated Metaiodo-benzylguanidine Scintigraphγ for Pheochromocytoma. Hormone Research, 1996, 46, 138-142.	1.8	27
146	Ibuprofen-Related Hypoglycemia in a Patient Receiving Sulfonylurea. Annals of Internal Medicine, 2001, 134, 344.	3.9	27
147	Flow-Mediated Dilation is Associated with Microalbuminuria Independent of Cardiovascular Risk Factors in Type 2 Diabetes. Journal of Atherosclerosis and Thrombosis, 2011, 18, 744-752.	2.0	27
148	Comparison of the Framingham Risk Score, UK Prospective Diabetes Study (UKPDS) Risk Engine, Japanese Atherosclerosis Longitudinal Study-Existing Cohorts Combine (JALS-ECC) and Maximum Carotid Intima-Media Thickness for Predicting Coronary Artery Stenosis in Patients with Asymptomatic Type 2 Diabetes. Journal of Atherosclerosis and Thrombosis, 2014, 21, 799-815.	2.0	27
149	Patients with type 2 diabetes having higher glomerular filtration rate showed rapid renal function decline followed by impaired glomerular filtration rate: Japan Diabetes Complications Study. Journal of Diabetes and Its Complications, 2017, 31, 473-478.	2.3	26
150	Impact of glucose tolerance status on the development of coronary artery disease among working-age men. Diabetes and Metabolism, 2017, 43, 261-264.	2.9	26
151	Frequent expression of CD30 in extranodal NK/T-cell lymphoma: Potential therapeutic target for anti-CD30 antibody-based therapy. Hematological Oncology, 2018, 36, 166-173.	1.7	26
152	Patient-based prediction algorithm of relapse after alloHSCT for acute Leukemia and its usefulness in the decision-making process using a machine learning approach. Cancer Medicine, 2019, 8, 5058-5067.	2.8	26
153	Eicosapentaenoic Acid Suppresses Basal and Insulin-Stimulated Endothelin-1 Production in Human Endothelial Cells. Hypertension Research, 2003, 26, 655-661.	2.7	25
154	Sterol Regulatory Element-binding Proteins Activate Insulin Gene Promoter Directly and Indirectly through Synergy with BETA2/E47. Journal of Biological Chemistry, 2005, 280, 34577-34589.	3.4	25
155	Thinness in Japanese Young Women. Epidemiology, 2009, 20, 464-465.	2.7	25
156	Comparison of lipid parameters to predict cardiovascular events in Japanese mild-to-moderate hypercholesterolemic patients with and without type 2 diabetes: Subanalysis of the MEGA study. Diabetes Research and Clinical Practice, 2016, 113, 14-22.	2.8	25
157	Association between Helicobacter pylori infection, eradication and diabetes mellitus. Journal of Diabetes Investigation, 2019, 10, 1341-1346.	2.4	25
158	Diabetes mellitus and risk of new-onset and recurrent heart failure: a systematic review and meta-analysis. ESC Heart Failure, 2020, 7, 2146-2174.	3.1	25
159	Inhibition of Ubiquitin Ligase F-box and WD Repeat Domain-containing 7 (Fbw7) Causes Hepatosteatosis through Krüppel-like Factor 5 (KLF5)/Peroxisome Proliferator-activated Receptor 1 α (PPAR1 α) Pathway but Not SREBP-1c Protein in Mice*. Journal of Biological Chemistry, 2011, 286, 40835-40846.	3.4	24
160	Comparison of different aspects of BMI history to identify undiagnosed diabetes in Japanese men and women: Toranomon Hospital Health Management Center Study 12 (TOPICS) Tj ETQ 0 0 rg 01 /Overlo	2.0	24
161	Late Onset Post-Transfusion Hepatitis E Developing during Chemotherapy for Acute Promyelocytic Leukemia. Internal Medicine, 2015, 54, 657-661.	0.7	24
162	Ability of Current Machine Learning Algorithms to Predict and Detect Hypoglycemia in Patients With Diabetes Mellitus: Meta-analysis. JMIR Diabetes, 2021, 6, e22458.	1.9	24

#	ARTICLE	IF	CITATIONS
163	WGEF is a novel RhoGEF expressed in intestine, liver, heart, and kidney. <i>Biochemical and Biophysical Research Communications</i> , 2004, 324, 1053-1058.	2.1	23
164	Outcome of One-year of Specialist Care of Patients with Type 2 Diabetes: A Multi-Center Prospective Survey (JDDM 2). <i>Internal Medicine</i> , 2006, 45, 589-597.	0.7	23
165	Cross-sectional association between BMI, glycemic control and energy intake in Japanese patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2007, 77, S23-S29.	2.8	22
166	Association of <i>Helicobacter pylori</i> Infection with Glycemic Control in Patients with Diabetes: A Meta-Analysis. <i>Journal of Diabetes Research</i> , 2014, 2014, 1-7.	2.3	22
167	Fulminant Type 1 Diabetes Mellitus Associated with Coxsackie Virus Type A2 Infection: A Case Report and Literature Review. <i>Internal Medicine</i> , 2016, 55, 643-646.	0.7	22
168	Relationship Between Number of Multiple Risk Factors and Coronary Artery Disease Risk With and Without Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 5084-5090.	3.6	22
169	Transgenic mice overexpressing SREBP-1a under the control of the PEPCK promoter exhibit insulin resistance, but not diabetes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2005, 1740, 427-433.	3.8	21
170	Oxygenomics in environmental stress. <i>Redox Report</i> , 2010, 15, 98-114.	4.5	21
171	Fasting and Post-Challenge Glucose as Quantitative Cardiovascular Risk Factors: A Meta-Analysis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 385-396.	2.0	21
172	Development of a Screening Score for Undiagnosed Diabetes and Its Application in Estimating Absolute Risk of Future Type 2 Diabetes in Japan: Toranomon Hospital Health Management Center Study 10 (TOPICS 10). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013, 98, 1051-1060.	3.6	21
173	Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). <i>Medicine (United States)</i> , 2017, 96, e6122.	1.0	21
174	Waist Circumference as a Cardiovascular and Metabolic Risk in Japanese Patients With Type 2 Diabetes. <i>Obesity</i> , 2009, 17, 585-592.	3.0	20
175	Genetic Variants of the Fatty Acid Desaturase Gene Cluster Are Associated with Plasma LDL Cholesterol Levels in Japanese Males. <i>Journal of Nutritional Science and Vitaminology</i> , 2013, 59, 325-335.	0.6	20
176	Trajectory of body mass index before the development of type 2 diabetes in Japanese men: Toranomon Hospital Health Management Center Study 15. <i>Journal of Diabetes Investigation</i> , 2015, 6, 289-294.	2.4	20
177	Risk factors associated with abnormal cognition in Japanese outpatients with diabetes, hypertension or dyslipidemia. <i>Diabetology International</i> , 2015, 6, 268-274.	1.4	20
178	Impact of body mass index and metabolic phenotypes on coronary artery disease according to glucose tolerance status. <i>Diabetes and Metabolism</i> , 2017, 43, 543-546.	2.9	20
179	Absence of Elovl6 attenuates steatohepatitis but promotes gallstone formation in a lithogenic diet-fed Ldlr ^{-/-} mouse model. <i>Scientific Reports</i> , 2015, 5, 17604.	3.3	20
180	The Regulation of Adiponectin Receptors Expression by Acute Exercise in Mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007, 115, 417-422.	1.2	19

#	ARTICLE	IF	CITATIONS
181	Fasting Insulin Levels and Metabolic Risk Factors in Type 2 Diabetic Patients at the First Visit in Japan: A 10-year, nationwide, observational study (JDDM 28). <i>Diabetes Care</i> , 2012, 35, 1853-1857.	8.6	19
182	Use of high-normal levels of haemoglobin A _{1c} and fasting plasma glucose for diabetes screening and for prediction: a meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2013, 29, 680-692.	4.0	19
183	Hypertension increases urinary excretion of immunoglobulin G, ceruloplasmin and transferrin in normoalbuminuric patients with type 2 diabetes mellitus. <i>Journal of Hypertension</i> , 2014, 32, 432-438.	0.5	19
184	Relationship between hemoglobin A1c and cardiovascular disease in mild-to-moderate hypercholesterolemic Japanese individuals: subanalysis of a large-scale randomized controlled trial. <i>Cardiovascular Diabetology</i> , 2011, 10, 58.	6.8	18
185	Direct effect of dasatinib on proliferation and cytotoxicity of natural killer cells in <i>in vitro</i> study. <i>Hematological Oncology</i> , 2013, 31, 156-163.	1.7	18
186	Role of Body Mass Index History in Predicting Risk of the Development of Hypertension in Japanese Individuals. <i>Hypertension</i> , 2014, 64, 247-252.	2.7	18
187	Attenuation of Weight Loss Through Improved Antilipolytic Effect in Adipose Tissue Via the SGLT2 Inhibitor Tofogliflozin. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 3647-3660.	3.6	18
188	Overt Proteinuria, Moderately Reduced eGFR and Their Combination Are Predictive of Severe Diabetic Retinopathy or Diabetic Macular Edema in Diabetes. , 2019, 60, 2685.		18
189	Pulse Pressure is a Stronger Predictor Than Systolic Blood Pressure for Severe Eye Diseases in Diabetes Mellitus. <i>Journal of the American Heart Association</i> , 2019, 8, e010627.	3.7	18
190	Vascular endothelial growth factor gene expression in a retinal pigmented cell is up-regulated by glucose deprivation through 3' UTR. <i>Life Sciences</i> , 2002, 71, 1607-1614.	4.3	17
191	Self-reported rate of eating is significantly associated with body mass index in Japanese patients with type 2 diabetes. Japan Diabetes Clinical Data Management Study Group (JDDM26). <i>Appetite</i> , 2012, 59, 252-255.	3.7	17
192	Management of Type IIb Dyslipidemia. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 105-114.	2.0	17
193	Weight Loss Maintenance for 2 Years after a 6-Month Randomised Controlled Trial Comparing Education-Only and Group-Based Support in Japanese Adults. <i>Obesity Facts</i> , 2014, 7, 376-387.	3.4	17
194	Unstable bodyweight and incident type 2 diabetes mellitus: A meta-analysis. <i>Journal of Diabetes Investigation</i> , 2017, 8, 501-509.	2.4	17
195	Successful 5-azacytidine treatment of myeloid sarcoma and leukemia cutis associated with myelodysplastic syndrome. <i>Medicine (United States)</i> , 2017, 96, e7975.	1.0	17
196	Vitamin B6 intake and incidence of diabetic retinopathy in Japanese patients with type 2 diabetes: analysis of data from the Japan Diabetes Complications Study (JDCCS). <i>European Journal of Nutrition</i> , 2020, 59, 1585-1594.	3.9	17
197	Efficacy and safety of empagliflozin as add-on to insulin in Japanese patients with type 2 diabetes: A randomized, double-blind, placebo-controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 417-426.	4.4	17
198	Associations of Systolic Blood Pressure and Diastolic Blood Pressure With the Incidence of Coronary Artery Disease or Cerebrovascular Disease According to Glucose Status. <i>Diabetes Care</i> , 2021, 44, 2124-2131.	8.6	17

#	ARTICLE	IF	CITATIONS
199	Influence of Adiponectin Gene Polymorphism SNP276 (G/T) on Adiponectin in Response to Exercise Training. <i>Endocrine Journal</i> , 2007, 54, 879-886.	1.6	16
200	Age, gender, insulin and blood glucose control status alter the risk of ischemic heart disease and stroke among elderly diabetic patients. <i>Cardiovascular Diabetology</i> , 2011, 10, 86.	6.8	16
201	Low BMI at age 20 years predicts gestational diabetes independent of BMI in early pregnancy in Japan: Tanaka Women's Clinic Study. <i>Diabetic Medicine</i> , 2013, 30, 70-73.	2.3	16
202	Utility of the Triglyceride Level for Predicting Incident Diabetes Mellitus According to the Fasting Status and Body Mass Index Category: The Ibaraki Prefectural Health Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014, 21, 1152-1169.	2.0	16
203	Role of fatty liver in the association between obesity and reduced hepatic insulin clearance. <i>Diabetes and Metabolism</i> , 2018, 44, 135-142.	2.9	16
204	Predictive ability of current machine learning algorithms for type 2 diabetes mellitus: A meta-analysis. <i>Journal of Diabetes Investigation</i> , 2022, 13, 900-908.	2.4	16
205	Impact of metabolic syndrome and metabolic dysfunction-associated fatty liver disease on cardiovascular risk by the presence or absence of type 2 diabetes and according to sex. <i>Cardiovascular Diabetology</i> , 2022, 21, .	6.8	16
206	Metabolic predictors of ischemic heart disease and cerebrovascular attack in elderly diabetic individuals: difference in risk by age. <i>Cardiovascular Diabetology</i> , 2013, 12, 10.	6.8	15
207	Quality and accuracy of Internet information concerning a healthy diet. <i>International Journal of Food Sciences and Nutrition</i> , 2013, 64, 1007-1013.	2.8	15
208	Ablation of Elovl6 protects pancreatic islets from high-fat diet-induced impairment of insulin secretion. <i>Biochemical and Biophysical Research Communications</i> , 2014, 450, 318-323.	2.1	15
209	Impact of individual components and their combinations within a family history of hypertension on the incidence of hypertension. <i>Medicine (United States)</i> , 2016, 95, e4564.	1.0	15
210	Meat intake and incidence of cardiovascular disease in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDCS). <i>European Journal of Nutrition</i> , 2019, 58, 281-290.	3.9	15
211	Nutrition education in Japanese medical schools: a follow-up survey. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2013, 22, 144-9.	0.4	15
212	Comparison of Education-Only versus Group-Based Intervention in Promoting Weight Loss: A Randomised Controlled Trial. <i>Obesity Facts</i> , 2011, 4, 222-228.	3.4	14
213	Pioglitazone treatment and cardiovascular event and death in subjects with type 2 diabetes without established cardiovascular disease (JDDM 36). <i>Diabetes Research and Clinical Practice</i> , 2015, 109, 485-492.	2.8	14
214	Expression of programmed death ligand 1 is associated with poor prognosis in myeloid sarcoma patients. <i>Hematological Oncology</i> , 2018, 36, 591-599.	1.7	14
215	Transgenic Mice Overexpressing SREBP-1a in Male ob/ob Mice Exhibit Lipodystrophy and Exacerbate Insulin Resistance. <i>Endocrinology</i> , 2018, 159, 2308-2323.	2.8	14
216	Association between Low Protein Intake and Mortality in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2020, 12, 1629.	4.1	14

#	ARTICLE	IF	CITATIONS
217	Vascular risk factors and diabetic neuropathy. <i>New England Journal of Medicine</i> , 2005, 352, 1925-7; author reply 1925-7.	27.0	14
218	Syndrome of Inappropriate Secretion of Antidiuretic Hormone(SIADH) and Gerhardt Syndrome Associated with Shy-Drager Syndrome.. <i>Internal Medicine</i> , 1994, 33, 773-778.	0.7	13
219	Effects of high glucose concentration and a thromboxane synthase inhibitor on the production of thromboxane A2 and prostaglandin I2 and E2 by cultured retinal endothelial cells. <i>Life Sciences</i> , 1995, 58, 239-243.	4.3	13
220	Primary Adrenal Lymphoma Presenting as Addisonian Crisis. Pitfalls in the Diagnosis of Bilateral Adrenal Swelling. <i>Hormone and Metabolic Research</i> , 1996, 28, 116-116.	1.5	13
221	The factors that affect exercise therapy for patients with type 2 diabetes in Japan: a nationwide survey. <i>Diabetology International</i> , 2015, 6, 19-25.	1.4	13
222	Comparison of clinical characteristics in patients with type 2 diabetes among whom different antihyperglycemic agents were prescribed as monotherapy or combination therapy by diabetes specialists. <i>Journal of Diabetes Investigation</i> , 2016, 7, 260-269.	2.4	13
223	Is the Proportion of Carbohydrate Intake Associated with the Incidence of Diabetes Complications? An Analysis of the Japan Diabetes Complications Study. <i>Nutrients</i> , 2017, 9, 113.	4.1	13
224	MAGI-1 expression is decreased in several types of human T-cell leukemia cell lines, including adult T-cell leukemia. <i>International Journal of Hematology</i> , 2018, 107, 337-344.	1.6	13
225	Familial predisposition to cardiovascular risk and disease contributes to cardiovascular risk and disease interacting with other cardiovascular risk factors in diabetes Implication for common soil (JDDM 14). <i>Atherosclerosis</i> , 2008, 201, 332-338.	0.8	12
226	Importance of Lipid Levels in Elderly Diabetic Individuals Baseline Characteristics and 1-Year Survey of Cardiovascular Events. <i>Circulation Journal</i> , 2008, 72, 218-225.	1.6	12
227	Contribution of first trimester fasting plasma insulin levels to the incidence of glucose intolerance in later pregnancy: Tanaka women's clinic study. <i>Diabetes Research and Clinical Practice</i> , 2011, 92, 293-298.	2.8	12
228	Comparison of clinicopathological characteristics between T-cell prolymphocytic leukemia and peripheral T-cell lymphoma, not otherwise specified. <i>European Journal of Haematology</i> , 2017, 98, 459-466.	2.2	12
229	Functional dissection of hematopoietic stem cell populations with a stemness-monitoring system based on NS-GFP transgene expression. <i>Scientific Reports</i> , 2017, 7, 11442.	3.3	12
230	Relationships among cardiorespiratory fitness, muscular fitness, and cardiometabolic risk factors in Japanese adolescents: Niigata screening for and preventing the development of non-communicable disease study-Agano (NICE EVIDENCE Study-Agano) 2. <i>Pediatric Diabetes</i> , 2018, 19, 593-602.	2.9	12
231	Risk of coronary artery disease according to glucose abnormality status and prior coronary artery disease in Japanese men. <i>Metabolism: Clinical and Experimental</i> , 2019, 101, 153991.	3.4	12
232	CREBH Improves Diet-Induced Obesity, Insulin Resistance, and Metabolic Disturbances by FGF21-Dependent and FGF21-Independent Mechanisms. <i>iScience</i> , 2020, 23, 100930.	4.1	12
233	A 52-week randomized controlled trial of ipragliflozin or sitagliptin in type 2 diabetes combined with metformin: The <i>SM</i> study. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 811-821.	4.4	12
234	Physical Fitness and Dyslipidemia Among Japanese: A Cohort Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2021, 31, 287-296.	2.4	12

#	ARTICLE	IF	CITATIONS
235	The Long-Term Effects of Self-Management Education for Patients With Type 2 Diabetes on Glycemic Control: Response to Norris et al.. <i>Diabetes Care</i> , 2002, 25, 2115-2116.	8.6	11
236	Risk Imparted by Various Parameters of Smoking in Japanese Men With Type 2 Diabetes on Their Development of Microalbuminuria: Analysis from the Tsukuba Kawai Diabetes Registry. <i>Diabetes Care</i> , 2007, 30, 1286-1288.	8.6	11
237	Are serum cholesterol levels associated with silent brain infarcts? The Seiryō Clinic Study. <i>Atherosclerosis</i> , 2010, 210, 674-677.	0.8	11
238	TFE3 inhibits myoblast differentiation in C2C12 cells via down-regulating gene expression of myogenin. <i>Biochemical and Biophysical Research Communications</i> , 2013, 430, 664-669.	2.1	11
239	Advanced glycation end products induce brain-derived neurotrophic factor release from human platelets through the Src-family kinase activation. <i>Cardiovascular Diabetology</i> , 2017, 16, 20.	6.8	11
240	Relationship between intake of fruit separately from vegetables and triglycerides - A meta-analysis. <i>Clinical Nutrition ESPEN</i> , 2018, 27, 53-58.	1.2	11
241	Enterohepatic Transcription Factor CREB3L3 Protects Atherosclerosis via SREBP Competitive Inhibition. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021, 11, 949-971.	4.5	11
242	Alcohol Use and Diabetes Mellitus. <i>Annals of Internal Medicine</i> , 2004, 141, 408.	3.9	11
243	Self-reported fast eating is a potent predictor of development of impaired glucose tolerance in Japanese men and women: Tsukuba Medical Center Study. <i>Diabetes Research and Clinical Practice</i> , 2011, 94, e72-e74.	2.8	10
244	Impact of introducing HbA1c into the diagnostic criteria on prevalence and cardiovascular risk profiles of individuals with newly diagnosed diabetes in Japan: The Toranomon Hospital Health Management Center Study 2 (TOPICS 2). <i>Diabetes Research and Clinical Practice</i> , 2012, 95, 283-290.	2.8	10
245	Prevalence and risk factors for diabetic maculopathy, and its relationship to diabetic retinopathy in elderly Japanese patients with type 2 diabetes mellitus. <i>Geriatrics and Gerontology International</i> , 2012, 12, 134-140.	1.5	10
246	Body Weight Change and Type 2 Diabetes. <i>Epidemiology</i> , 2013, 24, 778-779.	2.7	10
247	Early Diagnosis of Hepatic Intravascular Lymphoma: A Case Report and Literature Review. <i>Internal Medicine</i> , 2014, 53, 587-593.	0.7	10
248	The SIL index is a simple and objective prognostic indicator in diffuse large B-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2016, 57, 2763-2770.	1.3	10
249	Quantitative Relationship Between Cumulative Risk Alleles Based on Genome-Wide Association Studies and Type 2 Diabetes Mellitus: A Systematic Review and Meta-analysis. <i>Journal of Epidemiology</i> , 2018, 28, 3-18.	2.4	10
250	Effect of family-oriented diabetes programs on glycemic control: A meta-analysis. <i>Family Practice</i> , 2019, 36, 387-394.	1.9	10
251	A Prospective Cohort Study of Muscular and Performance Fitness and Risk of Hearing Loss: The Niigata Wellness Study. <i>American Journal of Medicine</i> , 2021, 134, 235-242.e4.	1.5	10
252	Maximum BMI and microvascular complications in a cohort of Japanese patients with type 2 diabetes: the Japan Diabetes Complications Study. <i>Journal of Diabetes and Its Complications</i> , 2016, 30, 790-797.	2.3	9

#	ARTICLE	IF	CITATIONS
253	Association between all-cause mortality and severity of depressive symptoms in patients with type 2 diabetes: Analysis from the Japan Diabetes Complications Study (JDCS). <i>Journal of Psychosomatic Research</i> , 2017, 99, 34-39.	2.6	9
254	Body flexibility and incident hypertension: The Niigata wellness study. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2021, 31, 702-709.	2.9	9
255	Distinct effects of chondroitin sulfate on hematopoietic cells and the stromal microenvironment in bone marrow hematopoiesis. <i>Experimental Hematology</i> , 2021, 96, 52-62.e5.	0.4	9
256	Rates and risk factors for amputation in people with diabetes in Japan: a historical cohort study using a nationwide claims database. <i>Journal of Foot and Ankle Research</i> , 2021, 14, 29.	1.9	9
257	Applications of physical performance measures to routine diabetes care for frailty prevention concept: fundamental data with grip strength, gait speed, timed chair stand speed, standing balance, and knee extension strength. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e001562.	2.8	9
258	Acute effect of beraprost sodium on lower limb circulation in patients with non-insulin-dependent diabetes mellitus-evaluation by color doppler ultrasonography and laser cutaneous blood flowmetry. <i>Prostaglandins</i> , 1996, 52, 375-384.	1.2	8
259	Disease model: hyperinsulinemia and insulin resistance Part B “polygenic and other animal models. <i>Trends in Molecular Medicine</i> , 2001, 7, 373-376.	6.7	8
260	Contribution of glimepiride to basal“prandial insulin therapy in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2011, 91, 148-153.	2.8	8
261	Present situation of exercise therapy for patients with diabetes mellitus in Japan: a nationwide survey. <i>Diabetology International</i> , 2012, 3, 86-91.	1.4	8
262	Impact on short-term glycaemic control of initiating diabetes care versus leaving diabetes untreated among individuals with newly screening-detected diabetes in Japan. <i>Journal of Epidemiology and Community Health</i> , 2014, 68, 1189-1195.	3.7	8
263	The association of level of reduction of Wilms“ tumor gene 1 mRNA transcript in bone marrow and outcome in acute myeloid leukemia patients. <i>Leukemia Research</i> , 2015, 39, 667-671.	0.8	8
264	Clinicopathological features of cryptococcal lymphadenitis and a review of literature. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2017, 57, 26-30.	0.8	8
265	Cardiovascular Disease in Japanese Patients with Type 2 Diabetes Mellitus. <i>Annals of Vascular Diseases</i> , 2018, 11, 2-14.	0.5	8
266	Predictors of the response of HbA1c and body weight after SGLT2 inhibition. <i>Diabetes and Metabolism</i> , 2018, 44, 172-174.	2.9	8
267	Higher pulse pressure predicts initiation of dialysis in Japanese patients with diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2019, 35, e3120.	4.0	8
268	Influence of an SGLT2 inhibitor, tofogliflozin, on the resting heart rate in relation to adipose tissue insulin resistance. <i>Diabetic Medicine</i> , 2020, 37, 1316-1325.	2.3	8
269	Association of increased hepatic insulin clearance and change in serum triglycerides or “hydroxybutyrate concentration via the sodium/glucose“cotransporter 2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2020, 22, 947-956.	4.4	8
270	Leisure“time physical activity and incidence of objectively assessed hearing loss: The Niigata Wellness Study. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2022, 32, 435-445.	2.9	8

#	ARTICLE	IF	CITATIONS
271	Dietary salt intake and diabetes complications in patients with diabetes: An overview. <i>Journal of General and Family Medicine</i> , 2017, 18, 16-20.	0.8	8
272	Quantification of BCR-ABL mRNA in Plasma/Serum of Patients with Chronic Myelogenous Leukemia. <i>International Journal of Medical Sciences</i> , 2012, 9, 901-908.	2.5	7
273	Prevalence of resistant hypertension and associated factors in Japanese subjects with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2015, 110, 18-25.	2.8	7
274	Development and evaluation of the Japanese version of the Audit of Diabetes-Dependent Quality of Life for patients with diabetes. <i>Diabetology International</i> , 2016, 7, 384-390.	1.4	7
275	Cladribine treatment for Erdheim-Chester disease involving the central nervous system and concomitant polycythemia vera: A case report. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2018, 58, 161-165.	0.8	7
276	Combination of diabetes mellitus and lack of habitual physical activity is a risk factor for functional disability in Japanese. <i>BMJ Open Diabetes Research and Care</i> , 2020, 8, e000901.	2.8	7
277	Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study. <i>JMIR Medical Informatics</i> , 2021, 9, e22148.	2.6	7
278	Sodium Intake and Incidence of Diabetes Complications in Elderly Patients with Type 2 Diabetes—Analysis of Data from the Japanese Elderly Diabetes Intervention Study (J-EDIT). <i>Nutrients</i> , 2021, 13, 689.	4.1	7
279	Diabetes Care in Emergency Settings. <i>Diabetes Care</i> , 1995, 18, 1310-1311.	8.6	6
280	Efficacy of Ibudilast on Lower Limb Circulation of Diabetic Patients with Minimally Impaired Baseline Flow. <i>Angiology</i> , 1995, 46, 699-703.	1.8	6
281	Association of living alone with the presence of undiagnosed diabetes in Japanese men: the role of modifiable risk factors for diabetes: Toranomon Hospital Health Management Center Study 13 (TOPICS) Tj ETQq1 20.784314 rgBT /Ove		
282	Reversible brain atrophy and cognitive impairment in an adolescent Japanese patient with primary adrenal Cushing's syndrome. <i>Neuropsychiatric Disease and Treatment</i> , 2014, 10, 1763.	2.2	6
283	Relationship between a Low Ankle Brachial Index and All-Cause Death and Cardiovascular Events in Subjects with and without Diabetes. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014, 21, .	2.0	6
284	The circulating level of leptin and blood pressure in patients with multiple system atrophy. <i>Journal of the Neurological Sciences</i> , 2014, 347, 349-351.	0.6	6
285	Type 1 Diabetes Mellitus and Isolated Adrenocorticotropin Deficiency Manifested by Parkinsonism: A Case Report and Literature Review. <i>Internal Medicine</i> , 2015, 54, 2629-2635.	0.7	6
286	Circulating Malondialdehyde-Modified LDL-Related Variables and Coronary Artery Stenosis in Asymptomatic Patients with Type 2 Diabetes. <i>Journal of Diabetes Research</i> , 2015, 2015, 1-8.	2.3	6
287	Cluster-randomized trial to improve the quality of diabetes management: The study for the efficacy assessment of the standard diabetes manual (SEAS-DM). <i>Journal of Diabetes Investigation</i> , 2016, 7, 539-543.	2.4	6
288	Carbohydrate intake during early pregnancy is inversely associated with abnormal glucose challenge test results in Japanese pregnant women. <i>Diabetes/Metabolism Research and Reviews</i> , 2017, 33, e2898.	4.0	6

#	ARTICLE	IF	CITATIONS
289	Severe hypoglycaemia is a major predictor of incident diabetic retinopathy in Japanese patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017, 43, 424-429.	2.9	6
290	Starvation-induced transcription factor CREBH negatively governs body growth by controlling GH signaling. <i>FASEB Journal</i> , 2021, 35, e21663.	0.5	6
291	Secular Trends in Dietary Intake over a 20-Year Period in People with Type 2 Diabetes in Japan: A Comparative Study of Two Nationwide Registries; Japan Diabetes Complications Study (JDCS) and Japan Diabetes Clinical Data Management Study (JDDM). <i>Nutrients</i> , 2021, 13, 3428.	4.1	6
292	Predicting long-term glycemic control of post-educational type II diabetic patients by evaluating serum 1,5-anhydroglucitol levels. <i>Diabetes Research and Clinical Practice</i> , 1996, 34, 83-88.	2.8	5
293	Requirement for hypertension and hyperlipidemia medication in U.S. and Japanese patients with diabetes. <i>American Journal of Medicine</i> , 2004, 117, 711-712.	1.5	5
294	Effect of weight reduction on concentration of plasma total homocysteine in obese Japanese men. <i>Obesity Research and Clinical Practice</i> , 2007, 1, 213-221.	1.8	5
295	Association of eating three meals irregularly with changes in BMI and weight among young Japanese men and women: A 2-year follow-up. <i>Physiology and Behavior</i> , 2016, 163, 81-87.	2.1	5
296	The Glasgow Prognostic Score as a pre-transplant risk assessment for allogeneic hematopoietic cell transplantation. <i>Clinical Transplantation</i> , 2017, 31, e13103.	1.6	5
297	Clinicopathological analysis of splenic red pulp low-grade B-cell lymphoma. <i>Pathology International</i> , 2020, 70, 280-286.	1.3	5
298	Association of estimated plasma volume and weight loss after long-term administration and subsequent discontinuation of the sodium-glucose cotransporter-2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2021, 23, 1660-1665.	4.4	5
299	Comparing Associations of Dietary Energy Density and Energy Intake, Macronutrients with Obesity in Patients with Type 2 Diabetes (JDDM 63). <i>Nutrients</i> , 2021, 13, 3167.	4.1	5
300	Altered microbiota by a high-fat diet accelerates lethal myeloid hematopoiesis associated with systemic SOCS3 deficiency. <i>IScience</i> , 2021, 24, 103117.	4.1	5
301	Influence of SGLT2 Inhibitor on Resting Heart Rate (RHR) and Factors Related to Its Changes. <i>Diabetes</i> , 2018, 67, .	0.6	5
302	Increased mutant frequency and altered mutation spectrum of the lacl transgene in Wilson disease rats with hepatitis. <i>Cancer Research</i> , 2000, 60, 5080-6.	0.9	5
303	Acute gliclazide administration enhances glucose and ketone body utilization in the perfused hind limb of normal and streptozotocin-diabetic rats. <i>Life Sciences</i> , 2002, 71, 647-654.	4.3	4
304	Enhancement of antigen presenting ability in the leukemic plasmacytoid dendritic cell line (PMDC05) by lentiviral vector-mediated transduction of CD80 gene. <i>Leukemia Research</i> , 2012, 36, 1541-1546.	0.8	4
305	Long-term risk factors for diabetic retinopathy and diabetic maculopathy in elderly Japanese patients with type 2 diabetes mellitus. <i>Geriatrics and Gerontology International</i> , 2012, 12, 141-144.	1.5	4
306	Importance of high-density lipoprotein cholesterol control during pravastatin treatment in hypercholesterolemic Japanese with type 2 diabetes mellitus: A post hoc analysis of MEGA study. <i>Diabetes Research and Clinical Practice</i> , 2013, 100, e31-e33.	2.8	4

#	ARTICLE	IF	CITATIONS
307	The <sc>D</sc>inakara equation for adjusting <sc>DLCO</sc> for hemoglobin in the <sc>HCT</sc>â€<sc>Cl</sc> is superior to the <sc>C</sc>otes equation for predicting highâ€risk patients in allogeneic hematopoietic stem cell transplantation. American Journal of Hematology, 2016, 91, E296.	4.1	4
308	Comparative effects of torasemide and furosemide on gap junction proteins and cardiac fibrosis in a rat model of dilated cardiomyopathy. BioFactors, 2017, 43, 187-194.	5.4	4
309	Anaplastic large cell lymphoma, with 1,25(OH)<sub>2</sub><sup>2</sup><sub>D</sub><sup>3</sup>-mediated hypercalcemia: A case report. Journal of Clinical and Experimental Hematopathology: JCEH, 2019, 59, 22-28.	0.8	4
310	Association of treatment-achieved HbA1c with incidence of coronary artery disease and severe eye disease in diabetes patients. Diabetes and Metabolism, 2020, 46, 331-334.	2.9	4
311	Family Support for Medical Nutritional Therapy and Dietary Intake among Japanese with Type 2 Diabetes (JDDM 56). Nutrients, 2020, 12, 2649.	4.1	4
312	Successful Treatment with Edoxaban for Disseminated Intravascular Coagulation in a Case of Aortic Dissection Complicated with Immune Thrombocytopenic Purpura. Internal Medicine, 2020, 59, 2035-2039.	0.7	4
313	Skipping breakfast, late-night eating and current smoking are associated with medication adherence in Japanese patients with diabetes. Primary Care Diabetes, 2020, 14, 753-759.	1.8	4
314	Prevention of postprandial hypotension-related syncope by caffeine in a patient with long-standing diabetes mellitus. Endocrine Journal, 2020, 67, 585-592.	1.6	4
315	Intensive oral care can reduce bloodstream infection with coagulase-negative staphylococci after neutrophil engraftment in allogeneic hematopoietic stem-cell transplantation. Supportive Care in Cancer, 2022, 30, 475-485.	2.2	4
316	Developing a health economic model for Asians with type 2 diabetes based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. BMJ Open Diabetes Research and Care, 2021, 9, e002177.	2.8	4
317	Impact of prior cerebrovascular disease and glucose status on incident cerebrovascular disease in Japanese. Cardiovascular Diabetology, 2021, 20, 174.	6.8	4
318	Higher Dietary Intake of Vitamin D Is Associated with Lower Incidence of Diabetic Nephropathy in Japanese Patients with Type 2 Diabetes. Diabetes, 2018, 67, 1561-P.	0.6	4
319	Morphological and functional adaptation of pancreatic islet blood vessels to insulin resistance is impaired in diabetic db/db mice. Biochimica Et Biophysica Acta - Molecular Basis of Disease, 2022, 1868, 166339.	3.8	4
320	YAP1/TAZ activity maintains vascular integrity and organismal survival. Biochemical and Biophysical Research Communications, 2022, 619, 117-123.	2.1	4
321	Restoration of myo-inositol uptake by eicosapentaenoic acid in human skin fibroblasts cultured in high-glucose medium. Life Sciences, 1995, 57, PL71-PL74.	4.3	3
322	Factors associated with inappropriate weight loss attempts by early adolescent girls in Japan. Eating and Weight Disorders, 2011, 16, e157-e163.	2.5	3
323	Manifestations of Fulminant CD8 T-cell Post-transplant Lymphoproliferative Disorder Following the Administration of Rituximab for Lymphadenopathy with a High Level of Epstein-Barr Virus (EBV) Replication after Allogeneic Hematopoietic Stem Cell Transplantation. Internal Medicine, 2014, 53, 2115-2119.	0.7	3
324	Simple selfâ€reported behavioral or psychological characteristics as risk factors for future type 2 diabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 14. Journal of Diabetes Investigation, 2015, 6, 236-241.	2.4	3

#	ARTICLE	IF	CITATIONS
325	A case of hypoglycemia attributable to atypical antipsychotic drugs. <i>Diabetology International</i> , 2015, 6, 341-346.	1.4	3
326	The Combination of Elevated Triglycerides and Abnormal Fasting Glucose Increases Risk of Cerebral Infarction in Patients With Mild to Moderate Hypercholesterolemia. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2015, 20, 169-173.	2.0	3
327	Refinement of the Glasgow Prognostic Score as a pre-transplant risk assessment for allogeneic hematopoietic cell transplantation. <i>International Journal of Hematology</i> , 2018, 108, 282-289.	1.6	3
328	Brain adaptations of insulin signaling kinases, GLUT 3, p-BADser155 and nitrotyrosine expression in various hypoglycemic models of mice. <i>Neurochemistry International</i> , 2020, 137, 104745.	3.8	3
329	Isolated Adrenocorticotrophic Hormone Deficiency Presenting with Severe Hyponatremia and Rhabdomyolysis: A Case Report and Literature Review. <i>American Journal of Case Reports</i> , 2019, 20, 1857-1863.	0.8	3
330	6-OR: Blood Pressure as Predictor of Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) According to Glucose Tolerance Status (GTS): Implications for Updated Guidelines. <i>Diabetes</i> , 2020, 69, 6-OR.	0.6	3
331	Evidence for the Diagnosis of Metabolic Syndrome in Japan. <i>The Japanese Journal of Nutrition and Dietetics</i> , 2011, 69, 205-213.	0.1	3
332	Background to Discuss Guidelines for Control of Plasma HDL-Cholesterol in Japan*. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012, 19, 207-212.	2.0	3
333	449-P: Impact of Body Mass Index (BMI) and Waist Circumference (WC) on Coronary Artery Disease (CAD) in Japanese with and without Diabetes Mellitus (DM). <i>Diabetes</i> , 2020, 69, 449-P.	0.6	3
334	1903-P: Low Level of and Decrease in Serum Amylase (AMY) Increases the Risk of Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2020, 69, 1903-P.	0.6	3
335	A Prospective Cohort Study of Muscular and Performance Fitness and Incident Glaucoma: The Niigata Wellness Study. <i>Journal of Physical Activity and Health</i> , 2020, 17, 1171-1178.	2.0	3
336	The effect of nilvadipine on bloodflow in the dorsal pedis artery in type 2 diabetic patients--a study using duplex Doppler ultrasonography.. <i>Postgraduate Medical Journal</i> , 1995, 71, 613-616.	1.8	2
337	Acute Effects of Thromboxane Dual Blocker (KDI-792) on Different Portions of Lower Limb Blood Flow--A Study Using Doppler Ultrasonography and Laser Doppler Flowmetry in Type 2 Diabetic Patients. <i>Prostaglandins</i> , 1997, 53, 395-409.	1.2	2
338	Plasma chloride concentration as a new diagnostic indicator of insulin insufficiency. <i>Diabetes Research and Clinical Practice</i> , 2005, 67, 137-143.	2.8	2
339	Importance of high-density lipoprotein cholesterol levels in elderly diabetic individuals with type IIb dyslipidemia: A 2-year survey of cardiovascular events. <i>Geriatrics and Gerontology International</i> , 2014, 14, 806-810.	1.5	2
340	Second trimester postload glucose level as an important predictor of low birth weight infants: Tanaka Women's Clinic Study. <i>Diabetes Research and Clinical Practice</i> , 2014, 105, e16-e19.	2.8	2
341	Fasting mediated increase in p-BADser155 and p-AKTser473 in the prefrontal cortex of mice. <i>Neuroscience Letters</i> , 2014, 579, 134-139.	2.1	2
342	Fatal tracheal aspergillosis during rituximab combined chemotherapy for diffuse large B-cell lymphoma that developed after lung transplantation. <i>Transplant Infectious Disease</i> , 2015, 17, 872-875.	1.7	2

#	ARTICLE	IF	CITATIONS
343	Correlation between SNP genotypes and periodontitis in Japanese type II diabetic patients: a preliminary study. <i>Odontology / the Society of the Nippon Dental University</i> , 2015, 103, 233-240.	1.9	2
344	Meta-analytic research on the relationship between cumulative risk alleles and risk of type 2 diabetes mellitus. <i>Diabetes/Metabolism Research and Reviews</i> , 2016, 32, 178-186.	4.0	2
345	Utility of nonblood-based risk assessment for predicting type 2 diabetes mellitus: A meta-analysis. <i>Preventive Medicine</i> , 2016, 91, 180-187.	3.4	2
346	Comparative evaluation of torasemide and spironolactone on adverse cardiac remodeling in a rat model of dilated cardiomyopathy. <i>Cardiovascular Therapeutics</i> , 2017, 35, e12283.	2.5	2
347	Marker chromosome is a strong poor prognosis factor after allogeneic HSCT for adverse-risk AML patients. <i>European Journal of Haematology</i> , 2020, 105, 616-625.	2.2	2
348	Le Carbone prevents liver damage in non-alcoholic steatohepatitis-hepatocellular carcinoma mouse model via AMPK β -SIRT1 signaling pathway activation. <i>Heliyon</i> , 2021, 7, e05888.	3.2	2
349	DA-EPOCH-R therapy for high-grade B-cell lymphoma with <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> rearrangements in a patient with renal dysfunction. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2021, 61, 42-47.	0.8	2
350	Developing a Health Economic Model for Asians with Type 2 Diabetes Based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>Diabetes</i> , 2018, 67, .	0.6	2
351	Impact of Medication Adherence and Glycemic Control on the Risk of Micro- and Macrovascular Diseases in Patients with Diabetes. <i>American Journal of Medicine</i> , 2022, 135, 461-470.e1.	1.5	2
352	Dietary intake and physical activity in Japanese patients with type 2 diabetes: the Japan Diabetes Complication and its Prevention prospective study (JDCP study 8). <i>Diabetology International</i> , 2022, 13, 344-357.	1.4	2
353	Transient Nephrogenic Diabetes insipidus Accompanied by Possible Psychogenic Polydipsia. <i>Hormone Research</i> , 1995, 44, 193-196.	1.8	1
354	Body Image: International Comparisons. <i>Nutrition Today</i> , 2010, 45, 113-117.	1.0	1
355	The Level of Orally Ingested Vitamin C Affected the Expression of Vitamin C Transporters and Vitamin C Accumulation in the Livers of ODS Rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011, 75, 2394-2397.	1.3	1
356	Commentary on the United Kingdom Prospective Diabetes Study outcomes model 2: Need for long-term follow up and quality of life data in Asian patients. <i>Journal of Diabetes Investigation</i> , 2014, 5, 281-283.	2.4	1
357	Efficacy of HMG-CoA reductase inhibitors in the prevention of cerebrovascular attack in 1016 patients older than 75 years among 4014 type 2 diabetic individuals. <i>International Journal of Cardiology</i> , 2014, 177, 860-866.	1.7	1
358	Food groups and weight gain in Japanese men. <i>Clinical Obesity</i> , 2014, 4, 157-164.	2.0	1
359	Fibromyalgia in a Patient with Cushing's Disease Accompanied by Central Hypothyroidism. <i>Internal Medicine</i> , 2016, 55, 3185-3190.	0.7	1
360	Rapid Normalization of High Glutamic Acid Decarboxylase Autoantibody Titers and Preserved Endogenous Insulin Secretion in a Patient with Diabetes Mellitus: A Case Report and Literature Review. <i>Internal Medicine</i> , 2016, 55, 485-489.	0.7	1

#	ARTICLE	IF	CITATIONS
361	Quantitative assessment of genetic testing for type 2 diabetes mellitus based on findings of genome-wide association studies. <i>Annals of Epidemiology</i> , 2016, 26, 816-818.e6.	1.9	1
362	Heterogeneity of intrahepatic iron deposition in transfusion-dependent iron overload patients with hematological malignancies. <i>Leukemia Research</i> , 2018, 70, 41-44.	0.8	1
363	Combined Effects of Energy Intake and Physical Activity on Obesity in Japanese Patients with Type 2 Diabetes (JDDM 50): A Cross-Sectional Study. <i>Diabetes Therapy</i> , 2019, 10, 1133-1138.	2.5	1
364	Safety and effective salvage regimen comprising a novel combination of brentuximab vedotin, L-asparaginase, and dexamethasone for refractory anaplastic large cell lymphoma, anaplastic lymphoma kinase negative. <i>Hematological Oncology</i> , 2019, 37, 212-214.	1.7	1
365	Gemcitabine, Dexamethasone, and Cisplatin Regimen as an Effective Salvage Therapy for High-grade B-cell Lymphoma with <i>MYC</i> and <i>BCL2</i> and/or <i>BCL6</i> Rearrangements. <i>Internal Medicine</i> , 2019, 58, 575-580.	0.7	1
366	Network Meta-Analysis of Drug Therapies for Lowering Uric Acid and Mortality Risk in Patients with Heart Failure. <i>Cardiovascular Drugs and Therapy</i> , 2020, 35, 1217-1225.	2.6	1
367	WT1-specific CD8 ⁺ cytotoxic T cells with the capacity for antigen-specific expansion accumulate in the bone marrow in MDS. <i>International Journal of Hematology</i> , 2021, 113, 723-734.	1.6	1
368	Dipeptidyl peptidase-4 inhibitor, anagliptin, alters hepatic insulin clearance in relation to the glycemic status in Japanese individuals with type 2 diabetes. <i>Journal of Diabetes Investigation</i> , 2021, 12, 1805-1815.	2.4	1
369	767-P: Family Support for Medical Nutritional Therapy and Dietary Intake among Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019, 68, .	0.6	1
370	Test Strip-Positive Proteinuria and Its Combination with Low eGFR Are Predictive of Treatment-Required Eye Diseases in Japanese Patients with Diabetes Mellitus. <i>Diabetes</i> , 2018, 67, .	0.6	1
371	Incidence and Risk Factors for Amputation in Patients with Diabetes in Japan—Historical Cohort Study Using a Nationwide Claims Database. <i>Diabetes</i> , 2018, 67, 637-P.	0.6	1
372	Distinct Effects of Chondroitin Sulfate on Hematopoietic Cells and the Stromal Microenvironment in Bone Marrow Hematopoiesis. <i>Blood</i> , 2018, 132, 3852-3852.	1.4	1
373	598-P: Lower Hematocrit Is Predictive of Treatment-Required Eye Diseases in Japanese Patients with Diabetes Mellitus. <i>Diabetes</i> , 2019, 68, .	0.6	1
374	789-P: Association of Zinc Intake with Obesity in Japanese Patients with Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019, 68, .	0.6	1
375	854-P: Personality, Self-Management Behaviors, and Glycemic Control among Japanese Patients with Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019, 68, 854-P.	0.6	1
376	1126-P: Randomized Controlled Trial of Ipragliflozin or Sitagliptin Combined with Metformin in Type 2 Diabetes: NISM Study. <i>Diabetes</i> , 2020, 69, 1126-P.	0.6	1
377	1267-P: Physical Fitness (PF), Weight Status, and Metabolic Risk in Japanese Adolescents. <i>Diabetes</i> , 2020, 69, .	0.6	1
378	Severity of hypertension as a predictor of initiation of dialysis among study participants with and without diabetes mellitus. <i>Journal of Investigative Medicine</i> , 2021, 69, 724-729.	1.6	1

#	ARTICLE	IF	CITATIONS
379	Higher Iron Intake Is Independently Associated with Obesity in Younger Japanese Type-2 Diabetes Mellitus Patients. <i>Nutrients</i> , 2022, 14, 211.	4.1	1
380	Differences in occupational stress by smoking intensity and gender in cross-sectional study of 59 355 Japanese employees using the Brief Job Stress Questionnaire (BJSQ): the Niigata Wellness Study. <i>BMJ Open</i> , 2022, 12, e055577.	1.9	1
381	Weight and cardiometabolic risk among adolescents in Agano city, Japan: NICE EVIDENCE Study-Agano 1. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020, 29, 856-866.	0.4	1
382	No Relationship Between Body Mass Index During Adolescence and All-Cause Mortality in Japanese Women—A 56.5-Year Observational Study. <i>Annals of Epidemiology</i> , 2009, 19, 590-591.	1.9	0
383	A Hierarchical Regression Model for Dietary Data Adjusting for Covariates Measurement Error by Regression Calibration: An Application to a Large Prospective Study for Diabetic Complications. <i>Japanese Journal of Biometrics</i> , 2010, 31, 49-62.	0.0	0
384	Accumulation of cardiovascular risks in Japanese women with abnormal glucose and mild to moderate hypercholesterolemia. <i>International Journal of Cardiology</i> , 2011, 152, 254-256.	1.7	0
385	Erratum to “Accumulation of cardiovascular risks in Japanese women with abnormal glucose and mild to moderate hypercholesterolemia” [International Journal of Cardiology 152 (2) (2011) 254–256]. <i>International Journal of Cardiology</i> , 2011, 153, 238.	1.7	0
386	Dose of 3- α -methylcholanthrene enhances vitamin C accumulation and mRNA expression of its transporter in the liver of ODS rats and in HepG2 cells. <i>Journal of Biochemical and Molecular Toxicology</i> , 2011, 25, 369-376.	3.0	0
387	Two Authors Reply. <i>American Journal of Epidemiology</i> , 2013, 177, 863-863.	3.4	0
388	Correction to: Comparison of Education-Only versus Group-Based Intervention in Promoting Weight Loss: A Randomised Controlled Trial. <i>Obesity Facts</i> , 2013, 6, 89-90.	3.4	0
389	Response to Comment on Heianza et al. Effect of Postmenopausal Status and Age at Menopause on Type 2 Diabetes and Prediabetes in Japanese Individuals: Toranomon Hospital Health Management Center Study 17 (TOPICS 17). <i>Diabetes Care</i> 2013;36:4007–4014. <i>Diabetes Care</i> , 2014, 37, e165-e166.	8.6	0
390	Efficacy of Habitual Exercise for Improving Lipid Profiles Depends on the PPRA ¹ Genotype in Japanese Males. <i>Journal of Nutritional Science and Vitaminology</i> , 2014, 60, 66-70.	0.6	0
391	Combination effect of hypertension and diabetes mellitus on urinary protein excretion. <i>Journal of Hypertension</i> , 2014, 32, 2278.	0.5	0
392	Role of the polycomb gene BCOR in hematopoiesis. <i>Experimental Hematology</i> , 2015, 43, S83.	0.4	0
393	Reply to: Relationship between leptin and blood pressure in patients with multiple system atrophy. <i>Journal of the Neurological Sciences</i> , 2015, 348, 285.	0.6	0
394	Potential impact of joint association of alanine aminotransferase and gamma-glutamyltransferase on insulin resistance in Japan: The Toranomon Hospital Health Management Center Study 19 (TOPICS 19). <i>Hepatology Research</i> , 2015, 45, 247-258.	3.4	0
395	Assessment of kidney dysfunction with cystatin C- and creatinine-based estimated glomerular filtration rate and predicting type 2 diabetes: Toranomon Hospital Health Management Center Study 21. <i>Diabetes Research and Clinical Practice</i> , 2016, 113, 60-68.	2.8	0
396	OBSOLETE: Diabetes Mellitus. , 2018, , .		0

#	ARTICLE	IF	CITATIONS
397	Glutamic Acid Decarboxylase Autoantibody-negative Slowly Progressive Type 1 Diabetes Mellitus: A Case Report and Literature Review. <i>Internal Medicine</i> , 2018, 57, 3581-3587.	0.7	0
398	The Glasgow prognostic score divides high-risk hematopoietic cell transplantation-specific comorbidity index patients into stratified subgroups in allogeneic hematopoietic cell transplantation. <i>Annals of Hematology</i> , 2020, 99, 671-673.	1.8	0
399	Meta-analytic research of the dose-response relationship between salt intake and risk of heart failure. <i>Hypertension Research</i> , 2021, 44, 885-887.	2.7	0
400	Carrot Consumption Frequency Associated with Reduced BMI and Obesity through the SNP Intermediary rs4445711. <i>Nutrients</i> , 2021, 13, 3478.	4.1	0
401	EFFECT OF EXERCISE TRAINING ON SERUM HIGH-SENSITIVITY C-REACTIVE PROTEIN CONCENTRATION IN HEALTHY MIDDLE-AGED AND ELDERLY SUBJECTS. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2007, 56, 179-190.	0.0	0
402	Early Responses At 3 Months and 12 Months After Starting Imatinib As Predictive Factors For The Achievement Of Deep MR In Japanese CML Patients. <i>Blood</i> , 2013, 122, 2744-2744.	1.4	0
403	Log Reduction Levels of WT1 mRNA Expression in BM after Chemotherapies Are Predictive Markers of Good Prognosis in AML Patients Achieved CR after Induction Therapy. <i>Blood</i> , 2014, 124, 2334-2334.	1.4	0
404	Clinical Significance of MYC, BCL2 and BCL6 Rearrangement and Protein Expression in GCB and Non-GCB Type Diffuse Large B-Cell Lymphoma. <i>Blood</i> , 2015, 126, 1445-1445.	1.4	0
405	The Sil Index Is a Useful Prognostic Indicator for Diffuse Large B-Cell Lymphoma. <i>Blood</i> , 2015, 126, 1512-1512.	1.4	0
406	Sedentary Behavior, Physical Activity and Kidney Stones. <i>Medicine and Science in Sports and Exercise</i> , 2016, 48, 226.	0.4	0
407	Evaluation of Liver Iron Deposition in Transfusion-Dependent Patients By Dual-Energy CT. <i>Blood</i> , 2016, 128, 3619-3619.	1.4	0
408	The Predictive Factors of Favorable Prognosis after Allo-HSCT for Refractory Acute Leukemia. <i>Blood</i> , 2016, 128, 2297-2297.	1.4	0
409	Genetically Reduced Chondroitin Sulfate Prevents the Progression of Diabetic Neuropathy. <i>Diabetes</i> , 2018, 67, .	0.6	0
410	Impact of Prior Coronary Artery Disease (CAD) and Glucose Tolerance Status (GTS) on Incident CAD in Japanese Men. <i>Diabetes</i> , 2018, 67, 1488-P.	0.6	0
411	Impact of Vitamin B6 Intake on the Risk of Diabetic Retinopathy—Analysis from Multicenter Prospective Study of Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2018, 67, 597-P.	0.6	0
412	Impact of Carbohydrate Intake on Obesity in Japanese Patients with Type 2 Diabetes—An Analysis of the JDCP Registry. <i>Diabetes</i> , 2018, 67, .	0.6	0
413	Dipstick Proteinuria as a Predictor of End-Stage Renal Disease in Japanese Adults With and Without Diabetes Mellitus (DM). <i>Diabetes</i> , 2018, 67, 2384-PUB.	0.6	0
414	Education for Family Members Is Effective for Improved Glycemic Control of Patients with Type 2 Rather than Type 1 Diabetes Mellitus—A Meta-analysis. <i>Diabetes</i> , 2018, 67, .	0.6	0

#	ARTICLE	IF	CITATIONS
415	Effects of Treatment-Achieved HbA1c on Incidence of Micro-/Macrovascular Complications in Patients with Diabetes Mellitus. <i>Diabetes</i> , 2018, 67, .	0.6	0
416	Effect of Number of Achieved Targets for Risk Factors on Coronary Artery Disease (CAD) in Those With and Without Diabetes Mellitus (DM). <i>Diabetes</i> , 2018, 67, 442-P.	0.6	0
417	Association between Serum Amylase Level and Incidence of Type 2 Diabetes. <i>Diabetes</i> , 2018, 67, .	0.6	0
418	Reduced Postprandial Hepatic Insulin Clearance via the DPP-4 Inhibitor Anagliptin Contributed to Improvement in Hyperglycemia in Patients with Type 2 Diabetes Mellitus. <i>Diabetes</i> , 2018, 67, .	0.6	0
419	Marker Chromosomes Are a New Cytogenetic Adverse Risk Factor in AML after Allo-HCT. <i>Blood</i> , 2018, 132, 5260-5260.	1.4	0
420	Clinical Features and Risk Factors of Post-Engraftment Bloodstream Infection in Allogeneic HCT. <i>Blood</i> , 2018, 132, 5712-5712.	1.4	0
421	634-P: Poor Adherence to Medication and HbA1c Level Predict Risk of Amputation in Patients with Diabetes Mellitus—Historical Cohort Study Using a Nationwide Claims Database. <i>Diabetes</i> , 2019, 68, .	0.6	0
422	1335-P: Weight Status and Cardiometabolic Risk Factors among Adolescents in Japan. <i>Diabetes</i> , 2019, 68, 1335-P.	0.6	0
423	1579-P: Higher Calcium Intake Is Associated with Lower Incidence of Diabetic Nephropathy in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019, 68, .	0.6	0
424	1219-P: Enhanced Response in Serum Ketone Level in Men Compared with Women by Administration of SGLT2 Inhibitor. <i>Diabetes</i> , 2019, 68, 1219-P.	0.6	0
425	1205-P: Baseline Lactate Level Is a Useful Predictor for Weight Loss after Long-Term SGLT2 Inhibitor Treatment. <i>Diabetes</i> , 2019, 68, .	0.6	0
426	2450-PUB: Association between Intake of Magnesium and Obesity in Japanese Patients with Type 2 Diabetes Mellitus. <i>Diabetes</i> , 2019, 68, .	0.6	0
427	2078-P: Association of Dietary Intake of Phosphorus with Obesity in Japanese Patients with Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019, 68, 2078-P.	0.6	0
428	698-P: Factors Significantly Associated with Adherence to Diabetes Medications: Findings from a Large Japanese Claims Database. <i>Diabetes</i> , 2019, 68, .	0.6	0
429	450-P: Impact of Prior Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) and Diabetes Mellitus (DM) on Incident CAD/CVD in Japanese. <i>Diabetes</i> , 2019, 68, .	0.6	0
430	1583-P: Intake of Fish and Related Nutrients in Association with Obesity in Japanese Patients with Type 2 Diabetes (T2D). <i>Diabetes</i> , 2019, 68, .	0.6	0
431	775-P: Significant Association of Food Group Intake with Obesity among Patients with Type 2 Diabetes Mellitus in Japan. <i>Diabetes</i> , 2019, 68, .	0.6	0
432	448-P: Predictors and Their Impact on Coronary Artery Disease (CAD) According to Glucose Tolerance Status (GTS) and Prior CAD: Historical Cohort Study in Japan. <i>Diabetes</i> , 2019, 68, .	0.6	0

#	ARTICLE	IF	CITATIONS
433	1478-P: Impact of Prior Cerebrovascular Disease (CVD) and Glucose Tolerance Status on Incident CVD in Japanese. <i>Diabetes</i> , 2019, 68, .	0.6	0
434	1215-P: Adipose Tissue Insulin Resistance Predicts Ketosis via an SGLT2 Inhibitor. <i>Diabetes</i> , 2019, 68, .	0.6	0
435	1910-P: SOCS3 Deficiency on a High-Fat Diet Accelerates Systemic Inflammation and Results in Lethal Myeloid Hematopoiesis without Obesity and Adiposity. <i>Diabetes</i> , 2019, 68, 1910-P.	0.6	0
436	446-P: Associations of Blood Pressure (BP) with Incidence of Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) According to Glucose Tolerance Status (GTS). <i>Diabetes</i> , 2019, 68, .	0.6	0
437	A Prospective Cohort Study of Physical Fitness and Incident Glaucoma: The Niigata Wellness Study. <i>Medicine and Science in Sports and Exercise</i> , 2019, 51, 222-222.	0.4	0
438	764-P: Low Dietary Energy Density (DED) Diet Is Associated with Favorable Dietary Pattern in Japanese Patients with Type 2 Diabetes (T2DM). <i>Diabetes</i> , 2019, 68, 764-P.	0.6	0
439	1514-P: Serum Amylase (AMY) Level and Its Chronological Change as a Predictor of Incident Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019, 68, .	0.6	0
440	Depletion of Pre-Transplant Skeletal Muscle Is a Significant Poor Prognostic Factor in Allogeneic Hematopoietic Cell Transplantation. <i>Blood</i> , 2019, 134, 3322-3322.	1.4	0
441	518-P: Association between Plasma Metabolites in the Urea Cycle and Diabetic Kidney Disease (DKD) in a Japanese Population. <i>Diabetes</i> , 2020, 69, .	0.6	0
442	1541-P: Dietary Patterns Significantly Associated with Obesity in Japanese with Type 2 Diabetes: JDDM. <i>Diabetes</i> , 2020, 69, 1541-P.	0.6	0
443	1133-P: Association of Higher Baseline BNP Levels with a Greater Reduction in Plasma Volume and Increase in Beta-Hydroxybutyrate via the SGLT2 Inhibitor Tofogliflozin in Type 2 Diabetes. <i>Diabetes</i> , 2020, 69, 1133-P.	0.6	0
444	841-P: Predictive Ability of Incident Type 2 Diabetes Mellitus (T2DM) Using Machine Learning Algorithms: A Meta-analysis. <i>Diabetes</i> , 2020, 69, 841-P.	0.6	0
445	1125-P: Association of Plasma Volume with Body Weight and BNP after Long-Term Administration and Subsequent Withdrawal of the SGLT2 Inhibitor Tofogliflozin. <i>Diabetes</i> , 2020, 69, .	0.6	0
446	1513-P: Severity of Hypertension (HT) as a Predictor of Initiation of Dialysis among Study Participants with and without Diabetes Mellitus (DM). <i>Diabetes</i> , 2020, 69, .	0.6	0
447	1988-P: Association between Screen Time Including Smartphone Use and Obesity and Its Associated Lifestyles in School Children in Japan. <i>Diabetes</i> , 2020, 69, .	0.6	0
448	1553-P: Combination of Diabetes Mellitus and Lack of Habitual Physical Activity Is a Risk Factor for Functional Disability in Japanese. <i>Diabetes</i> , 2020, 69, .	0.6	0
449	389-P: Ability for Detecting or Predicting Hypoglycemia with the Aid of Machine Learning Techniques: A Meta-analysis. <i>Diabetes</i> , 2020, 69, .	0.6	0
450	1485-P: Difference between Comprehensive Physical Fitness Age and Calendar Age Is a Potent Predictor of Incident Metabolic Syndrome (MetS). <i>Diabetes</i> , 2020, 69, 1485-P.	0.6	0

#	ARTICLE	IF	CITATIONS
451	663-P: Medication Adherence (MA) Is an Independent Predictor for Glycemic Control Even after Adjustment for Lifestyle Confounders in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2020, 69, .	0.6	0
452	A Prospective Cohort Study Of Physical Fitness And Incident Hearing Loss: The Niigata Wellness Study. <i>Medicine and Science in Sports and Exercise</i> , 2020, 52, 421-421.	0.4	0
453	<i>Genetic Manipulation Resulting in Decreased Donor Chondroitin sulfate Synthesis Mitigates Gvhd Following Allogeneic Hematopoietic Cell Transplantation in a Murine Model.</i> <i>Blood</i> , 2020, 136, 25-26.	1.4	0