

Twan van den Beucken

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3465055/publications.pdf>

Version: 2024-02-01

42
papers

3,152
citations

257450

24
h-index

254184

43
g-index

43
all docs

43
docs citations

43
times ranked

7110
citing authors

#	ARTICLE	IF	CITATIONS
1	The unfolded protein response protects human tumor cells during hypoxia through regulation of the autophagy genes MAP1LC3B and ATG5. <i>Journal of Clinical Investigation</i> , 2010, 120, 127-141.	8.2	675
2	Gene expression during acute and prolonged hypoxia is regulated by distinct mechanisms of translational control. <i>EMBO Journal</i> , 2006, 25, 1114-1125.	7.8	328
3	Hypoxia promotes stem cell phenotypes and poor prognosis through epigenetic regulation of DICER. <i>Nature Communications</i> , 2014, 5, 5203.	12.8	195
4	PERK/eIF2 \pm signaling protects therapy resistant hypoxic cells through induction of glutathione synthesis and protection against ROS. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013, 110, 4622-4627.	7.1	193
5	Taking advantage of tumor cell adaptations to hypoxia for developing new tumor markers and treatment strategies. <i>Journal of Enzyme Inhibition and Medicinal Chemistry</i> , 2009, 24, 1-39.	5.2	167
6	Hypoxia-mediated downregulation of miRNA biogenesis promotes tumour progression. <i>Nature Communications</i> , 2014, 5, 5202.	12.8	151
7	Control of the hypoxic response through regulation of mRNA translation. <i>Seminars in Cell and Developmental Biology</i> , 2005, 16, 487-501.	5.0	141
8	Translational control of gene expression during hypoxia. <i>Cancer Biology and Therapy</i> , 2006, 5, 749-755.	3.4	126
9	Two phases of disulfide bond formation have differing requirements for oxygen. <i>Journal of Cell Biology</i> , 2013, 203, 615-627.	5.2	113
10	The hypoxic proteome is influenced by gene-specific changes in mRNA translation. <i>Radiotherapy and Oncology</i> , 2005, 76, 177-186.	0.6	105
11	Affinity maturation of Fab antibody fragments by fluorescent-activated cell sorting of yeast-displayed libraries. <i>FEBS Letters</i> , 2003, 546, 288-294.	2.8	92
12	Targeting hypoxia tolerance in cancer. <i>Drug Resistance Updates</i> , 2004, 7, 25-40.	14.4	81
13	Construction and diversification of yeast cell surface displayed libraries by yeast mating: application to the affinity maturation of Fab antibody fragments. <i>Gene</i> , 2004, 342, 211-218.	2.2	75
14	Cell Surface Profiling Using High-Throughput Flow Cytometry: A Platform for Biomarker Discovery and Analysis of Cellular Heterogeneity. <i>PLoS ONE</i> , 2014, 9, e105602.	2.5	65
15	Hypoxia increases genome-wide bivalent epigenetic marking by specific gain of H3K27me3. <i>Epigenetics and Chromatin</i> , 2016, 9, 46.	3.9	63
16	Hypoxia-induced Expression of Carbonic Anhydrase 9 Is Dependent on the Unfolded Protein Response. <i>Journal of Biological Chemistry</i> , 2009, 284, 24204-24212.	3.4	57
17	Phosphorylation of eIF2 \pm is required for mRNA translation inhibition and survival during moderate hypoxia. <i>Radiotherapy and Oncology</i> , 2007, 83, 353-361.	0.6	54
18	Building novel binding ligands to B7.1 and B7.2 based on human antibody single variable light chain domains 1 Edited by I. A. Wilson. <i>Journal of Molecular Biology</i> , 2001, 310, 591-601.	4.2	47

#	ARTICLE	IF	CITATIONS
19	The mTOR target 4Eâ€BP1 contributes to differential protein expression during normoxia and hypoxia through changes in mRNA translation efficiency. <i>Proteomics</i> , 2008, 8, 1019-1028.	2.2	45
20	Translational control is a major contributor to hypoxia induced gene expression. <i>Radiotherapy and Oncology</i> , 2011, 99, 379-384.	0.6	37
21	Hypoxia disrupts the Fanconi anemia pathway and sensitizes cells to chemotherapy through regulation of UBE2T. <i>Radiotherapy and Oncology</i> , 2011, 101, 190-197.	0.6	36
22	Cigarette Smoke Extract Induces a Phenotypic Shift in Epithelial Cells; Involvement of HIF1± in Mesenchymal Transition. <i>PLoS ONE</i> , 2014, 9, e107757.	2.5	34
23	Identification of essential transcription factors for adequate DNA damage response after benzo(a)pyrene and aflatoxin B1 exposure by combining transcriptomics with functional genomics. <i>Toxicology</i> , 2017, 390, 74-82.	4.2	31
24	Regulation of TRIB3 mRNA and Protein in Breast Cancer. <i>PLoS ONE</i> , 2012, 7, e49439.	2.5	28
25	Regulation of Cited2 expression provides a functional link between translational and transcriptional responses during hypoxia. <i>Radiotherapy and Oncology</i> , 2007, 83, 346-352.	0.6	24
26	Deficient carbonic anhydrase 9 expression in UPR-impaired cells is associated with reduced survival in an acidic microenvironment. <i>Radiotherapy and Oncology</i> , 2009, 92, 437-442.	0.6	23
27	Quantitative analysis of ChIP-seq data uncovers dynamic and sustained H3K4me3 and H3K27me3 modulation in cancer cells under hypoxia. <i>Epigenetics and Chromatin</i> , 2016, 9, 48.	3.9	23
28	Proteomic analysis of gene expression following hypoxia and reoxygenation reveals proteins involved in the recovery from endoplasmic reticulum and oxidative stress. <i>Radiotherapy and Oncology</i> , 2007, 83, 340-345.	0.6	21
29	Deregulation of cap-dependent mRNA translation increases tumour radiosensitivity through reduction of the hypoxic fraction. <i>Radiotherapy and Oncology</i> , 2011, 99, 385-391.	0.6	21
30	Canonical autophagy does not contribute to cellular radioresistance. <i>Radiotherapy and Oncology</i> , 2015, 114, 406-412.	0.6	21
31	Radioprotective effects of ATP in human blood ex vivo. <i>Biochemical and Biophysical Research Communications</i> , 2008, 367, 383-387.	2.1	19
32	Valproic acid promotes mitochondrial dysfunction in primary human hepatocytes in vitro; impact of C/EBP±-controlled gene expression. <i>Archives of Toxicology</i> , 2020, 94, 3463-3473.	4.2	11
33	Dynamic Interplay between the Transcriptome and Methylome in Response to Oxidative and Alkylating Stress. <i>Chemical Research in Toxicology</i> , 2016, 29, 1428-1438.	3.3	8
34	Human-induced pluripotent stem cells as a model for studying sporadic Alzheimerâ€™s disease. <i>Neurobiology of Learning and Memory</i> , 2020, 175, 107318.	1.9	8
35	In vivo optical imaging of MMP2 immuno protein antibody: tumor uptake is associated with MMP2 activity. <i>Scientific Reports</i> , 2016, 6, 22198.	3.3	8
36	Inferring transcription factor activity from microarray data reveals novel targets for toxicological investigations. <i>Toxicology</i> , 2017, 389, 101-107.	4.2	7

#	ARTICLE	IF	CITATIONS
37	Translational regulation is a key determinant of the cellular response to benzo[a]pyrene. <i>Toxicology Letters</i> , 2018, 295, 144-152.	0.8	6
38	A cross-omics approach to investigate temporal gene expression regulation by 5-hydroxymethylcytosine via TBH-derived oxidative stress showed involvement of different regulatory kinases. <i>Toxicology in Vitro</i> , 2018, 48, 318-328.	2.4	4
39	Systems biology approaches to interpreting genomic data. <i>Current Opinion in Toxicology</i> , 2019, 18, 1-7.	5.0	3
40	Phosphorylation of eIF2 β promotes cell survival in response to benzo[a]pyrene exposure. <i>Toxicology in Vitro</i> , 2019, 54, 330-337.	2.4	2
41	Persistent transcriptional responses show the involvement of feed-forward control in a repeated dose toxicity study. <i>Toxicology</i> , 2017, 375, 58-63.	4.2	1
42	RNF8-Independent Lys63 Poly-Ubiquitylation Prevents Genomic Instability in Response to Replication-Associated DNA Damage. <i>PLoS ONE</i> , 2014, 9, e89997.	2.5	1