

Christian Bogdan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3496651/publications.pdf>

Version: 2024-02-01

46
papers

5,146
citations

147801

31
h-index

223800

46
g-index

47
all docs

47
docs citations

47
times ranked

7157
citing authors

#	ARTICLE	IF	CITATIONS
1	Cytokine-Mediated Regulation of ARG1 in Macrophages and Its Impact on the Control of Salmonella enterica Serovar Typhimurium Infection. <i>Cells</i> , 2021, 10, 1823.	4.1	15
2	Macrophages as host, effector and immunoregulatory cells in leishmaniasis: Impact of tissue micro-environment and metabolism. <i>Cytokine: X</i> , 2020, 2, 100041.	1.4	58
3	Arginase impedes the resolution of colitis by altering the microbiome and metabolome. <i>Journal of Clinical Investigation</i> , 2020, 130, 5703-5720.	8.2	44
4	Resolution of Cutaneous Leishmaniasis and Persistence of <i>Leishmania major</i> in the Absence of Arginase 1. <i>Journal of Immunology</i> , 2019, 202, 1453-1464.	0.8	25
5	SPIONs functionalized with small peptides for binding of lipopolysaccharide, a pathophysiologically relevant microbial product. <i>Colloids and Surfaces B: Biointerfaces</i> , 2019, 174, 95-102.	5.0	6
6	Transcription factor Fra-1 targets arginase-1 to enhance macrophage-mediated inflammation in arthritis. <i>Journal of Clinical Investigation</i> , 2019, 129, 2669-2684.	8.2	51
7	Rhinophyma-Like Cutaneous Leishmaniasis due to <i>Leishmania aethiopica</i> Treated Successfully with Liposomal Amphotericin B. <i>American Journal of Tropical Medicine and Hygiene</i> , 2019, 100, 231-232.	1.4	4
8	The Brief Case: Cutaneous Sporotrichosis in an Immunocompetent Patient after Travel to Peru. <i>Journal of Clinical Microbiology</i> , 2018, 56, .	3.9	2
9	Monocyte-Derived Signals Activate Human Natural Killer Cells in Response to <i>Leishmania</i> Parasites. <i>Frontiers in Immunology</i> , 2018, 9, 24.	4.8	18
10	Type I Interferon Signaling Is Required for CpG-Oligodesoxynucleotide-Induced Control of <i>Leishmania major</i> , but Not for Spontaneous Cure of Subcutaneous Primary or Secondary <i>L. major</i> Infection. <i>Frontiers in Immunology</i> , 2018, 9, 79.	4.8	25
11	Characterization of the Protein Tyrosine Phosphatase LmpRL-1 Secreted by <i>Leishmania major</i> via the Exosome Pathway. <i>Infection and Immunity</i> , 2017, 85, .	2.2	34
12	Function of Macrophage and Parasite Phosphatases in Leishmaniasis. <i>Frontiers in Immunology</i> , 2017, 8, 1838.	4.8	47
13	TNF-Mediated Restriction of Arginase 1 Expression in Myeloid Cells Triggers Type 2 NO Synthase Activity at the Site of Infection. <i>Cell Reports</i> , 2016, 15, 1062-1075.	6.4	102
14	Nitric oxide synthase in innate and adaptive immunity: an update. <i>Trends in Immunology</i> , 2015, 36, 161-178.	6.8	657
15	Fatal Leishmaniasis in the Absence of TNF Despite a Strong Th1 Response. <i>Frontiers in Microbiology</i> , 2015, 6, 1520.	3.5	36
16	Hypoxia in <i>Leishmania major</i> Skin Lesions Impairs the NO-Dependent Leishmanicidal Activity of Macrophages. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2339-2346.	0.7	59
17	Natural killer cells in experimental and human leishmaniasis. <i>Frontiers in Cellular and Infection Microbiology</i> , 2012, 2, 69.	3.9	68
18	Leishmaniasis in rheumatology, haematology and oncology: epidemiological, immunological and clinical aspects and caveats: Figure 1. <i>Annals of the Rheumatic Diseases</i> , 2012, 71, i60-i66.	0.9	71

#	ARTICLE	IF	CITATIONS
19	Leishmania-Infected Macrophages Are Targets of NK Cell-Derived Cytokines but Not of NK Cell Cytotoxicity. <i>Infection and Immunity</i> , 2011, 79, 2699-2708.	2.2	36
20	IL-18, but not IL-15, contributes to the IL-12-dependent induction of NK cell effector functions by <i>Leishmania infantum</i> in vivo. <i>European Journal of Immunology</i> , 2010, 40, 1708-1717.	2.9	45
21	Regulation of Lymphocytes by Nitric Oxide. <i>Methods in Molecular Biology</i> , 2010, 677, 375-393.	0.9	77
22	Endothelial nitric oxide synthase limits the inflammatory response in mouse cutaneous leishmaniasis. <i>Immunobiology</i> , 2010, 215, 826-832.	1.9	25
23	Generation, Culture and Flow-Cytometric Characterization of Primary Mouse Macrophages. <i>Methods in Molecular Biology</i> , 2009, 531, 203-224.	0.9	55
24	Toll-like receptor-induced arginase 1 in macrophages thwarts effective immunity against intracellular pathogens. <i>Nature Immunology</i> , 2008, 9, 1399-1406.	14.5	558
25	Mechanisms and consequences of persistence of intracellular pathogens: leishmaniasis as an example. <i>Cellular Microbiology</i> , 2008, 10, 1221-1234.	2.1	132
26	The innate immune response against <i>Leishmania</i> parasites. <i>Immunobiology</i> , 2008, 213, 377-387.	1.9	142
27	NK cell activation in visceral leishmaniasis requires TLR9, myeloid DCs, and IL-12, but is independent of plasmacytoid DCs. <i>Journal of Experimental Medicine</i> , 2007, 204, 893-906.	8.5	168
28	TLR9 signaling is essential for the innate NK cell response in murine cutaneous leishmaniasis. <i>European Journal of Immunology</i> , 2007, 37, 3424-3434.	2.9	140
29	Cytokine-mediated control of lipopolysaccharide-induced activation of small intestinal epithelial cells. <i>Immunology</i> , 2007, 122, 306-315.	4.4	33
30	Minute numbers of contaminant CD8+ T cells or CD11b+CD11c+ NK cells are the source of IFN- γ in IL-12/IL-18-stimulated mouse macrophage populations. <i>Blood</i> , 2005, 105, 1319-1328.	1.4	86
31	Translational Control of Inducible Nitric Oxide Synthase by IL-13 and Arginine Availability in Inflammatory Macrophages. <i>Journal of Immunology</i> , 2003, 171, 4561-4568.	0.8	160
32	Response to 'Species differences in macrophage NO production are important'. <i>Nature Immunology</i> , 2002, 3, 102-102.	14.5	21
33	Visceral Leishmaniasis in a German Child Who Had Never Entered a Known Endemic Area: Case Report and Review of the Literature. <i>Clinical Infectious Diseases</i> , 2001, 32, 302-306.	5.8	125
34	The Production of IFN- γ by IL-12/IL-18-Activated Macrophages Requires STAT4 Signaling and Is Inhibited by IL-4. <i>Journal of Immunology</i> , 2001, 166, 3075-3082.	0.8	168
35	Regulation of type 2 nitric oxide synthase by type 1 interferons in macrophages infected with <i>Leishmania major</i> . <i>European Journal of Immunology</i> , 2000, 30, 2257-2267.	2.9	58
36	The role of nitric oxide in innate immunity. <i>Immunological Reviews</i> , 2000, 173, 17-26.	6.0	572

#	ARTICLE	IF	CITATIONS
37	Fibroblasts as Host Cells in Latent Leishmaniasis. <i>Journal of Experimental Medicine</i> , 2000, 191, 2121-2130.	8.5	193
38	IFN- γ inhibits the production of latent transforming growth factor- β 1 by mouse inflammatory macrophages. <i>European Journal of Immunology</i> , 1998, 28, 1181-1188.	2.9	15
39	Type 1 Interferon (IFN- α / β) and Type 2 Nitric Oxide Synthase Regulate the Innate Immune Response to a Protozoan Parasite. <i>Immunity</i> , 1998, 8, 77-87.	14.3	354
40	The immune response to Leishmania: mechanisms of parasite control and evasion. <i>International Journal for Parasitology</i> , 1998, 28, 121-134.	3.1	246
41	2,4-diamino-6-hydroxypyrimidine, an inhibitor of tetrahydrobiopterin synthesis, downregulates the expression of iNOS protein and mRNA in primary murine macrophages. <i>FEBS Letters</i> , 1995, 363, 69-74.	2.8	27
42	Cytokine interactions in experimental cutaneous leishmaniasis. Interleukin 4 synergizes with interferon- γ to activate murine macrophages for killing of <i>Leishmania major</i> amastigotes. <i>European Journal of Immunology</i> , 1991, 21, 327-333.	2.9	88
43	Cytokine interactions in experimental cutaneous leishmaniasis. II. Endogenous tumor necrosis factor- α production by macrophages is induced by the synergistic action of interferon (IFN)- γ and interleukin (IL) 4 and accounts for the antiparasitic effect mediated by IFN- γ and IL 4. <i>European Journal of Immunology</i> , 1991, 21, 1669-1675.	2.9	60
44	Tumor necrosis factor- α in combination with interferon- γ , but not with interleukin 4 activates murine macrophages for elimination of <i>Leishmania major</i> amastigotes. <i>European Journal of Immunology</i> , 1990, 20, 1131-1135.	2.9	185
45	Immunization of susceptible hosts with a soluble antigen fraction from <i>Leishmania major</i> leads to aggravation of murine leishmaniasis mediated by CD4+ T cells. <i>European Journal of Immunology</i> , 1990, 20, 2533-2540.	2.9	28
46	Detection of Potentially Diagnostic Leishmanial Antigens by Western Blot Analysis of Sera from Patients with Kala-Azar or Multilesional Cutaneous Leishmaniasis. <i>Journal of Infectious Diseases</i> , 1990, 162, 1417-1418.	4.0	27