

Isabel Marzo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3504275/publications.pdf>

Version: 2024-02-01

101
papers

12,133
citations

57758

44
h-index

34986

98
g-index

105
all docs

105
docs citations

105
times ranked

12095
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Harnessing the Potential of NK Cell-Based Immunotherapies against Multiple Myeloma. <i>Cells</i> , 2022, 11, 392. | 4.1 | 7 |
| 2 | Metformin sensitizes leukemic cells to cytotoxic lymphocytes by increasing expression of intercellular adhesion molecule-1 (ICAM-1). <i>Scientific Reports</i> , 2022, 12, 1341. | 3.3 | 11 |
| 3 | Preclinical Studies of Granulysin-Based Anti-MUC1-Tn Immunotoxins as a New Antitumoral Treatment. <i>Biomedicines</i> , 2022, 10, 1223. | 3.2 | 2 |
| 4 | Multifunctional Heterometallic Ir ^{III} -Au ^I Probes as Promising Anticancer and Antiangiogenic Agents. <i>Chemistry - A European Journal</i> , 2021, 27, 9885-9897. | 3.3 | 17 |
| 5 | Future prospects for mitosis-targeted antitumor therapies. <i>Biochemical Pharmacology</i> , 2021, 190, 114655. | 4.4 | 24 |
| 6 | Dual Emissive Ir(III) Complexes for Photodynamic Therapy and Bioimaging. <i>Pharmaceutics</i> , 2021, 13, 1382. | 4.5 | 9 |
| 7 | Expanded NK cells from umbilical cord blood and adult peripheral blood combined with daratumumab are effective against tumor cells from multiple myeloma patients. <i>Oncolmmunology</i> , 2021, 10, 1853314. | 4.6 | 24 |
| 8 | Synthesis of New Thiourea-Metal Complexes with Promising Anticancer Properties. <i>Molecules</i> , 2021, 26, 6891. | 3.8 | 13 |
| 9 | Novel ureido-dihydropyridine scaffolds as theranostic agents. <i>Bioorganic Chemistry</i> , 2020, 105, 104364. | 4.1 | 5 |
| 10 | Expanded and activated allogeneic NK cells are cytotoxic against B-chronic lymphocytic leukemia (B-CLL) cells with sporadic cases of resistance. <i>Scientific Reports</i> , 2020, 10, 19398. | 3.3 | 23 |
| 11 | Heterobimetallic propargyl gold complexes with π -bound copper or silver with enhanced anticancer activity. <i>Dalton Transactions</i> , 2020, 49, 11736-11742. | 3.3 | 15 |
| 12 | Luminescent Bimetallic Ir ^{III} /Au ^I Peptide Bioconjugates as Potential Theranostic Agents. <i>Chemistry - A European Journal</i> , 2020, 26, 12085-12085. | 3.3 | 1 |
| 13 | Luminescent Re(I)/Au(I) Species As Selective Anticancer Agents for HeLa Cells. <i>Inorganic Chemistry</i> , 2020, 59, 8960-8970. | 4.0 | 24 |
| 14 | Novel Forms of Immunomodulation for Cancer Therapy. <i>Trends in Cancer</i> , 2020, 6, 518-532. | 7.4 | 17 |
| 15 | Luminescent Bimetallic Ir ^{III} /Au ^I Peptide Bioconjugates as Potential Theranostic Agents. <i>Chemistry - A European Journal</i> , 2020, 26, 12158-12167. | 3.3 | 19 |
| 16 | Immunogenic Cell Death and Immunotherapy of Multiple Myeloma. <i>Frontiers in Cell and Developmental Biology</i> , 2019, 7, 50. | 3.7 | 139 |
| 17 | Bioactive and luminescent indole and isatin based gold(I) derivatives. <i>Dalton Transactions</i> , 2019, 48, 3098-3108. | 3.3 | 17 |
| 18 | Response: Commentary: Immunogenic Cell Death and Immunotherapy of Multiple Myeloma. <i>Frontiers in Cell and Developmental Biology</i> , 2019, 7, 306. | 3.7 | 4 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Study of the anticancer properties of optically active titanocene oximate compounds. <i>Journal of Organometallic Chemistry</i> , 2019, 881, 150-158. | 1.8 | 5 |
| 20 | Biological evaluation of water soluble arene Ru(II) enantiomers with amino-oxime ligands. <i>Journal of Inorganic Biochemistry</i> , 2018, 183, 32-42. | 3.5 | 12 |
| 21 | Bioactive Heterobimetallic Re(I)/Au(I) Complexes Containing Bidentate N-Heterocyclic Carbenes. <i>Organometallics</i> , 2018, 37, 3993-4001. | 2.3 | 27 |
| 22 | Ylide Ligands as Building Blocks for Bioactive Group 11 Metal Complexes. <i>Chemistry - A European Journal</i> , 2018, 24, 11693-11702. | 3.3 | 15 |
| 23 | Multifaceted anticancer activity of BH3 mimetics: Current evidence and future prospects. <i>Biochemical Pharmacology</i> , 2017, 136, 12-23. | 4.4 | 52 |
| 24 | Highly active group 11 metal complexes with β -hydrazidophosphonate ligands. <i>Dalton Transactions</i> , 2017, 46, 13745-13755. | 3.3 | 13 |
| 25 | Trackable Metallodrugs Combining Luminescent Re(I) and Bioactive Au(I) Fragments. <i>Inorganic Chemistry</i> , 2017, 56, 15159-15170. | 4.0 | 48 |
| 26 | Synthesis of luminescent squaramide monoesters: cytotoxicity and cell imaging studies in HeLa cells. <i>RSC Advances</i> , 2016, 6, 14171-14177. | 3.6 | 21 |
| 27 | High-order TRAIL oligomer formation in TRAIL-coated lipid nanoparticles enhances DR5 cross-linking and increases antitumour effect against colon cancer. <i>Cancer Letters</i> , 2016, 383, 250-260. | 7.2 | 42 |
| 28 | Inhibition of autophagy with chloroquine potentiates carfilzomib-induced apoptosis in myeloma cells in vitro and in vivo. <i>Cancer Letters</i> , 2016, 382, 1-10. | 7.2 | 74 |
| 29 | Cytotoxicity and biodistribution studies of luminescent Au(I) and Ag(I) N-heterocyclic carbenes. Searching for new biological targets. <i>Dalton Transactions</i> , 2016, 45, 15026-15033. | 3.3 | 58 |
| 30 | MLL-Rearranged Acute Lymphoblastic Leukemias Activate BCL-2 through H3K79 Methylation and Are Sensitive to the BCL-2-Specific Antagonist ABT-199. <i>Cell Reports</i> , 2015, 13, 2715-2727. | 6.4 | 118 |
| 31 | Cyclometalated Iminophosphorane Gold(III) and Platinum(II) Complexes. A Highly Permeable Cationic Platinum(II) Compound with Promising Anticancer Properties. <i>Journal of Medicinal Chemistry</i> , 2015, 58, 5825-5841. | 6.4 | 88 |
| 32 | Bcl-2 family of proteins as drug targets for cancer chemotherapy: the long way of BH3 mimetics from bench to bedside. <i>Current Opinion in Pharmacology</i> , 2015, 23, 74-81. | 3.5 | 75 |
| 33 | Immunotherapy with liposome-bound TRAIL overcomes partial protection to soluble TRAIL-induced apoptosis offered by down-regulation of Bim in leukemic cells. <i>Clinical and Translational Oncology</i> , 2015, 17, 657-667. | 2.4 | 27 |
| 34 | Human NK cells activated by EBV-transformed lymphoblastoid cells overcome anti-apoptotic mechanisms of drug resistance in haematological cancer cells. <i>Oncotarget</i> , 2015, 4, e991613. | 4.6 | 36 |
| 35 | Highly Cytotoxic Bioconjugated Gold(I) Complexes with Cysteine-Containing Dipeptides. <i>Chemistry - A European Journal</i> , 2015, 21, 11088-11095. | 3.3 | 49 |
| 36 | Two death pathways induced by sorafenib in myeloma cells: Puma-mediated apoptosis and necroptosis. <i>Clinical and Translational Oncology</i> , 2015, 17, 121-132. | 2.4 | 21 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | In Vitro and in Vivo Evaluation of Water-Soluble Iminophosphorane Ruthenium(II) Compounds. A Potential Chemotherapeutic Agent for Triple Negative Breast Cancer. <i>Journal of Medicinal Chemistry</i> , 2014, 57, 9995-10012. | 6.4 | 87 |
| 38 | Granulysin induces apoptotic cell death and cleavage of the autophagy regulator Atg5 in human hematological tumors. <i>Biochemical Pharmacology</i> , 2014, 87, 410-423. | 4.4 | 29 |
| 39 | Synthesis, Characterization, and Cytotoxic Activity of Au(I),S-Heterocyclic Carbenes Derived from Peptides Containing L-Thiazolylalanine. <i>European Journal of Inorganic Chemistry</i> , 2014, 2014, 2512-2519. | 2.0 | 35 |
| 40 | Gold(I) thiolates containing amino acid moieties. Cytotoxicity and structure-activity relationship studies. <i>Dalton Transactions</i> , 2014, 43, 17054-17066. | 3.3 | 33 |
| 41 | Luminescent Re(I) and Re(I)/Au(I) complexes as cooperative partners in cell imaging and cancer therapy. <i>Chemical Science</i> , 2014, 5, 4434-4446. | 7.4 | 74 |
| 42 | Antimitotic drugs in cancer chemotherapy: Promises and pitfalls. <i>Biochemical Pharmacology</i> , 2013, 86, 703-710. | 4.4 | 72 |
| 43 | Liposomes Decorated with Apo2L/TRAIL Overcome Chemoresistance of Human Hematologic Tumor Cells. <i>Molecular Pharmaceutics</i> , 2013, 10, 893-904. | 4.6 | 70 |
| 44 | Direct Interaction of Bax and Bak Proteins with Bcl-2 Homology Domain 3 (BH3)-only Proteins in Living Cells Revealed by Fluorescence Complementation. <i>Journal of Biological Chemistry</i> , 2013, 288, 4935-4946. | 3.4 | 74 |
| 45 | Acute Lymphoblastic Leukemia Is a Bcl-2 Dependent Disease: Proteomic Profiling and Pre-Clinical Efficacy Of a Selective Bcl-2 Antagonist ABT-199. <i>Blood</i> , 2013, 122, 3919-3919. | 1.4 | 2 |
| 46 | Organometallic Palladium Complexes with a Water-Soluble Iminophosphorane Ligand As Potential Anticancer Agents. <i>Organometallics</i> , 2012, 31, 5772-5781. | 2.3 | 70 |
| 47 | Cytotoxic hydrophilic iminophosphorane coordination compounds of d8 metals. Studies of their interactions with DNA and HSA. <i>Journal of Inorganic Biochemistry</i> , 2012, 116, 204-214. | 3.5 | 56 |
| 48 | Targeting the Apo2L/TRAIL system for the therapy of autoimmune diseases and cancer. <i>Biochemical Pharmacology</i> , 2012, 83, 1475-1483. | 4.4 | 45 |
| 49 | Bortezomib resistance in a myeloma cell line is associated to PSM β 5 overexpression and polyploidy. <i>Leukemia Research</i> , 2012, 36, 212-218. | 0.8 | 75 |
| 50 | Cell fate after mitotic arrest in different tumor cells is determined by the balance between slippage and apoptotic threshold. <i>Toxicology and Applied Pharmacology</i> , 2012, 258, 384-393. | 2.8 | 24 |
| 51 | Iminophosphorane-organogold(III) complexes induce cell death through mitochondrial ROS production. <i>Journal of Inorganic Biochemistry</i> , 2011, 105, 1306-1313. | 3.5 | 57 |
| 52 | Bim is the key mediator of glucocorticoid-induced apoptosis and of its potentiation by rapamycin in human myeloma cells. <i>Biochimica Et Biophysica Acta - Molecular Cell Research</i> , 2010, 1803, 311-322. | 4.1 | 19 |
| 53 | Different contribution of BH3-only proteins and caspases to doxorubicin-induced apoptosis in p53-deficient leukemia cells. <i>Biochemical Pharmacology</i> , 2010, 79, 1746-1758. | 4.4 | 26 |
| 54 | Detection of <i>Clostridium tyrobutyricum</i> spores using polyclonal antibodies and flow cytometry. <i>Journal of Applied Microbiology</i> , 2010, 108, 488-498. | 3.1 | 17 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Cooperation between Apo2L/TRAIL and bortezomib in multiple myeloma apoptosis. <i>Biochemical Pharmacology</i> , 2009, 77, 804-812. | 4.4 | 51 |
| 56 | Bcl-2 family members as molecular targets in cancer therapy. <i>Biochemical Pharmacology</i> , 2008, 76, 939-946. | 4.4 | 75 |
| 57 | Exposure of any of two proapoptotic domains of presenilin 1-associated protein/mitochondrial carrier homolog 1 on the surface of mitochondria is sufficient for induction of apoptosis in a Bax/Bak-independent manner. <i>European Journal of Cell Biology</i> , 2008, 87, 325-334. | 3.6 | 18 |
| 58 | Individual Variation of Scavenger Receptor Expression in Human Macrophages with Oxidized Low-Density Lipoprotein Is Associated with a Differential Inflammatory Response. <i>Journal of Immunology</i> , 2007, 179, 3242-3248. | 0.8 | 64 |
| 59 | Conjugation of a novel Apaf-1 inhibitor to peptide-based cell-membrane transporters. <i>Peptides</i> , 2007, 28, 958-968. | 2.4 | 31 |
| 60 | Apoptosis by IL-2 deprivation in human CD8+ T cell blasts predominates over death receptor ligation, requires Bim expression and is associated with Mcl-1 loss. <i>Molecular Immunology</i> , 2007, 44, 1446-1453. | 2.2 | 18 |
| 61 | Mechanism of apoptosis induced by IFN- γ in human myeloma cells: Role of Jak1 and Bim and potentiation by rapamycin. <i>Cellular Signalling</i> , 2007, 19, 844-854. | 3.6 | 38 |
| 62 | Membrane expression of DR4, DR5 and caspase-8 levels, but not Mcl-1, determine sensitivity of human myeloma cells to Apo2L/TRAIL. <i>Experimental Cell Research</i> , 2007, 313, 2378-2388. | 2.6 | 53 |
| 63 | Human CD8+ T α cell blasts are more sensitive than CD4+ T α cell blasts to regulation by APO2L/TRAIL. <i>European Journal of Immunology</i> , 2005, 35, 1812-1821. | 2.9 | 27 |
| 64 | Farnesyltransferase Inhibitor BMS-214662 Induces Apoptosis in Myeloma Cells through PUMA Up-Regulation, Bax and Bak Activation, and Mcl-1 Elimination. <i>Molecular Pharmacology</i> , 2005, 67, 1991-1998. | 2.3 | 34 |
| 65 | Down-regulation of normal human T cell blast activation: roles of APO2L/TRAIL, FasL, and c-FLIP, Bim, or Bcl-x isoform expression. <i>Journal of Leukocyte Biology</i> , 2005, 77, 568-578. | 3.3 | 37 |
| 66 | Apo2L/TRAIL is an indirect mediator of apoptosis induced by interferon- γ in human myeloma cells. <i>FEBS Letters</i> , 2005, 579, 6217-6222. | 2.8 | 20 |
| 67 | Farnesyltransferase inhibitor BMS-214662 induces apoptosis in B-cell chronic lymphocytic leukemia cells. <i>Leukemia</i> , 2004, 18, 1599-1604. | 7.2 | 17 |
| 68 | Role of caspases and apoptosis-inducing factor (AIF) in cladribine-induced apoptosis of B cell chronic lymphocytic leukemia. <i>Leukemia</i> , 2002, 16, 2106-2114. | 7.2 | 36 |
| 69 | Cladribine induces apoptosis in human leukaemia cells by caspase-dependent and -independent pathways acting on mitochondria. <i>Biochemical Journal</i> , 2001, 359, 537-546. | 3.7 | 83 |
| 70 | A Role of the Mitochondrial Apoptosis-Inducing Factor in Granulysin-Induced Apoptosis. <i>Journal of Immunology</i> , 2001, 167, 1222-1229. | 0.8 | 103 |
| 71 | Cladribine induces apoptosis in human leukaemia cells by caspase-dependent and -independent pathways acting on mitochondria. <i>Biochemical Journal</i> , 2001, 359, 537. | 3.7 | 52 |
| 72 | Purification and Liposomal Reconstitution of Permeability Transition Pore Complex. <i>Methods in Enzymology</i> , 2000, 322, 243-252. | 1.0 | 15 |

| # | ARTICLE | IF | CITATIONS |
|----|--|------|-----------|
| 73 | Bcl-2 and Bax regulate the channel activity of the mitochondrial adenine nucleotide translocator. <i>Oncogene</i> , 2000, 19, 329-336. | 5.9 | 322 |
| 74 | Mitochondrial Release of Caspase-2 and -9 during the Apoptotic Process. <i>Journal of Experimental Medicine</i> , 1999, 189, 381-394. | 8.5 | 678 |
| 75 | Molecular characterization of mitochondrial apoptosis-inducing factor. <i>Nature</i> , 1999, 397, 441-446. | 27.8 | 3,697 |
| 76 | Lonidamine triggers apoptosis via a direct, Bcl-2-inhibited effect on the mitochondrial permeability transition pore. <i>Oncogene</i> , 1999, 18, 2537-2546. | 5.9 | 194 |
| 77 | Arsenite Induces Apoptosis via a Direct Effect on the Mitochondrial Permeability Transition Pore. <i>Experimental Cell Research</i> , 1999, 249, 413-421. | 2.6 | 283 |
| 78 | Mitochondrial permeability transition in apoptosis and necrosis. <i>Cell Biology and Toxicology</i> , 1998, 14, 141-145. | 5.3 | 121 |
| 79 | The thiol crosslinking agent diamide overcomes the apoptosis-inhibitory effect of Bcl-2 by enforcing mitochondrial permeability transition. <i>Oncogene</i> , 1998, 16, 1055-1063. | 5.9 | 149 |
| 80 | Subcellular and submitochondrial mode of action of Bcl-2-like oncoproteins. <i>Oncogene</i> , 1998, 16, 2265-2282. | 5.9 | 385 |
| 81 | Editorial. <i>Experimental Gerontology</i> , 1998, 33, 543-553. | 2.8 | 25 |
| 82 | Cytofluorometric detection of mitochondrial alterations in early CD95/Fas/APO-1-triggered apoptosis of Jurkat T lymphoma cells. Comparison of seven mitochondrion-specific fluorochromes. <i>Immunology Letters</i> , 1998, 61, 157-163. | 2.5 | 195 |
| 83 | La mitochondrie, chef d'orchestre de la mort cellulaire. <i>Biofutur</i> , 1998, 1998, 32-36. | 0.0 | 0 |
| 84 | Bax and Adenine Nucleotide Translocator Cooperate in the Mitochondrial Control of Apoptosis. , 1998, 281, 2027-2031. | | 1,061 |
| 85 | Loss of Δ^6 -desaturase activity leads to impaired docosahexaenoic acid synthesis in Y-79 retinoblastoma cells. <i>Prostaglandins Leukotrienes and Essential Fatty Acids</i> , 1998, 59, 293-297. | 2.2 | 8 |
| 86 | The central role of the mitochondrial megachannel in apoptosis: evidence obtained with intact cells, isolated mitochondria, and purified protein complexes. <i>Biomedicine and Pharmacotherapy</i> , 1998, 52, 248-251. | 5.6 | 74 |
| 87 | Caspases disrupt mitochondrial membrane barrier function. <i>FEBS Letters</i> , 1998, 427, 198-202. | 2.8 | 123 |
| 88 | The Permeability Transition Pore Complex: A Target for Apoptosis Regulation by Caspases and Bcl-2-related Proteins. <i>Journal of Experimental Medicine</i> , 1998, 187, 1261-1271. | 8.5 | 657 |
| 89 | Coopération mortelle entre la protéine pro-apoptotique Bax et le translocateur Δ adénine nucléotidique pour le contrôle mitochondrial de l'apoptose.. <i>Medecine/Sciences</i> , 1998, 14, 1399. | 0.2 | 0 |
| 90 | A Cytofluorometric Assay of Nuclear Apoptosis Induced in a Cell-Free System: Application to Ceramide-Induced Apoptosis. <i>Experimental Cell Research</i> , 1997, 236, 397-403. | 2.6 | 73 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 91 | Nitric oxide induces apoptosis via triggering mitochondrial permeability transition. FEBS Letters, 1997, 410, 373-377. | 2.8 | 220 |
| 92 | The apoptosis-necrosis paradox. Apoptogenic proteases activated after mitochondrial permeability transition determine the mode of cell death. Oncogene, 1997, 15, 1573-1581. | 5.9 | 443 |
| 93 | Role of the Mitochondrial Permeability Transition Pore in Apoptosis. Bioscience Reports, 1997, 17, 67-76. | 2.4 | 193 |
| 94 | Glutathione depletion is an early and calcium elevation is a late event of thymocyte apoptosis. Journal of Immunology, 1997, 158, 4612-9. | 0.8 | 205 |
| 95 | CPP32 inhibition prevents Fas-induced ceramide generation and apoptosis in human cells. FEBS Letters, 1996, 390, 233-237. | 2.8 | 78 |
| 96 | Biosynthesis of docosahexaenoic acid in human cells: evidence that two different Δ^6 -desaturase activities may exist. Lipids and Lipid Metabolism, 1996, 1301, 263-272. | 2.6 | 47 |
| 97 | Biosynthesis of unsaturated fatty acids in the main cell lineages of human leukemia and lymphoma. Lipids and Lipid Metabolism, 1995, 1257, 140-148. | 2.6 | 33 |
| 98 | mtDNA-depleted U937 cells are sensitive to TNF and Fas-mediated cytotoxicity. FEBS Letters, 1995, 376, 15-18. | 2.8 | 32 |
| 99 | Self-Staining of Polyunsaturated Fatty Acids in Argentation Thin-Layer Chromatography. Analytical Biochemistry, 1994, 220, 210-212. | 2.4 | 17 |
| 100 | Alternative route for the biosynthesis of polyunsaturated fatty acids in K562 cells. Biochemical Journal, 1993, 291, 841-845. | 3.7 | 25 |
| 101 | Synthesis and antiproliferative study of phosphorescent multimetallic Re(I)/Au(I) complexes containing fused imidazo[4,5-f]quinoxaline core. Applied Organometallic Chemistry, 0, , . | 3.5 | 4 |