

Andreas Peter

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/350468/publications.pdf>

Version: 2024-02-01

42
papers

2,094
citations

279798

23
h-index

289244

40
g-index

43
all docs

43
docs citations

43
times ranked

5241
citing authors

#	ARTICLE	IF	CITATIONS
1	Testing for SARS-CoV-2 seroprevalence: experiences of a tertiary eye centre. <i>BMJ Open Ophthalmology</i> , 2021, 6, e000688.	1.6	0
2	An Activity Trackerâ€œGuided Physical Activity Program for Patients Undergoing Radiotherapy: Protocol for a Prospective Phase III Trial (OnkoFit I and II Trials). <i>JMIR Research Protocols</i> , 2021, 10, e28524.	1.0	1
3	Elevated circulating follistatin associates with an increased risk of type 2 diabetes. <i>Nature Communications</i> , 2021, 12, 6486.	12.8	31
4	Intensifying sleep slow oscillations does not improve metabolic control in healthy men. <i>Psychoneuroendocrinology</i> , 2019, 99, 1-7.	2.7	10
5	Sex-Specific Associations of Testosterone With Metabolic Traits. <i>Frontiers in Endocrinology</i> , 2019, 10, 90.	3.5	13
6	The Gly385(388)Arg Polymorphism of the FGFR4 Receptor Regulates Hepatic Lipogenesis Under Healthy Diet. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019, 104, 2041-2053.	3.6	8
7	Effects of resveratrol supplementation on liver fat content in overweight and insulinâ€resistant subjects: A randomized, doubleâ€blind, placeboâ€controlled clinical trial. <i>Diabetes, Obesity and Metabolism</i> , 2018, 20, 1793-1797.	4.4	66
8	Androgen receptor overexpression in prostate cancer in type 2 diabetes. <i>Molecular Metabolism</i> , 2018, 8, 158-166.	6.5	22
9	Mitotic figure in the peripheral blood smear. <i>American Journal of Hematology</i> , 2018, 93, 154-154.	4.1	0
10	The hepatokines fetuin-A and fetuin-B are upregulated in the state of hepatic steatosis and may differently impact on glucose homeostasis in humans. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2018, 314, E266-E273.	3.5	56
11	cGMP-dependent protein kinase I (cGKI) modulates human hepatic stellate cell activation. <i>Metabolism: Clinical and Experimental</i> , 2018, 88, 22-30.	3.4	18
12	Single Nucleotide Polymorphisms in the G-Protein Coupled Receptor Kinase 5 (GRK5) Gene are associated with Plasma LDL-Cholesterol Levels in Humans. <i>Scientific Reports</i> , 2018, 8, 7745.	3.3	3
13	Effect of increased enteral protein intake on plasma and urinary urea concentrations in preterm infants born at ≥ 32 weeks gestation and ≤ 1500 g birth weight enrolled in a randomized controlled trial â€œ a secondary analysis. <i>BMC Pediatrics</i> , 2018, 18, 154.		17
14	Unusual high blood glucose in ketoacidosis as first presentation of type 1 diabetes mellitus. <i>Endocrinology, Diabetes and Metabolism Case Reports</i> , 2018, 2018, .	0.5	3
15	Soluble urokinase receptor (suPAR) predicts microalbuminuria in patients at risk for type 2 diabetes mellitus. <i>Scientific Reports</i> , 2017, 7, 40627.	3.3	40
16	Point-of-care testing for emergency assessment of coagulation in patients treated with direct oral anticoagulants. <i>Critical Care</i> , 2017, 21, 32.	5.8	58
17	Hypothalamic and Striatal Insulin Action Suppresses Endogenous Glucose Production and May Stimulate Glucose Uptake During Hyperinsulinemia in Lean but Not in Overweight Men. <i>Diabetes</i> , 2017, 66, 1797-1806.	0.6	87
18	Bezafibrate ameliorates diabetes via reduced steatosis and improved hepatic insulin sensitivity in diabetic TallyHo mice. <i>Molecular Metabolism</i> , 2017, 6, 256-266.	6.5	27

#	ARTICLE	IF	CITATIONS
19	Impact of end-stage renal disease on glucose metabolism—a matched cohort analysis. <i>Nephrology Dialysis Transplantation</i> , 2017, 32, 670-676.	0.7	22
20	Effect of Increased Enteral Protein Intake on Growth in Human Milk-Fed Preterm Infants. <i>JAMA Pediatrics</i> , 2017, 171, 16.	6.2	48
21	Diagnostic Accuracy of a Novel Chromogenic Direct Thrombin Inhibitor Assay: Clinical Experiences for Dabigatran Monitoring. <i>Thrombosis and Haemostasis</i> , 2017, 117, 2369-2375.	3.4	11
22	Relationship of Serum Trimethylamine N-Oxide (TMAO) Levels with early Atherosclerosis in Humans. <i>Scientific Reports</i> , 2016, 6, 26745.	3.3	224
23	The Genetic Variant I148M in <i>PNPLA3</i> Is Associated With Increased Hepatic Retinyl-Palmitate Storage in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2015, 100, E1568-E1574.	3.6	52
24	Urinary Neutrophil Gelatinase-Associated Lipocalin (NGAL) and proteinuria predict severity of acute kidney injury in Puumala virus infection. <i>BMC Infectious Diseases</i> , 2015, 15, 464.	2.9	22
25	Associations of Erythrocyte Fatty Acids in the De Novo Lipogenesis Pathway with Proxies of Liver Fat Accumulation in the EPIC-Potsdam Study. <i>PLoS ONE</i> , 2015, 10, e0127368.	2.5	25
26	Low-frequency and rare exome chip variants associate with fasting glucose and type 2 diabetes susceptibility. <i>Nature Communications</i> , 2015, 6, 5897.	12.8	173
27	Reticulocyte haemoglobin content as a marker of iron deficiency. <i>Archives of Disease in Childhood: Fetal and Neonatal Edition</i> , 2015, 100, F198-F202.	2.8	37
28	Point-of-Care Testing of Coagulation in Patients Treated With Non-Vitamin K Antagonist Oral Anticoagulants. <i>Stroke</i> , 2015, 46, 2741-2747.	2.0	62
29	The Brain Response to Peripheral Insulin Declines with Age: A Contribution of the Blood-Brain Barrier?. <i>PLoS ONE</i> , 2015, 10, e0126804.	2.5	80
30	Plasma Concentrations of the Vasoactive Peptide Fragments Mid-Regional Pro-Adrenomedullin, C-Terminal Pro-Endothelin 1 and Copeptin in Hemodialysis Patients: Associated Factors and Prediction of Mortality. <i>PLoS ONE</i> , 2014, 9, e86148.	2.5	17
31	Cinnamon Extract Improves Insulin Sensitivity in the Brain and Lowers Liver Fat in Mouse Models of Obesity. <i>PLoS ONE</i> , 2014, 9, e92358.	2.5	80
32	Comment on Hedderson et al. Prepregnancy SHBG Concentrations and Risk for Subsequently Developing Gestational Diabetes Mellitus. <i>Diabetes Care</i> 2014;37:1296-1303. <i>Diabetes Care</i> , 2014, 37, e278-e279.	8.6	1
33	Central Insulin Administration Improves Whole-Body Insulin Sensitivity via Hypothalamus and Parasympathetic Outputs in Men. <i>Diabetes</i> , 2014, 63, 4083-4088.	0.6	135
34	Lower Plasma Creatinine and Urine Albumin in Individuals at Increased Risk of Type 2 Diabetes with Factor V Leiden Mutation. <i>Isrn Endocrinology</i> , 2014, 2014, 1-3.	2.0	6
35	PNPLA3 variant I148M is associated with altered hepatic lipid composition in humans. <i>Diabetologia</i> , 2014, 57, 2103-2107.	6.3	41
36	Hepatic Glucokinase Expression Is Associated with Lipogenesis and Fatty Liver in Humans. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011, 96, E1126-E1130.	3.6	94

#	ARTICLE	IF	CITATIONS
37	Increased Cyclosporine Concentrations in the Absence of Cyclosporine Administration. <i>Clinical Chemistry</i> , 2011, 57, 670-673.	3.2	13
38	Relationships between hepatic stearoyl-CoA desaturase-1 activity and mRNA expression with liver fat content in humans. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2011, 300, E321-E326.	3.5	28
39	Relationships of Circulating Sex Hormone-Binding Globulin With Metabolic Traits in Humans. <i>Diabetes</i> , 2010, 59, 3167-3173.	0.6	130
40	Hepatic Lipid Composition and Stearoyl-Coenzyme A Desaturase 1 mRNA Expression Can Be Estimated from Plasma VLDL Fatty Acid Ratios. <i>Clinical Chemistry</i> , 2009, 55, 2113-2120.	3.2	113
41	Individual Stearoyl-CoA Desaturase 1 Expression Modulates Endoplasmic Reticulum Stress and Inflammation in Human Myotubes and Is Associated With Skeletal Muscle Lipid Storage and Insulin Sensitivity In Vivo. <i>Diabetes</i> , 2009, 58, 1757-1765.	0.6	134
42	Induction of stearoyl-CoA desaturase protects human arterial endothelial cells against lipotoxicity. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2008, 295, E339-E349.	3.5	86