In Kyoung Lim

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3544439/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1 | Role of TNF-α-Inducing Protein Secreted by Helicobacter pylori as a Tumor Promoter in Gastric Cancer and Emerging Preventive Strategies. Toxins, 2021, 13, 181. | 3.4 | 13 |
| 2 | Mitochondrial nucleoid remodeling and biogenesis are regulated by the p53-p21WAF1-PKCζ pathway in p16INK4a-silenced cells. Aging, 2020, 12, 6700-6732. | 3.1 | 7 |
| 3 | Translational downregulation of Twist1 expression by antiproliferative gene, B-cell translocation gene 2, in the triple negative breast cancer cells. Cell Death and Disease, 2019, 10, 410. | 6.3 | 9 |
| 4 | Inhibition of TNFα-interacting protein α (Tipα)-associated gastric carcinogenesis by BTG2/TIS21 via downregulating cytoplasmic nucleolin expression. Experimental and Molecular Medicine, 2018, 50, e449-e449. | 7.7 | 9 |
| 5 | Phylogenetic characterization of norovirus strains detected from sporadic gastroenteritis in Seoul during 2014–2016. Gut Pathogens, 2018, 10, 36. | 3.4 | 12 |
| 6 | TIS21/BTG2 inhibits breast cancer growth and progression by differential regulation of mTORc1 and mTORc2–AKT1–NFAT1–PHLPP2 signaling axis. Journal of Cancer Research and Clinical Oncology, 2018, 144, 1445-1462. | 2.5 | 9 |
| 7 | PPARα-Target Gene Expression Requires TIS21 Gene in Liver of the C57BL/6 Mice under Fasting Condition. Molecules and Cells, 2018, 41, 140-149. | 2.6 | 2 |
| 8 | TIS21/BTG2 inhibits doxorubicin-induced stress fiber-vimentin networks via Nox4-ROS-ABI2-DRF-linked signal cascade. Cellular Signalling, 2017, 30, 179-190. | 3.6 | 7 |
| 9 | Shifting p53-induced senescence to cell death by TIS21 /BTG2/Pc3 gene through posttranslational modification of p53 protein. Cellular Signalling, 2016, 28, 1172-1185. | 3.6 | 14 |
| 10 | Strong immunoexpression of midkine is associated with multiple lymph node metastases in BRAFV600E papillary thyroid carcinoma Journal of Clinical Oncology, 2016, 34, e17552-e17552. | 1.6 | 1 |
| 11 | Regulations of Reversal of Senescence by PKC Isozymes in Response to 12-O-Tetradecanoylphorbol-13-Acetate via Nuclear Translocation of pErk1/2. Molecules and Cells, 2016, 39, 266-279. | 2.6 | 4 |
| 12 | Stress-induced NF-κB activation differentiates promyelocytic leukemia cells to macrophages in response to all-trans-retinoic acid. Cellular Signalling, 2015, 27, 694-706. | 3.6 | 4 |
| 13 | Downregulation of PEA-15 reverses G1 arrest, and nuclear and chromatin changes of senescence phenotype via pErk1/2 translocation to nuclei. Cellular Signalling, 2015, 27, 1102-1109. | 3.6 | 8 |
| 14 | Protein Methylation and Interaction with the Antiproliferative Gene, BTG2 ^{/TIS21/Pc3} . Yonsei Medical Journal, 2014, 55, 292. | 2.2 | 4 |
| 15 | Sensitization of metformin-cytotoxicity by dichloroacetate via reprogramming glucose metabolism in cancer cells. Cancer Letters, 2014, 346, 300-308. | 7.2 | 52 |
| 16 | Câ€reactive protein induces G2/M phase cell cycle arrest and apoptosis in monocytes through the upregulation of Bâ€cell translocation gene 2 expression. FEBS Letters, 2014, 588, 625-631. | 2.8 | 19 |
| 17 | Inhibition of bladder cancer invasion by Sp1â€mediated <scp>BTG</scp> 2 expression via inhibition of <scp>DNA</scp> methyltransferaseÂ1. FEBS Journal, 2014, 281, 5581-5601. | 4.7 | 26 |
| 18 | Accumulation of cytolytic CD8+ T cells in B16-melanoma and proliferation of mature T cells in TIS21-knockout mice after T cell receptor stimulation. Experimental Cell Research, 2014, 327, 209-221. | 2.6 | 11 |

IN KYOUNG LIM

| # | Article | IF | CITATIONS |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | Signals regulating necrosis of cardiomyoblast by BTG2/TIS21/PC3 via activation of GSK3β and opening of mitochondrial permeability transition pore in response to H2O2. Biochemical and Biophysical Research Communications, 2013, 434, 559-565. | 2.1 | 11 |
| 20 | TIS21/BTG2 inhibits invadopodia formation by downregulating reactive oxygen species level in MDA-MB-231 cells. Journal of Cancer Research and Clinical Oncology, 2013, 139, 1657-1665. | 2.5 | 20 |
| 21 | B-cell translocation gene 2 mediates crosstalk between PI3K/Akt1 and NFκB pathways which enhances transcription of MnSOD by accelerating lκBα degradation in normal and cancer cells. Cell Communication and Signaling, 2013, 11, 69. | 6.5 | 23 |
| 22 | C-terminus-deleted FoxM1 is expressed in cancer cell lines and induces chromosome instability. Carcinogenesis, 2013, 34, 1907-1917. | 2.8 | 10 |
| 23 | TIS21/BTG2/PC3 inhibits interleukin-6 expression via downregulation of STAT3 pathway. Cellular Signalling, 2013, 25, 2391-2399. | 3.6 | 10 |
| 24 | B-cell translocation gene 2: Expression in the rat ovary and potential association with adenine nucleotide translocase 2 in mitochondria. Molecular and Cellular Endocrinology, 2013, 367, 31-40. | 3.2 | 5 |
| 25 | Regulation of Btg2/TIS21/PC3 expression via reactive oxygen species–protein kinase C–ÎFκΒ pathway under stress conditions. Cellular Signalling, 2013, 25, 2400-2412. | 3.6 | 42 |
| 26 | Abstract 1857: Dichloroacetate (DCA), pyruvate dehydrogenase kinase (PDK) inhibitor, enhances the cytotoxic effect induced by metformin , 2013, , . | | 0 |
| 27 | BTG2 suppresses cancer cell migration through inhibition of Src-FAK signaling by downregulation of reactive oxygen species generation in mitochondria. Clinical and Experimental Metastasis, 2012, 29, 901-913. | 3.3 | 39 |
| 28 | TIS21/BTG2/PC3 enhances downregulation of c-Myc during differentiation of HL-60 cells by activating Erk1/2 and inhibiting Akt in response to all-trans-retinoic acid. European Journal of Cancer, 2012, 48, 2474-2485. | 2.8 | 22 |
| 29 | TIS21/BTG2/PC3 accelerates the repair of DNA double strand breaks by enhancing Mre11 methylation and blocking damage signal transfer to the Chk2T68–p53S20 pathway. DNA Repair, 2012, 11, 965-975. | 2.8 | 28 |
| 30 | Abstract 495: Overexpression of TIS21/BTG2/PC3inhibits formation of invadopodia in the highly metastatic breast cancer cells MDA-MB-231. , 2012, , . | | 0 |
| 31 | Loss of p21Sdi1 expression in senescent cells after DNA damage accompanied with increase of miR-93 expression and reduced p53 interaction with p21Sdi1 gene promoter. Biochemical and Biophysical Research Communications, 2011, 407, 406-411. | 2.1 | 6 |
| 32 | Reduction of exportin 6 activity leads to actin accumulation via failure of RanGTP restoration and NTF2 sequestration in the nuclei of senescent cells. Experimental Cell Research, 2011, 317, 941-954. | 2.6 | 8 |
| 33 | Abstract 211: TIS21/BTG2/PC3triggers cancer cell death, instead of cellular senescence, by enhancing proapoptotic gene expression at the downstream of p53. , 2011, , . | | 0 |
| 34 | Abstract 1451: Inhibition of cell migrationviareduced phosphorylation of focal adhesion kinase by TIS21/BTG2/PC3in A549 cells. , 2011, , . | | 0 |
| 35 | Abstract 3112:TIS21BTG2/PC3enhances down-regulation of c-Myc expression during all-trans-retinoic acid-induced differentiation of acute promyelocytic leukemia cells. , 2010, , . | | 0 |
| 36 | Phosphorylated Extracellular Signal-regulated Protein Kinases 1 and 2 Phosphorylate Sp1 on Serine 59 and Regulate Cellular Senescence via Transcription of p21Sdi1/Cip1/Waf1. Journal of Biological Chemistry, 2009, 284, 15475-15486. | 3.4 | 32 |

IN KYOUNG LIM

| # | Article | IF | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 37 | Skp2 enhances polyubiquitination and degradation of TIS21/BTG2/PC3, tumor suppressor protein, at the downstream of FoxM1. Experimental Cell Research, 2009, 315, 3152-3162. | 2.6 | 20 |
| 38 | Friendship in research: the Japan–Korea symposia on cancer and ageing. Journal of Cancer Research and Clinical Oncology, 2008, 134, 813-817. | 2.5 | 0 |
| 39 | Pleiotrophin inhibits transforming growth factor β1â€induced apoptosis in hepatoma cell lines. Molecular Carcinogenesis, 2008, 47, 784-796. | 2.7 | 13 |
| 40 | TIS21 negatively regulates hepatocarcinogenesis by disruption of cyclin B1-Forkhead box M1 regulation loop. Hepatology, 2008, 47, 1533-1543. | 7.3 | 69 |
| 41 | TIS21/BTG2 Negatively Regulates Estradiol-Stimulated Expansion of Hematopoietic Stem Cells by Derepressing Akt Phosphorylation and Inhibiting mTOR Signal Transduction. Stem Cells, 2008, 26, 2339-2348. | 3.2 | 25 |
| 42 | Enhanced glycogenesis is involved in cellular senescence via GSK3/GS modulation. Aging Cell, 2008, 7, 894-907. | 6.7 | 73 |
| 43 | B Cell Translocation Gene 2 Enhances Susceptibility of HeLa Cells to Doxorubicin-induced Oxidative Damage. Journal of Biological Chemistry, 2008, 283, 33110-33118. | 3.4 | 34 |
| 44 | Establishing MD-PhD Programs in the Graduate School of Medicine in Korea. Journal of the Korean Medical Association, 2008, 51, 500. | 0.3 | 0 |
| 45 | TIS21 /BTG2/PC3 as a link between ageing and cancer: cell cycle regulator and endogenous cell death molecule. Journal of Cancer Research and Clinical Oncology, 2006, 132, 417-426. | 2.5 | 95 |
| 46 | Direct activation of TGF-β1 transcription by androgen and androgen receptor complex in Huh7 human hepatoma cells and its tumor in nude mice. Journal of Cellular Biochemistry, 2006, 97, 393-411. | 2.6 | 40 |
| 47 | Formation of elongated giant mitochondria in DFO-induced cellular senescence: Involvement of enhanced fusion process through modulation of Fis1. Journal of Cellular Physiology, 2006, 209, 468-480. | 4.1 | 234 |
| 48 | Phosphorylation of Serine 147 of tis21/BTG2/pc3 by p-Erk1/2 Induces Pin-1 Binding in Cytoplasm and Cell Death. Journal of Biological Chemistry, 2005, 280, 21256-21263. | 3.4 | 46 |
| 49 | Nuclear Accumulation of Globular Actin as a Cellular Senescence Marker. Cancer Research, 2004, 64, 572-580. | 0.9 | 65 |
| 50 | Compartmental inhibition of hepatic glutathione reductase activities by 1,3-bis(2-chloroethyl)-N-nitrosourea (BCNU) in Sprague–Dawley and Fischer-344 rats. Toxicology Letters, 2004, 147, 219-228. | 0.8 | 13 |
| 51 | TIS21/BTG2/PC3 is expressed through PKC-Ĩ´ pathway and inhibits binding of cyclin B1-Cdc2 and its activity, independent of p53 expression. Experimental Cell Research, 2004, 299, 159-170. | 2.6 | 53 |
| 52 | Erratum to "Spectrum of molecular changes during hepatocarcinogenesis induced by DEN and other chemicals in Fischer 344 male rats―[Mechanisms of Ageing and Development 123 (2002) 1665–1680]. Mechanisms of Ageing and Development, 2003, 124, 697-708. | 4.6 | 22 |
| 53 | Constitutive Induction of p-Erk1/2 Accompanied by Reduced Activities of Protein Phosphatases 1 and 2A and MKP3 Due to Reactive Oxygen Species during Cellular Senescence. Journal of Biological Chemistry, 2003, 278, 37497-37510. | 3.4 | 113 |
| 54 | Spectrum of molecular changes during hepatocarcinogenesis induced by DEN and other chemicals in Fischer 344 male rats. Mechanisms of Ageing and Development, 2002, 123, 1665-1680. | 4.6 | 36 |

IN KYOUNG LIM

| # | Article | IF | CITATIONS |
|----|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 55 | Marked inhibition of testosterone biosynthesis by the hepatotoxin nodularin due to apoptosis of Leydig cells. Molecular Carcinogenesis, 2002, 34, 151-163. | 2.7 | 16 |
| 56 | Induction of thermal and chemical stability of O6-methylguanine-DNA methyltransferase by Ca2+. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2002, 1599, 36-44. | 2.3 | 1 |
| 57 | Sequential changes in hepatocarcinogenesis induced by diethylnitrosamine plus thioacetamide in Fischer 344 rats: Induction of gankyrin expression in liver fibrosis, pRB degradation in cirrhosis, and methylation ofp16INK4A exon 1 in hepatocellular carcinoma. Molecular Carcinogenesis, 2001, 30, 138-150. | 2.7 | 50 |
| 58 | Translocational Inefficiency of Intracellular Proteins in Senescence of Human Diploid Fibroblasts. Annals of the New York Academy of Sciences, 2001, 928, 176-181. | 3.8 | 12 |
| 59 | Cytoplasmic retention of p-Erk1/2 and nuclear accumulation of actin proteins during cellular senescence in human diploid fibroblasts. Mechanisms of Ageing and Development, 2000, 119, 113-130. | 4.6 | 58 |
| 60 | Effect of nodularin on the expression of glutathione S-transferase placental form and proliferating cell nuclear antigen in N-nitrosodiethylamine initiated hepatocarcinogenesis in the male Fischer 344 rat. Carcinogenesis, 1999, 20, 1541-1548. | 2.8 | 24 |
| 61 | Differential expression of O   6 -methylguanine-DNA methyltransferase during diethylnitrosamine-induced carcinogenesis and liver regeneration in Sprague-Dawley male rats. Journal of Cancer Research and Clinical Oncology, 1999, 125, 493-499. | 2.5 | 13 |
| 62 | Regulation of selection of liver nodules initiated withN-nitrosodiethylamine and promoted with nodularin injections in Fischer 344 male rats by reciprocal expression of transforming growth factor-?1 and its receptors. Molecular Carcinogenesis, 1999, 26, 83-92. | 2.7 | 15 |
| 63 | Expression of rat BTG3 gene, Rbtg3, is regulated by redox changes. Gene, 1999, 240, 165-173. | 2.2 | 12 |
| 64 | Identification of highly methylated arginine residues in an endogenous 20-kDa polypeptide in cancer cells. Life Sciences, 1999, 65, 737-745. | 4.3 | 9 |
| 65 | ICE-like protease (caspase) is involved in transforming growth factor ?1-mediated apoptosis in FaO rat hepatoma cell line. Hepatology, 1998, 27, 415-421. | 7.3 | 48 |
| 66 | Induction of growth inhibition of 293 cells by downregulation of the cyclin E and cyclin-dependent kinase 4 proteins due to overexpression of TIS21. Molecular Carcinogenesis, 1998, 23, 25-35. | 2.7 | 67 |
| 67 | Biological methylation of myelin basic protein: Enzymology and biological significance. International Journal of Biochemistry and Cell Biology, 1997, 29, 743-751. | 2.8 | 73 |
| 68 | Protection of 5α-dihydrotestosterone against TGF-β-induced apoptosis in FaO cells and induction of mitosis in HepG2 cells. , 1997, 72, 351-355. | | 14 |
| 69 | Two synaptotagmin genes, Syt 1 and Syt 4, are differentially regulated in adult brain and during postnatal development following kainic acid-induced seizures. Molecular Brain Research, 1996, 40, 229-239. | 2.3 | 53 |
| 70 | Pronase-Based Assay Method forO6-Methylguanine–DNA Methyltransferase. Analytical Biochemistry, 1996, 236, 284-289. | 2.4 | 8 |
| 71 | Experimental Oncology Differential expression of TIS21 and TIS1 genes in the various organs of Balb/c mice, thymic carcinoma tissues and human cancer cell lines. Journal of Cancer Research and Clinical Oncology, 1995, 121, 279-284. | 2.5 | 37 |
| 72 | Increased drug resistance following retroviral gene transfer of a chimeric P-enolipyruvate carboxykinase (GTIP)-bacterial O6 alkylguanine-DNA alkyltransferase gene into NRK cells. Carcinogenesis, 1990, 11, 737-743. | 2.8 | 7 |