

Ralf Bender

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3555406/publications.pdf>

Version: 2024-02-01

40
papers

5,059
citations

394286

19
h-index

302012

39
g-index

43
all docs

43
docs citations

43
times ranked

9186
citing authors

#	ARTICLE	IF	CITATIONS
1	Performing Meta-analyses with Very Few Studies. <i>Methods in Molecular Biology</i> , 2022, 2345, 91-102.	0.4	8
2	GRADE Guidelines: 29. Rating the certainty in time-to-event outcomesâ€”Study limitations due to censoring of participants with missing data in intervention studies. <i>Journal of Clinical Epidemiology</i> , 2021, 129, 126-137.	2.4	4
3	The NNTnet metric is not new, not easy to use, and not routinely applied in medical research. <i>Journal of Clinical Epidemiology</i> , 2021, 129, 198.	2.4	2
4	On weakly informative prior distributions for the heterogeneity parameter in Bayesian randomâ€”effects metaâ€”analysis. <i>Research Synthesis Methods</i> , 2021, 12, 448-474.	4.2	55
5	A simulation study to compare different estimation approaches for network meta-analysis and corresponding methods to evaluate the consistency assumption. <i>BMC Medical Research Methodology</i> , 2020, 20, 36.	1.4	6
6	Methods to calculate uncertainty in the estimated overall effect size from a randomâ€”effects metaâ€”analysis. <i>Research Synthesis Methods</i> , 2019, 10, 23-43.	4.2	123
7	Limitations of the incidence density ratio as approximation of the hazard ratio. <i>Trials</i> , 2019, 20, 485.	0.7	9
8	The number needed to treat in pairwise and network meta-analysis and its graphical representation. <i>Journal of Clinical Epidemiology</i> , 2019, 111, 11-22.	2.4	32
9	Methods for evidence synthesis in the case of very few studies. <i>Research Synthesis Methods</i> , 2018, 9, 382-392.	4.2	132
10	Contribution to the discussion of â€œWhen should metaâ€”analysis avoid making hidden normality assumptions?â€” <i>Biometrical Journal</i> , 2018, 60, 1077-1078.	0.6	0
11	Adverse event development in clinical oncology trials. <i>Lancet Oncology</i> , The, 2016, 17, e263-e264.	5.1	3
12	Biometrical issues in the analysis of adverse events within the benefit assessment of drugs. <i>Pharmaceutical Statistics</i> , 2016, 15, 292-296.	0.7	22
13	Methods to estimate the betweenâ€”study variance and its uncertainty in metaâ€”analysis. <i>Research Synthesis Methods</i> , 2016, 7, 55-79.	4.2	891
14	A note on calculating asymptotic confidence intervals for the adjusted risk difference and number needed to treat in the Cox regression model. <i>Statistics in Medicine</i> , 2014, 33, 798-810.	0.8	6
15	Estimation of numbers needed to treat should be based on absolute risks. <i>Journal of Clinical Epidemiology</i> , 2014, 67, 238-239.	2.4	2
16	Absolute risks rather than incidence rates should be used to estimate the number needed to treat from time-to-event data. <i>Journal of Clinical Epidemiology</i> , 2013, 66, 1038-1044.	2.4	11
17	Unsolved issues of mixed treatment comparison metaâ€”analysis: network size and inconsistency. <i>Research Synthesis Methods</i> , 2012, 3, 300-311.	4.2	31
18	The assessment of heterogeneity is mandatory in clinical trials and systematic reviews. <i>Journal of Clinical Epidemiology</i> , 2011, 64, 452-452.	2.4	0

#	ARTICLE	IF	CITATIONS
19	Multiplicity issues in clinical trials. <i>Biometrical Journal</i> , 2011, 53, 873-874.	0.6	1
20	Methods to calculate relative risks, risk differences, and numbers needed to treat from logistic regression. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 7-8.	2.4	14
21	Common problems related to the use of number needed to treat. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 820-825.	2.4	69
22	Logistic regression was preferred to estimate risk differences and numbers needed to be exposed adjusted for covariates. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 1223-1231.	2.4	15
23	No inconsistent trial assessments by NICE and IQWiG: different assessment goals may lead to different assessment results regarding subgroup analyses. <i>Journal of Clinical Epidemiology</i> , 2010, 63, 1305-1307.	2.4	18
24	Estimating adjusted NNTs in randomised controlled trials with binary outcomes: A simulation study. <i>Contemporary Clinical Trials</i> , 2010, 31, 498-505.	0.8	14
25	Using and Interpreting Adjusted NNT Measures in Biomedical Research. <i>Open Dentistry Journal</i> , 2010, 4, 72-76.	0.2	2
26	Using and Interpreting Adjusted NNT Measures in Biomedical Research-!2009-09-25~!2009-10-14~!2010-07-16~!. <i>Open Dentistry Journal</i> , 2010, 4, 72-76.	0.2	4
27	Attention should be given to multiplicity issues in systematic reviews. <i>Journal of Clinical Epidemiology</i> , 2008, 61, 857-865.	2.4	117
28	Estimating adjusted NNT measures in logistic regression analysis. <i>Statistics in Medicine</i> , 2007, 26, 5586-5595.	0.8	57
29	Effect of Berkson measurement error on parameter estimates in Cox regression models. <i>Lifetime Data Analysis</i> , 2007, 13, 261-272.	0.4	20
30	Causes of death in obesity: Relevant increase in cardiovascular but not in all-cancer mortality. <i>Journal of Clinical Epidemiology</i> , 2006, 59, 1064-1071.	2.4	41
31	Generating survival times to simulate Cox proportional hazards models by Ralf Bender, Thomas Augustin and Maria Blettner, <i>Statistics in Medicine</i> 2005;24:1713-1723. <i>Statistics in Medicine</i> , 2006, 25, 1978-1979.	0.8	15
32	Tutorial: Using Confidence Curves in Medical Research. <i>Biometrical Journal</i> , 2005, 47, 237-247.	0.6	16
33	Generating survival times to simulate Cox proportional hazards models. <i>Statistics in Medicine</i> , 2005, 24, 1713-1723.	0.8	573
34	Confidence intervals for adjusted NNEs A simulation study. <i>Journal of Clinical Epidemiology</i> , 2003, 56, 205-206.	2.4	7
35	Breastfeeding promotion in non-UNICEF-certified hospitals and long-term breastfeeding success in Germany. <i>Acta Paediatrica, International Journal of Paediatrics</i> , 2003, 92, 653-658.	0.7	31
36	Body Weight, Blood Pressure, and Mortality in a Cohort of Obese Patients. <i>American Journal of Epidemiology</i> , 2002, 156, 239-245.	1.6	29

#	ARTICLE	IF	CITATIONS
37	Calculating the "number needed to be exposed" with adjustment for confounding variables in epidemiological studies. <i>Journal of Clinical Epidemiology</i> , 2002, 55, 525-530.	2.4	104
38	Adjusting for multiple testing "when and how?". <i>Journal of Clinical Epidemiology</i> , 2001, 54, 343-349.	2.4	2,133
39	Calculating Confidence Intervals for the Number Needed to Treat. <i>Contemporary Clinical Trials</i> , 2001, 22, 102-110.	2.0	187
40	Effect of Age on Excess Mortality in Obesity. <i>JAMA - Journal of the American Medical Association</i> , 1999, 281, 1498.	3.8	246