
## Jordi Segura

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3575189/publications.pdf Version: 2024-02-01



LODDI SECURA

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 1  | On the road of dried blood spot sampling for antidoping tests: Detection of GHRPâ€2 abuse. Drug<br>Testing and Analysis, 2021, 13, 510-522.   | 2.6 | 5         |
| 2  | If you play with fire, you may get burned. Drug Testing and Analysis, 2020, 12, 582-587.  | 2.6 | 3         |
| 3  | Automation of RNA-based biomarker extraction from dried blood spots for the detection of blood doping. Bioanalysis, 2020, 12, 729-736.  | 1.5 | 15        |
| 4  | Detection of Stimulated Erythropoiesis by the RNA-Based 5'-Aminolevulinate Synthase 2 Biomarker in<br>Dried Blood Spot Samples. Clinical Chemistry, 2019, 65, 1563-1571.  | 3.2 | 21        |
| 5  | Whole Blood Storage in CPDA1 Blood Bags Alters Erythrocyte Membrane Proteome. Oxidative<br>Medicine and Cellular Longevity, 2018, 2018, 1-12.   | 4.0 | 18        |
| 6  | Detection of erythropoiesisâ€ <b>s</b> timulating agents in a single dried blood spot. Drug Testing and Analysis,<br>2018, 10, 1496-1507.   | 2.6 | 25        |
| 7  | Evaluation of two glucuronides resistant to enzymatic hydrolysis as markers of testosterone oral administration. Journal of Steroid Biochemistry and Molecular Biology, 2017, 165, 212-218.   | 2.5 | 25        |
| 8  | Quantifying endogenous androgens, estrogens, pregnenolone and progesterone metabolites in human<br>urine by gas chromatography tandem mass spectrometry. Talanta, 2017, 169, 20-29.   | 5.5 | 40        |
| 9  | Determination of Recent Growth Hormone Abuse Using a Single Dried Blood Spot. Clinical Chemistry, 2016, 62, 1353-1360.  | 3.2 | 22        |
| 10 | Targeting tryptophan and tyrosine metabolism by liquid chromatography tandem mass spectrometry.<br>Journal of Chromatography A, 2016, 1434, 91-101.   | 3.7 | 72        |
| 11 | Bioanalytical techniques in discrimination between therapeutic and abusive use of drugs in sport.<br>Bioanalysis, 2016, 8, 965-980.   | 1.5 | 11        |
| 12 | Formation of Δ1 and Δ6 testosterone metabolites by human hepatocytes. Steroids, 2015, 95, 66-72.  | 1.8 | 7         |
| 13 | Detection and differentiation of 22kDa and 20kDa Growth Hormone proteoforms in human plasma by<br>LC-MS/MS. Biochimica Et Biophysica Acta - Proteins and Proteomics, 2015, 1854, 284-290.   | 2.3 | 9         |
| 14 | Urinary cysteinyl progestogens: Occurrence and origin. Journal of Steroid Biochemistry and<br>Molecular Biology, 2015, 152, 53-61.  | 2.5 | 10        |
| 15 | Screening for anabolic steroids in sports: Analytical strategy based on the detection of phase I and phase II intact urinary metabolites by liquid chromatography tandem mass spectrometry. Journal of Chromatography A, 2015, 1389, 65-75. | 3.7 | 37        |
| 16 | Ultra high performance liquid chromatography tandem mass spectrometric detection of<br>glucuronides resistant to enzymatic hydrolysis: Implications to doping control analysis. Analytica<br>Chimica Acta, 2015, 895, 35-44.                | 5.4 | 17        |
| 17 | Evaluation of the reporting level to detect triamcinolone acetonide misuse in sports. Journal of<br>Steroid Biochemistry and Molecular Biology, 2015, 145, 94-102.  | 2.5 | 18        |
| 18 | Investigation of endogenous corticosteroids profiles in human urine based on liquid<br>chromatography tandem mass spectrometry. Analytica Chimica Acta, 2014, 812, 92-104.  | 5.4 | 60        |

Jordi Segura

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Analysis of urinary human growth hormone (hGH) using hydrogel nanoparticles and isoform<br>differential immunoassays after short recombinant hGH treatment: Preliminary results. Journal of<br>Pharmaceutical and Biomedical Analysis, 2013, 85, 194-197. | 2.8 | 14        |
| 20 | Determination of five di-(2-ethylhexyl)phthalate metabolites in urine by UPLC–MS/MS, markers of<br>blood transfusion misuse in sports. Journal of Chromatography B: Analytical Technologies in the<br>Biomedical and Life Sciences, 2012, 908, 113-121.   | 2.3 | 36        |
| 21 | Alterations of the erythrocyte membrane proteome and cytoskeleton network during storage – a possible tool to identify autologous blood transfusion. Drug Testing and Analysis, 2012, 4, 882-890.   | 2.6 | 17        |
| 22 | Plasticizers excreted in urine: indication of autologous blood transfusion in sports. Transfusion, 2012, 52, 647-657.   | 1.6 | 35        |
| 23 | Progress in the Removal of Di-[2-Ethylhexyl]-Phthalate as Plasticizer in Blood Bags. Transfusion<br>Medicine Reviews, 2012, 26, 27-37.  | 2.0 | 78        |
| 24 | Growth hormone secretagogues: out of competition. Analytical and Bioanalytical Chemistry, 2012, 402, 1101-1108.   | 3.7 | 18        |
| 25 | Current strategic approaches for the detection of blood doping practices. Forensic Science International, 2011, 213, 42-48.   | 2.2 | 9         |
| 26 | Distinction Between Endogenous and Exogenous Erythropoietin: Marker Methods. Growth Hormone, 2011, , 151-161.   | 0.2 | 3         |
| 27 | Effect of physical fitness and endurance exercise on indirect biomarkers of growth hormone and<br>insulin misuse: Immunoassay-based measurement in urine samples. Journal of Pharmaceutical and<br>Biomedical Analysis, 2010, 53, 1003-1010.              | 2.8 | 12        |
| 28 | Urinary diâ€(2â€ethylhexyl)phthalate metabolites in athletes as screening measure for illicit blood<br>doping: a comparison study with patients receiving blood transfusion. Transfusion, 2010, 50, 145-149.  | 1.6 | 39        |
| 29 | Research Spotlight: Bioanalysis and Analytical Services Research Group at The Municipal Institute for<br>Medical Research IMIM-Hospital del Mar, Spain. Bioanalysis, 2009, 1, 1403-1409.  | 1.5 | 0         |
| 30 | Evaluation of different scan methods for the urinary detection of corticosteroid metabolites by<br>liquid chromatography tandem mass spectrometry. Journal of Mass Spectrometry, 2009, 44, 929-944.   | 1.6 | 46        |
| 31 | Evaluation of immunoassays for the measurement of insulin and C-peptide as indirect biomarkers of insulin misuse in sport: Values in selected population of athletes. Journal of Pharmaceutical and Biomedical Analysis, 2009, 49, 793-799.               | 2.8 | 14        |
| 32 | Is antiâ€doping analysis so far from clinical, legal or forensic targets?: The added value of close<br>relationships between related disciplines. Drug Testing and Analysis, 2009, 1, 479-484.  | 2.6 | 16        |
| 33 | Generation of 5 and 17 kDa human growth hormone fragments through limited proteolysis. Growth<br>Factors, 2009, 27, 255-264.  | 1.7 | 9         |
| 34 | Growth Hormone in Sport: Beyond Beijing 2008. Therapeutic Drug Monitoring, 2009, 31, 3-13.  | 2.0 | 23        |
| 35 | Immunoassays for the measurement of IGF-II, IGFBP-2 and -3, and ICTP as indirect biomarkers of recombinant human growth hormone misuse in sport. Journal of Pharmaceutical and Biomedical Analysis, 2008, 48, 844-852.                                    | 2.8 | 15        |
| 36 | Detection of the administration of 17 <i>β</i> â€nortestosterone in boars by gas chromatography/mass spectrometry. Rapid Communications in Mass Spectrometry, 2008, 22, 1863-1870.  | 1.5 | 16        |

Jordi Segura

| #  | Article  | IF  | CITATIONS |
|----|--|-----|-----------|
| 37 | Urinary metabolic profile of 19â€norsteroids in humans: glucuronide and sulphate conjugates after<br>oral administration of 19â€norâ€4â€androstenediol. Rapid Communications in Mass Spectrometry, 2008, 22,<br>3035-3042.                                   | 1.5 | 29        |
| 38 | Clarification on the detection of epoetin delta and epoetin omega using isoelectric focusing.<br>American Journal of Hematology, 2008, 83, 754-754.  | 4.1 | 8         |
| 39 | Structural analysis of the glycosylation of gene-activated erythropoietin (epoetin delta, Dynepo).<br>Analytical Biochemistry, 2008, 383, 243-254.   | 2.4 | 78        |
| 40 | Gas chromatography–mass spectrometry method for the analysis of 19-nor-4-androstenediol and<br>metabolites in human plasma: Application to pharmacokinetic studies after oral administration of a<br>prohormone supplement. Steroids, 2008, 73, 751-759.     | 1.8 | 10        |
| 41 | Characterisation of the 5 kDa growth hormone isoform. Growth Factors, 2008, 26, 152-162.   | 1.7 | 13        |
| 42 | High-Throughput and Sensitive Screening by Ultra-Performance Liquid Chromatography Tandem Mass<br>Spectrometry of Diuretics and other Doping Agents. European Journal of Mass Spectrometry, 2008, 14,<br>191-200.  | 1.0 | 63        |
| 43 | Recombinant erythropoietin found in seized blood bags from sportsmen. Haematologica, 2008, 93, 313-314.  | 3.5 | 28        |
| 44 | Intermittent hypoxia exposure in a hypobaric chamber and erythropoietin abuse interpretation.<br>Journal of Sports Sciences, 2007, 25, 1241-1250.  | 2.0 | 11        |
| 45 | Quantitation of 17β-nandrolone metabolites in boar and horse urine by gas chromatography–mass<br>spectrometry. Analytica Chimica Acta, 2007, 586, 184-195.   | 5.4 | 32        |
| 46 | Evaluation of protein <b><i>N</i></b> â€glycosylation in 2â€DE: Erythropoietin as a study case. Proteomics, 2007, 7, 4278-4291.  | 2.2 | 49        |
| 47 | Procedures for monitoring recombinant erythropoietin and analogues in doping control. Analytical and Bioanalytical Chemistry, 2007, 388, 1521-1529.  | 3.7 | 44        |
| 48 | Assessing the instability of the isoelectric focusing patterns of erythropoietin in urine.<br>Electrophoresis, 2006, 27, 4387-4395.  | 2.4 | 30        |
| 49 | Evaluation of immunoassays for the measurement of insulin-like growth factor-I and procollagen type<br>III peptide, indirect biomarkers of recombinant human growth hormone misuse in sport. Clinical<br>Chemistry and Laboratory Medicine, 2005, 43, 75-85. | 2.3 | 30        |
| 50 | Evaluation of immunoassays for the measurement of erythropoietin (EPO) as an indirect biomarker of<br>recombinant human EPO misuse in sport. Journal of Pharmaceutical and Biomedical Analysis, 2004, 35,<br>1169-1177.                                      | 2.8 | 26        |
| 51 | Plasma and urinary markers of oral testosterone undecanoate misuse. Steroids, 2002, 67, 39-50.   | 1.8 | 47        |
| 52 | Usefulness of Saliva for Measurement of 3,4-Methylenedioxymethamphetamine and Its Metabolites:<br>Correlation with Plasma Drug Concentrations and Effect of Salivary pH. Clinical Chemistry, 2001, 47,<br>1788-1795.   | 3.2 | 120       |
| 53 | Oral Testosterone Administration Detected by Testosterone Glucuronidation Measured in Blood<br>Spots Dried on Filter Paper. Clinical Chemistry, 2000, 46, 515-522.   | 3.2 | 37        |
| 54 | Immunological screening of drugs of abuse and gas chromatographic–mass spectrometric confirmation of opiates and cocaine in hair. Biomedical Applications, 1999, 724, 9-21.  | 1.7 | 43        |

| #  | Article   | IF  | CITATIONS |
|----|---|-----|-----------|
| 55 | Recent progress in the detection of the administration of natural hormones: Special focus on<br>Testosterone. Toxin Reviews, 1999, 18, 125-144. | 1.5 | 6         |
| 56 | Detection of testosterone esters in human plasma. Journal of Mass Spectrometry, 1995, 30, 1393-1404.  | 1.6 | 35        |