
Susan M O'brien

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3583897/publications.pdf Version: 2024-02-01



| # | Article | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Targeting BTK with Ibrutinib in Relapsed Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2013, 369, 32-42. | 13.9 | 2,019 |
| 2 | Idelalisib and Rituximab in Relapsed Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2014, 370, 997-1007. | 13.9 | 1,535 |
| 3 | Ibrutinib versus Ofatumumab in Previously Treated Chronic Lymphoid Leukemia. New England Journal of Medicine, 2014, 371, 213-223. | 13.9 | 1,427 |
| 4 | iwCLL guidelines for diagnosis, indications for treatment, response assessment, and supportive management of CLL. Blood, 2018, 131, 2745-2760. | 0.6 | 1,069 |
| 5 | Inotuzumab Ozogamicin versus Standard Therapy for Acute Lymphoblastic Leukemia. New England Journal of Medicine, 2016, 375, 740-753. | 13.9 | 1,047 |
| 6 | Safety and activity of blinatumomab for adult patients with relapsed or refractory B-precursor acute lymphoblastic leukaemia: a multicentre, single-arm, phase 2 study. Lancet Oncology, The, 2015, 16, 57-66. | 5.1 | 1,031 |
| 7 | Early Results of a Chemoimmunotherapy Regimen of Fludarabine, Cyclophosphamide, and Rituximab As Initial Therapy for Chronic Lymphocytic Leukemia. Journal of Clinical Oncology, 2005, 23, 4079-4088. | 0.8 | 899 |
| 8 | Acalabrutinib (ACP-196) in Relapsed Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2016, 374, 323-332. | 13.9 | 785 |
| 9 | Three-year follow-up of treatment-naÃ ⁻ ve and previously treated patients with CLL and SLL receiving single-agent ibrutinib. Blood, 2015, 125, 2497-2506. | 0.6 | 618 |
| 10 | The Bruton tyrosine kinase inhibitor PCI-32765 thwarts chronic lymphocytic leukemia cell survival and tissue homing in vitro and in vivo. Blood, 2012, 119, 1182-1189. | 0.6 | 564 |
| 11 | lbrutinib–Rituximab or Chemoimmunotherapy for Chronic Lymphocytic Leukemia. New England Journal of Medicine, 2019, 381, 432-443. | 13.9 | 545 |
| 12 | Fludarabine, cyclophosphamide, and rituximab treatment achieves long-term disease-free survival in IGHV-mutated chronic lymphocytic leukemia. Blood, 2016, 127, 303-309. | 0.6 | 441 |
| 13 | lbrutinib as initial therapy for elderly patients with chronic lymphocytic leukaemia or small lymphocytic lymphoma: an open-label, multicentre, phase 1b/2 trial. Lancet Oncology, The, 2014, 15, 48-58. | 5.1 | 438 |
| 14 | Inotuzumab ozogamicin, an anti-CD22–calecheamicin conjugate, for refractory and relapsed acute lymphocytic leukaemia: a phase 2 study. Lancet Oncology, The, 2012, 13, 403-411. | 5.1 | 401 |
| 15 | Chronic lymphocytic leukaemia. Nature Reviews Disease Primers, 2017, 3, 16096. | 18.1 | 363 |
| 16 | Chemoimmunotherapy With a Modified Hyper-CVAD and Rituximab Regimen Improves Outcome in De Novo Philadelphia Chromosome–Negative Precursor B-Lineage Acute Lymphoblastic Leukemia. Journal of Clinical Oncology, 2010, 28, 3880-3889. | 0.8 | 361 |
| 17 | Single-agent ibrutinib in treatment-naÃ ⁻ ve and relapsed/refractory chronic lymphocytic leukemia: a 5-year experience. Blood, 2018, 131, 1910-1919. | 0.6 | 339 |
| 18 | Final analysis from RESONATE: Up to six years of followâ€up on ibrutinib in patients with previously treated chronic lymphocytic leukemia or small lymphocytic lymphoma. American Journal of Hematology, 2019, 94, 1353-1363. | 2.0 | 305 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Outcomes of patients with chronic lymphocytic leukemia after discontinuing ibrutinib. Blood, 2015, 125, 2062-2067. | 0.6 | 303 |
| 20 | Ph-like acute lymphoblastic leukemia: a high-risk subtype in adults. Blood, 2017, 129, 572-581. | 0.6 | 285 |
| 21 | Management of adverse events associated with idelalisib treatment: expert panel opinion. Leukemia and Lymphoma, 2015, 56, 2779-2786. | 0.6 | 268 |
| 22 | Acalabrutinib Versus Ibrutinib in Previously Treated Chronic Lymphocytic Leukemia: Results of the First Randomized Phase III Trial. Journal of Clinical Oncology, 2021, 39, 3441-3452. | 0.8 | 266 |
| 23 | Results of inotuzumab ozogamicin, a CD22 monoclonal antibody, in refractory and relapsed acute lymphocytic leukemia. Cancer, 2013, 119, 2728-2736. | 2.0 | 265 |
| 24 | Early T-cell precursor acute lymphoblastic leukemia/lymphoma (ETP-ALL/LBL) in adolescents and adults: a high-risk subtype. Blood, 2016, 127, 1863-1869. | 0.6 | 253 |
| 25 | Prolonged lymphocytosis during ibrutinib therapy is associated with distinct molecular characteristics and does not indicate a suboptimal response to therapy. Blood, 2014, 123, 1810-1817. | 0.6 | 246 |
| 26 | Combination of hyper-CVAD with ponatinib as first-line therapy for patients with Philadelphia chromosome-positive acute lymphoblastic leukaemia: a single-centre, phase 2 study. Lancet Oncology, The, 2015, 16, 1547-1555. | 5.1 | 245 |
| 27 | A phase 2 study of idelalisib plus rituximab in treatment-naÃ⁻ve older patients with chronic lymphocytic leukemia. Blood, 2015, 126, 2686-2694. | 0.6 | 224 |
| 28 | Complex karyotype is a stronger predictor than del(17p) for an inferior outcome in relapsed or refractory chronic lymphocytic leukemia patients treated with ibrutinibâ€based regimens. Cancer, 2015, 121, 3612-3621. | 2.0 | 220 |
| 29 | Long-term outcome of acute promyelocytic leukemia treated with all-trans-retinoic acid, arsenic trioxide, and gemtuzumab. Blood, 2017, 129, 1275-1283. | 0.6 | 214 |
| 30 | Inotuzumab ozogamicin versus standard of care in relapsed or refractory acute lymphoblastic leukemia: Final report and longâ€ŧerm survival followâ€up from the randomized, phase 3 INOâ€VATE study. Cancer, 2019, 125, 2474-2487. | 2.0 | 210 |
| 31 | Other Malignancies in Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. Journal of Clinical Oncology, 2009, 27, 904-910. | 0.8 | 203 |
| 32 | Characterization of atrial fibrillation adverse events reported in ibrutinib randomized controlled registration trials. Haematologica, 2017, 102, 1796-1805. | 1.7 | 200 |
| 33 | Duvelisib, a novel oral dual inhibitor of PI3K-δ,γ, is clinically active in advanced hematologic malignancies. Blood, 2018, 131, 877-887. | 0.6 | 199 |
| 34 | Inotuzumab ozogamicin in combination with low-intensity chemotherapy for older patients with Philadelphia chromosome-negative acute lymphoblastic leukaemia: a single-arm, phase 2 study. Lancet Oncology, The, 2018, 19, 240-248. | 5.1 | 192 |
| 35 | Novel Targeted Agents and the Need to Refine Clinical End Points in Chronic Lymphocytic Leukemia. Journal of Clinical Oncology, 2012, 30, 2820-2822. | 0.8 | 182 |
| 36 | Randomized trial of ibrutinib vs ibrutinib plus rituximab in patients with chronic lymphocytic leukemia. Blood, 2019, 133, 1011-1019. | 0.6 | 168 |

| # | Article | IF | CITATIONS |
|----|--|------------------|--------------|
| 37 | Hepatic adverse event profile of inotuzumab ozogamicin in adult patients with relapsed or refractory acute lymphoblastic leukaemia: results from the open-label, randomised, phase 3 INO-VATE study. Lancet Haematology,the, 2017, 4, e387-e398. | 2.2 | 158 |
| 38 | International reference analysis of outcomes in adults with B-precursor Ph-negative relapsed/refractory acute lymphoblastic leukemia. Haematologica, 2016, 101, 1524-1533. | 1.7 | 154 |
| 39 | Monoclonal antibodies in acute lymphoblastic leukemia. Blood, 2015, 125, 4010-4016. | 0.6 | 144 |
| 40 | Economic Burden of Chronic Lymphocytic Leukemia in the Era of Oral Targeted Therapies in the United States. Journal of Clinical Oncology, 2017, 35, 166-174. | 0.8 | 131 |
| 41 | Second cancers in patients with chronic lymphocytic leukemia who received frontline fludarabine, cyclophosphamide and rituximab therapy: distribution and clinical outcomes. Leukemia and Lymphoma, 2015, 56, 1643-1650. | 0.6 | 130 |
| 42 | Acalabrutinib monotherapy in patients with relapsed/refractory chronic lymphocytic leukemia: updated phase 2 results. Blood, 2020, 135, 1204-1213. | 0.6 | 130 |
| 43 | Impact of BCR-ABL transcript type on outcome in patients with chronic-phase CML treated with tyrosine kinase inhibitors. Blood, 2016, 127, 1269-1275. | 0.6 | 119 |
| 44 | Evolution of CLL treatment — from chemoimmunotherapy to targeted and individualized therapy. Nature Reviews Clinical Oncology, 2018, 15, 510-527. | 12.5 | 114 |
| 45 | Results of the hyperfractionated cyclophosphamide, vincristine, doxorubicin, and dexamethasone regimen in elderly patients with acute lymphocytic leukemia. Cancer, 2008, 113, 2097-2101. | 2.0 | 109 |
| 46 | Defining the course and prognosis of adults with acute lymphocytic leukemia in first salvage after induction failure or short first remission duration. Cancer, 2010, 116, 5568-5574. | 2.0 | 104 |
| 47 | Longâ€ŧerm outcomes for patients with chronic lymphocytic leukemia who discontinue ibrutinib. Cancer, 2017, 123, 2268-2273. | 2.0 | 103 |
| 48 | Minimal residual disease assessed by multiâ€parameter flow cytometry is highly prognostic in adult patients with acute lymphoblastic leukaemia. British Journal of Haematology, 2016, 172, 392-400. | 1.2 | 102 |
| 49 | Long-term outcomes for ibrutinib–rituximab and chemoimmunotherapy in CLL: updated results of the E1912 trial. Blood, 2022, 140, 112-120. | 0.6 | 93 |
| 50 | Augmented Berlinâ€Frankfurtâ€Münster therapy in adolescents and young adults (AYAs) with acute lymphoblastic leukemia (ALL). Cancer, 2014, 120, 3660-3668. | 2.0 | 91 |
| 51 | Philadelphia-Positive Acute Lymphoblastic Leukemia: Current Treatment Options. Current Oncology Reports, 2012, 14, 387-394. | 1.8 | 83 |
| 52 | Long-term results of first salvage treatment in CLL patients treated initially with FCR (fludarabine,) Tj ETQq0 0 0 r | gBT /Over 0.6 | oçk 10 Tf 50 |

| 53 | Relevance of the immunoglobulin VH somatic mutation status in patients with chronic lymphocytic leukemia treated with fludarabine, cyclophosphamide, and rituximab (FCR) or related chemoimmunotherapy regimens. Blood, 2009, 113, 3168-3171. | 0.6 | 82 |
|----|---|-----|----|
| 54 | Collection and transfusion of granulocyte concentrates from donors primed with granulocyte stimulating factor and response of myelosuppressed patients with established infection. Journal of Clinical Apheresis, 1995, 10, 188-193. | 0.7 | 75 |

| # | Article | IF | CITATIONS |
|----|---|-----|-----------|
| 55 | Hyper VAD plus nelarabine in newly diagnosed adult Tâ€cell acute lymphoblastic leukemia and Tâ€lymphoblastic lymphoma. American Journal of Hematology, 2018, 93, 91-99. | 2.0 | 74 |
| 56 | Differential impact of minimal residual disease negativity according to the salvage status in patients with relapsed/refractory <scp>B</scp> â€eell acute lymphoblastic leukemia. Cancer, 2017, 123, 294-302. | 2.0 | 70 |
| 57 | Initial treatment of CLL: integrating biology and functional status. Blood, 2015, 126, 463-470. | 0.6 | 69 |
| 58 | Outcome of patients with relapsed/refractory acute lymphoblastic leukemia after blinatumomab failure: No change in the level of CD19 expression. American Journal of Hematology, 2018, 93, 371-374. | 2.0 | 68 |
| 59 | Cladribine and low-dose cytarabine alternating with decitabine as front-line therapy for elderly patients with acute myeloid leukaemia: a phase 2 single-arm trial. Lancet Haematology,the, 2018, 5, e411-e421. | 2.2 | 66 |
| 60 | Targeted therapies for CLL: Practical issues with the changing treatment paradigm. Blood Reviews, 2016, 30, 233-244. | 2.8 | 63 |
| 61 | Inotuzumab ozogamicin in combination with lowâ€intensity chemotherapy (miniâ€HCVD) with or without blinatumomab versus standard intensive chemotherapy (HCVAD) as frontline therapy for older patients with Philadelphia chromosomeâ€negative acute lymphoblastic leukemia: A propensity score analvsis. Cancer. 2019. 125. 2579-2586. | 2.0 | 63 |
| 62 | Statins enhance efficacy of venetoclax in blood cancers. Science Translational Medicine, 2018, 10, . | 5.8 | 61 |
| 63 | Five-Year Experience with Single-Agent Ibrutinib in Patients with Previously Untreated and Relapsed/Refractory Chronic Lymphocytic Leukemia/Small Lymphocytic Leukemia. Blood, 2016, 128, 233-233. | 0.6 | 60 |
| 64 | ATM gene deletion in patients with adult acute lymphoblastic leukemia. , 2000, 88, 1057-1062. | | 54 |
| 65 | Chronic Myelogenous Leukemia, Version 1.2014. Journal of the National Comprehensive Cancer Network: JNCCN, 2013, 11, 1327-1340. | 2.3 | 52 |
| 66 | ALPINE: zanubrutinib versus ibrutinib in relapsed/refractory chronic lymphocytic leukemia/small lymphocytic lymphoma. Future Oncology, 2020, 16, 517-523. | 1.1 | 52 |
| 67 | Prognostic impact of pretreatment cytogenetics in adult <scp>P</scp> hiladelphia chromosome–negative acute lymphoblastic leukemia in the era of minimal residual disease. Cancer, 2017, 123, 459-467. | 2.0 | 49 |
| 68 | Acalabrutinib in treatment-naive chronic lymphocytic leukemia. Blood, 2021, 137, 3327-3338. | 0.6 | 47 |
| 69 | Intensive chemotherapy induction followed by interferon-alpha maintenance in patients with Philadelphia chromosome-positive chronic myelogenous leukemia. Cancer, 1991, 68, 1201-1207. | 2.0 | 45 |
| 70 | Duvelisib, an oral dual PI3Kâ€̂f,γ inhibitor, shows clinical and pharmacodynamic activity in chronic lymphocytic leukemia and small lymphocytic lymphoma in a phase 1 study. American Journal of Hematology, 2018, 93, 1318-1326. | 2.0 | 45 |
| 71 | Fludarabine and cytosine arabinoside in the treatment of refractory or relapsed acute lymphocytic leukemia. Cancer, 1993, 72, 2155-2160. | 2.0 | 43 |
| 72 | Efficacy and safety analysis by age cohort of inotuzumab ozogamicin in patients with relapsed or refractory acute lymphoblastic leukemia enrolled in INOâ€VATE. Cancer, 2018, 124, 1722-1732. | 2.0 | 43 |

| # | Article | IF | CITATIONS |
|----|--|-----|-----------|
| 73 | Hyper-CVAD regimen in combination with ofatumumab as frontline therapy for adults with Philadelphia chromosome-negative B-cell acute lymphoblastic leukaemia: a single-arm, phase 2 trial. Lancet Haematology,the, 2020, 7, e523-e533. | 2.2 | 43 |
| 74 | Liposomal Grb2 antisense oligodeoxynucleotide (BP1001) in patients with refractory or relapsed haematological malignancies: a single-centre, open-label, dose-escalation, phase 1/1b trial. Lancet Haematology,the, 2018, 5, e136-e146. | 2.2 | 42 |
| 75 | Ibrutinib restores immune cell numbers and function in first-line and relapsed/refractory chronic lymphocytic leukemia. Leukemia Research, 2020, 97, 106432. | 0.4 | 40 |
| 76 | Optimal Management of Adverse Events From Copanlisib in the Treatment of Patients With Non-Hodgkin Lymphomas. Clinical Lymphoma, Myeloma and Leukemia, 2019, 19, 135-141. | 0.2 | 37 |
| 77 | Prognostic factors for outcome in patients with refractory and relapsed acute lymphocytic leukemia treated with inotuzumab ozogamicin, a <scp>CD</scp> 22 monoclonal antibody. American Journal of Hematology, 2015, 90, 193-196. | 2.0 | 35 |
| 78 | The absolute percent deviation of <i><scp>IGHV</scp></i> mutation rather than a 98% cutâ€off predicts survival of chronic lymphocytic leukaemia patients treated with fludarabine, cyclophosphamide and rituximab. British Journal of Haematology, 2018, 180, 33-40. | 1.2 | 33 |
| 79 | <scp>S</scp> ignificance of recurrence of minimal residual disease detected by multiâ€parameter flow cytometry in patients with acute lymphoblastic leukemia in morphological remission. American Journal of Hematology, 2017, 92, 279-285. | 2.0 | 32 |
| 80 | Combination of topotecan with cytarabine or etoposide in patients with refractory or relapsed acute myeloid leukemia: results of a randomized phase I/II study. Investigational New Drugs, 1999, 17, 89-95. | 1.2 | 31 |
| 81 | Advances in the Genetics and Therapy of Acute Lymphoblastic Leukemia. American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting, 2016, 35, e314-e322. | 1.8 | 28 |
| 82 | Longâ€ŧerm efficacy of firstâ€line ibrutinib treatment for chronic lymphocytic leukaemia in patients with <i>TP53</i> aberrations: a pooled analysis from four clinical trials. British Journal of Haematology, 2022, 196, 947-953. | 1.2 | 28 |
| 83 | Bone marrow necrosis in acute leukemia: Clinical characteristic and outcome. American Journal of Hematology, 2015, 90, 769-773. | 2.0 | 27 |
| 84 | Outcomes with ibrutinib by line of therapy and postâ€ibrutinib discontinuation in patients with chronic lymphocytic leukemia: Phase 3 analysis. American Journal of Hematology, 2019, 94, 554-562. | 2.0 | 27 |
| 85 | Monitoring and Managing BTK Inhibitor Treatment-Related Adverse Events in Clinical Practice. Frontiers in Oncology, 2021, 11, 720704. | 1.3 | 27 |
| 86 | Philadelphia chromosomeâ€positive acute lymphoblastic leukemia at first relapse in the era of tyrosine kinase inhibitors. American Journal of Hematology, 2019, 94, 1388-1395. | 2.0 | 26 |
| 87 | Survival of Young Patients with Chronic Lymphocytic Leukemia Failing Fludarabine Therapy: A Basis for the Use of Myeloablative Therapies. Leukemia and Lymphoma, 1995, 18, 493-496. | 0.6 | 25 |
| 88 | The early achievement of measurable residual disease negativity in the treatment of adults with Philadelphiaâ€negative Bâ€cell acute lymphoblastic leukemia is a strong predictor for survival. American Journal of Hematology, 2020, 95, 144-150. | 2.0 | 25 |
| 89 | Pooled analysis of safety data from clinical trials evaluating acalabrutinib monotherapy in mature B-cell malignancies. Leukemia, 2021, 35, 3201-3211. | 3.3 | 25 |
| 90 | Clinical relevance of intracellular vascular endothelial growth factor levels in B-cell chronic lymphocytic leukemia. Blood, 2000, 96, 768-770. | 0.6 | 25 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Longâ€ŧerm followâ€up of salvage therapy using a combination of inotuzumab ozogamicin and mini–hyperâ€CVD with or without blinatumomab in relapsed/refractory Philadelphia chromosome–negative acute lymphoblastic leukemia. Cancer, 2021, 127, 2025-2038. | 2.0 | 24 |
| 92 | Outcomes of acute lymphoblastic leukemia with <i>KMT2A</i> (<i>MLL</i>) rearrangement: the MD Anderson experience. Blood Advances, 2021, 5, 5415-5419. | 2.5 | 24 |
| 93 | Updated Efficacy Including Genetic and Clinical Subgroup Analysis and Overall Safety in the Phase 3 RESONATETM Trial of Ibrutinib Versus Ofatumumab in Previously Treated Chronic Lymphocytic Leukemia/Small Lymphocytic Lymphoma. Blood, 2014, 124, 3331-3331. | 0.6 | 24 |
| 94 | Acalabrutinib and its use in treatment of chronic lymphocytic leukemia. Future Oncology, 2019, 15, 579-589. | 1.1 | 23 |
| 95 | Ongoing Results of a Phase 1B/2 Dose-Escalation and Cohort-Expansion Study of the Selective, Noncovalent, Reversible Bruton'S Tyrosine Kinase Inhibitor, Vecabrutinib, in B-Cell Malignancies. Blood, 2019, 134, 3041-3041. | 0.6 | 23 |
| 96 | Ibrutinib, fludarabine, cyclophosphamide, and obinutuzumab (iFCG) regimen for chronic lymphocytic leukemia (CLL) with mutated IGHV and without TP53 aberrations. Leukemia, 2021, 35, 3421-3429. | 3.3 | 22 |
| 97 | Novel Treatments for Chronic Lymphocytic Leukemia and Moving Forward. American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting, 2014, , e317-e325. | 1.8 | 17 |
| 98 | Molecular response with blinatumomab in relapsed/refractory B-cell precursor acute lymphoblastic leukemia. Blood Advances, 2019, 3, 3033-3037. | 2.5 | 16 |
| 99 | Measurable residual disease does not preclude prolonged progression-free survival in CLL treated with ibrutinib. Blood, 2021, 138, 2810-2827. | 0.6 | 16 |
| 100 | Novel agents in chronic lymphocytic leukemia. Hematology American Society of Hematology Education Program, 2016, 2016, 137-145. | 0.9 | 14 |
| 101 | Ibrutinib, Fludarabine, Cyclophosphamide, and Obinutuzumab (iFCG) for First-Line Treatment of IGHV-Mutated CLL and without Del(17p)/Mutated TP53. Blood, 2019, 134, 357-357. | 0.6 | 14 |
| 102 | Blastic Plasmacytoid Dendritic Cell Neoplasm (BPDCN) Commonly Presents in the Setting of Prior or Concomitant Hematologic Malignancies (PCHM): Patient Characteristics and Outcomes in the Rapidly Evolving Modern Targeted Therapy Era. Blood, 2019, 134, 2723-2723. | 0.6 | 14 |
| 103 | Approaches to Chronic Lymphocytic Leukemia Therapy in the Era of New Agents: The Conundrum of Many Options. American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting, 2018, 38, 580-591. | 1.8 | 13 |
| 104 | Salvage Chemotherapy with Inotuzumab Ozogamicin (INO) Combined with Mini-Hyper-CVD for Adult Patients with Relapsed/Refractory (R/R) Acute Lymphoblastic Leukemia (ALL). Blood, 2015, 126, 3721-3721. | 0.6 | 13 |
| 105 | Updated Results of a Phase II Study of Reduced-Intensity Chemotherapy with Mini-Hyper-CVD in Combination with Inotuzumab Ozogamicin, with or without Blinatumomab, in Older Adults with Newly Diagnosed Philadelphia Chromosome-Negative Acute Lymphoblastic Leukemia. Blood, 2019, 134, 823-823. | 0.6 | 12 |
| 106 | A Phase I Study of Fludarabine, Cytarabine, and Oxaliplatin Therapy in Patients With Relapsed or Refractory Acute Myeloid Leukemia. Clinical Lymphoma, Myeloma and Leukemia, 2014, 14, 395-400.e1. | 0.2 | 11 |
| 107 | Prognostic implications of cytogenetics in adults with acute lymphoblastic leukemia treated with inotuzumab ozogamicin. American Journal of Hematology, 2019, 94, 408-416. | 2.0 | 11 |
| 108 | Inotuzumab Ozogamicin in Combination with Low-Intensity Chemotherapy (mini-hyper-CVD) As Frontline Therapy for Older Patients with Acute Lymphoblastic Leukemia (ALL): Interim Result of a Phase II Clinical Trial. Blood, 2016, 128, 588-588. | 0.6 | 11 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 109 | Hyper VAD plus ofatumumab versus hyper VAD plus rituximab as frontline therapy in adults with Philadelphia chromosome–negative acute lymphoblastic leukemia: A propensity score analysis. Cancer, 2021, 127, 3381-3389. | 2.0 | 10 |
| 110 | B cell receptor inhibition as a target for CLL therapy. Best Practice and Research in Clinical Haematology, 2016, 29, 2-14. | 0.7 | 9 |
| 111 | The safety of Bruton's tyrosine kinase inhibitors for the treatment of chronic lymphocytic leukemia. Expert Opinion on Drug Safety, 2017, 16, 1079-1088. | 1.0 | 9 |
| 112 | A phase <scp>II</scp> trial of eltrombopag for patients with chronic lymphocytic leukaemia (<scp>CLL</scp>) and thrombocytopenia. British Journal of Haematology, 2019, 185, 606-608. | 1.2 | 8 |
| 113 | Mechanisms of ibrutinib resistance in chronic lymphocytic leukemia and alternative treatment strategies. Expert Review of Hematology, 2020, 13, 871-883. | 1.0 | 8 |
| 114 | Using ibrutinib in earlier lines of treatment results in better outcomes for patients with chronic lymphocytic leukemia/small lymphocytic lymphoma. Leukemia and Lymphoma, 2021, 62, 3278-3282. | 0.6 | 7 |
| 115 | Reassessing the role of chemoimmunotherapy in chronic lymphocytic leukemia. Expert Review of Hematology, 2020, 13, 31-38. | 1.0 | 5 |
| 116 | Phase II Study of Blinatumomab in Patients with B-Cell Acute Lymphoblastic Leukemia (B-ALL) with Positive Measurable Residual Disease (MRD). Blood, 2019, 134, 1299-1299. | 0.6 | 4 |
| 117 | Causes of Discontinuation and Long-Term Outcomes of Patients with CLL after Discontinuing Ibrutinib. Blood, 2016, 128, 4390-4390. | 0.6 | 4 |
| 118 | Increased incidence of Pegaspargase-induced hypertriglyceridemia and associated pancreatitis observed in the Hispanic adult patient population. Leukemia and Lymphoma, 2022, 63, 2992-2995. | 0.6 | 4 |
| 119 | Efficacy and Safety Outcomes in the Phase 3 INO-Vate Trial By Baseline CD22 Positivity Assessed By Local Laboratories. Blood, 2019, 134, 1344-1344. | 0.6 | 3 |
| 120 | A 20-Year Review of Imatinib in Chronic Phase Chronic Myeloid Leukemia Patients after Failure with Interferon Therapy. Blood, 2019, 134, 2927-2927. | 0.6 | 3 |
| 121 | Externally validated predictive clinical model for untreated del(17p13.1) chronic lymphocytic leukemia patients. American Journal of Hematology, 2015, 90, 967-969. | 2.0 | 2 |
| 122 | Characteristics and Clinical Outcomes of Patients with Acute Lymphoblastic Leukemia with KMT2A (MLL) Rearrangement. Blood, 2019, 134, 2582-2582. | 0.6 | 2 |
| 123 | First-line therapy for young patients with CLL. Hematology American Society of Hematology Education Program, 2016, 2016, 146-148. | 0.9 | 1 |
| 124 | MLL-rearranged mixed phenotype acute leukemia masquerading as B-cell ALL. Leukemia and Lymphoma, 2017, 58, 1498-1501. | 0.6 | 1 |
| 125 | ATM gene deletion in patients with adult acute lymphoblastic leukemia. , 2000, 88, 1057. | | 1 |
| 126 | Human-Leukocyte-Histocompatibility Antigens Predict Response to Rituximab and Donor Lymphocyte Infusion (DLI) After Non-Myeloablative Allogeneic Stem Transplantation (NST) for Chronic Lymphocytic Leukemia (CLL). Blood, 2010, 116, 2548-2548. | 0.6 | 0 |

| # | Article | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Comparison of Hyper-CVAD Plus Ofatumumab to Hyper-CVAD Plus Rituximab in Patients with Newly Diagnosed Philadelphia Chromosome-Negative CD20-Positive B-Cell Acute Lymphoblastic Leukemia: A Propensity Score Analysis. Blood, 2020, 136, 42-43. | 0.6 | 0 |
| 128 | Clinical implications of the 2018 iwCLL Guidelines update. Clinical Advances in Hematology and Oncology, 2018, 16 Suppl 15, 1-16. | 0.3 | 0 |
| 129 | Characterization of low-grade arthralgia, myalgia, and musculoskeletal pain with ibrutinib therapy: pooled analysis of clinical trials in patients with chronic lymphocytic leukemia and mantle cell lymphoma. Leukemia and Lymphoma, 2022, 63, 1580-1588. | 0.6 | 0 |