

Laura M Torres

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3598514/publications.pdf>

Version: 2024-02-01

29
papers

429
citations

687363

13
h-index

752698

20
g-index

30
all docs

30
docs citations

30
times ranked

489
citing authors

#	ARTICLE	IF	CITATIONS
1	Evaluation of the effects, on canopy arthropods, of two agricultural management systems to control pests in olive groves from north-east of Portugal. <i>Chemosphere</i> , 2007, 67, 131-139.	8.2	56
2	Influence of fruit traits on oviposition preference of the olive fly, <i>Bactrocera oleae</i> (Rossi) (Diptera: Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50). <i>Scientia Horticulturae</i> , 2012, 145, 127-135.	3.6	31
3	Insect-associated fungi from naturally mycosed vine mealybug <i>Planococcus ficus</i> (Signoret) (Hemiptera: Pseudococcidae). <i>Biocontrol Science and Technology</i> , 2018, 28, 122-141.	1.3	30
4	Entomopathogenic fungi in Portuguese vineyards soils: suggesting a <i>Galleria-Tenebrio-bait method</i> ™ as bait-insects <i>Galleria</i> and <i>Tenebrio</i> significantly underestimate the respective recoveries of <i>Metarhizium (robertsii)</i> and <i>Beauveria (bassiana)</i> . <i>MycoKeys</i> , 2018, 38, 1-23.	1.9	29
5	Effects of pollen, sugars and honeydew on lifespan and nutrient levels of <i>Episyrphus balteatus</i> . <i>BioControl</i> , 2015, 60, 47-57.	2.0	26
6	Egg parasitoids of the genus <i>Trichogramma</i> (Hymenoptera, Trichogrammatidae) in olive groves of the Mediterranean region. <i>Biological Control</i> , 2007, 40, 48-56.	3.0	24
7	Effect of floral resources on longevity and nutrient levels of <i>Episyrphus balteatus</i> (Diptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50).	3.0	21
8	Does habitat heterogeneity affect the diversity of epigeic arthropods in vineyards?. <i>Agricultural and Forest Entomology</i> , 2018, 20, 366-379.	1.3	20
9	The effect of sooty mold on fluorescence and gas exchange properties of olive tree. <i>Turkish Journal of Biology</i> , 2013, 37, 620-628.	0.8	17
10	Higher longevity and fecundity of <i>Chrysoperla carnea</i> , a predator of olive pests, on some native flowering Mediterranean plants. <i>Agronomy for Sustainable Development</i> , 2016, 36, 1.	5.3	15
11	Is a biofix necessary for predicting the flight phenology of <i>Lobesia botrana</i> in Douro Demarcated Region vineyards?. <i>Crop Protection</i> , 2018, 110, 57-64.	2.1	15
12	The use of the cumulative degree-days to predict olive fly, <i>Bactrocera oleae</i> (Rossi), activity in traditional olive groves from the northeast of Portugal. <i>Journal of Pest Science</i> , 2011, 84, 187-197.	3.7	14
13	Natural mortality of immature stages of <i>Bactrocera oleae</i> (Diptera: Tephritidae) in traditional olive groves from north-eastern Portugal. <i>Biocontrol Science and Technology</i> , 2012, 22, 837-854.	1.3	12
14	Soil Arthropods in the Douro Demarcated Region Vineyards: General Characteristics and Ecosystem Services Provided. <i>Sustainability</i> , 2021, 13, 7837.	3.2	12
15	A cohort-based modelling approach for managing olive moth <i>Prays oleae</i> (Bernard, 1788) populations in olive orchards. <i>Ecological Modelling</i> , 2015, 296, 46-56.	2.5	11
16	Effects of ten naturally occurring sugars on the reproductive success of the green lacewing, <i>Chrysoperla carnea</i> . <i>BioControl</i> , 2016, 61, 57-67.	2.0	9
17	Identification of predator-prey relationships between coccinellids and <i>Saissetia oleae</i> (Hemiptera: Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50). <i>Entomologia Experimentalis et Applicata</i> , 2018, 82, 101-108.	3.7	8
18	Soil Chemical Properties Barely Perturb the Abundance of Entomopathogenic <i>Fusarium oxysporum</i> : A Case Study Using a Generalized Linear Mixed Model for Microbial Pathogen Occurrence Count Data. <i>Pathogens</i> , 2018, 7, 89.	2.8	8

#	ARTICLE	IF	CITATIONS
19	Ants (Hymenoptera: Formicidae) and Spiders (Araneae) Co-occurring on the Ground of Vineyards from Douro Demarcated Region. <i>Sociobiology</i> , 2017, 64, 404.	0.5	7
20	Native Mediterranean plants as potential food sources for natural enemies of insect pests in olive groves. <i>Ecological Research</i> , 2017, 32, 459-459.	1.5	6
21	The functional agrobiodiversity in the Douro demarcated region viticulture: utopia or reality? Arthropods as a case-study – A review. <i>Ciencia E Tecnica Vitivinicola</i> , 2019, 34, 102-114.	0.9	6
22	Effect of Soil Chemical Properties on the Occurrence and Distribution of Entomopathogenic Fungi in Portuguese Grapevine Fields. <i>Pathogens</i> , 2021, 10, 137.	2.8	6
23	The use of trap captures to forecast infestation by the olive fly, <i>Bactrocera oleae</i> (Rossi) (Diptera: Tephritidae), in traditional olive groves in north-eastern Portugal. <i>International Journal of Pest Management</i> , 2013, 59, 279-286.	1.8	5
24	Entomopathogenic fungi in Portuguese vineyards soils: suggesting a “Galleria-Tenebrio-bait method” as bait-insects <i>Galleria</i> and <i>Tenebrio</i> significantly underestimate the respective recoveries of <i>Metarhizium</i> (robertsii) and <i>Beauveria</i> (bassiana). <i>Mycology</i> , 0, 38, 1-23.	1.9	4
25	Evaluating potential sugar food sources from the olive grove agroecosystems for <i>Prays oleae</i> parasitoid <i>Chelonus elaeaphilus</i> . <i>Biocontrol Science and Technology</i> , 2017, 27, 686-695.	1.3	3
26	Hymenoptera parasitoid complex of <i>Prays oleae</i> (Bernard) (Lepidoptera: Praydidae) in Portugal. <i>Turkish Journal of Zoology</i> , 2017, 41, 502-512.	0.9	3
27	Does natural vegetation from olive groves benefit the olive moth, <i>Prays oleae</i> ? <i>Journal of Applied Entomology</i> , 2021, 145, 406-416.	1.8	2
28	Parasitoids of <i>Lobesia botrana</i> (Lepidoptera: Tortricidae) in the Douro Demarcated Region vineyards and the prospects for enhancing conservation biological control. <i>Bulletin of Entomological Research</i> , 2022, , 1-10.	1.0	2
29	Confusão sexual contra a traça-da-uva, <i>Lobesia botrana</i> , na região Demarcada do Douro usando dois modelos de difusores de feromona: ISONET-LTT BIO [®] E ISONET-LTT [®] . <i>Ciencia E Tecnica Vitivinicola</i> , 2022, 37, 100-115.	0.9	1