
Tatiana V Loboda

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3610526/publications.pdf Version: 2024-02-01



Τλτιλνίλ VI οβορλ

#	Article	IF	CITATIONS
1	Cloudâ€toâ€Ground Lightning and Nearâ€6urface Fire Weather Control Wildfire Occurrence in Arctic Tundra. Geophysical Research Letters, 2022, 49, .	4.0	12
2	Consequences of a future increase in fire: The human health perspective. One Earth, 2021, 4, 487-488.	6.8	2
3	Spatio-temporal patterns of optimal Landsat data for burn severity index calculations: Implications for high northern latitudes wildfire research. Remote Sensing of Environment, 2021, 258, 112393.	11.0	13
4	Satellite Observations and Malaria: New Opportunities for Research and Applications. Trends in Parasitology, 2021, 37, 525-537.	3.3	34
5	Impacts of wildfire and landscape factors on organic soil properties in Arctic tussock tundra. Environmental Research Letters, 2021, 16, 085004.	5.2	7
6	Characterizing Small-Town Development Using Very High Resolution Imagery within Remote Rural Settings of Mozambique. Remote Sensing, 2021, 13, 3385.	4.0	2
7	Missing Burns in the High Northern Latitudes: The Case for Regionally Focused Burned Area Products. Remote Sensing, 2021, 13, 4145.	4.0	11
8	Spaceâ€Based Observations for Understanding Changes in the Arcticâ€Boreal Zone. Reviews of Geophysics, 2020, 58, e2019RG000652.	23.0	39
9	A systematic evaluation of influence of image selection process on remote sensing-based burn severity indices in North American boreal forest and tundra ecosystems. ISPRS Journal of Photogrammetry and Remote Sensing, 2020, 159, 63-77.	11.1	28
10	Malaria Exposure in Ann Township, Myanmar, as a Function of Land Cover and Land Use: Combining Satellite Earth Observations and Field Surveys. GeoHealth, 2020, 4, e2020GH000299.	4.0	5
11	Long-term trends in anthropogenic land use in Siberia and the Russian Far East: a case study synthesis from Landsat. Environmental Research Letters, 2020, 15, 105007.	5.2	6
12	Modeling cloud-to-ground lightning probability in Alaskan tundra through the integration of Weather Research and Forecast (WRF) model and machine learning method. Environmental Research Letters, 2020, 15, 115009.	5.2	5
13	Mapping fractional cover of major fuel type components across Alaskan tundra. Remote Sensing of Environment, 2019, 232, 111324.	11.0	12
14	Mapping remote rural settlements at 30â€⁻m spatial resolution using geospatial data-fusion. Remote Sensing of Environment, 2019, 233, 111386.	11.0	28
15	Oil palm plantations in Peninsular Malaysia: Determinants and constraints on expansion. PLoS ONE, 2019, 14, e0210628.	2.5	61
16	Strong cooling induced by stand-replacing fires through albedo in Siberian larch forests. Scientific Reports, 2018, 8, 4821.	3.3	23
17	Surface forcing of non-stand-replacing fires in Siberian larch forests. Environmental Research Letters, 2018, 13, 045008.	5.2	16
18	Examining aspiration's imprint on the landscape: Lessons from Mozambique's Limpopo National Park. Global Environmental Change, 2018, 51, 43-53.	7.8	9

TATIANA V LOBODA

#	Article	IF	CITATIONS
19	Spatial distribution of young forests and carbon fluxes within recent disturbances in Russia. Global Change Biology, 2017, 23, 138-153.	9.5	12
20	Multi-Decadal Surface Water Dynamics in North American Tundra. Remote Sensing, 2017, 9, 497.	4.0	41
21	Expansion of Industrial Plantations Continues to Threaten Malayan Tiger Habitat. Remote Sensing, 2017, 9, 747.	4.0	15
22	Static and dynamic controls on fire activity at moderate spatial and temporal scales in the Alaskan boreal forest. Ecosphere, 2016, 7, e01572.	2.2	16
23	A MODIS-based burned area assessment for Russian croplands: Mapping requirements and challenges. Remote Sensing of Environment, 2016, 184, 506-521.	11.0	95
24	Mapping stand age dynamics of the Siberian larch forests from recent Landsat observations. Remote Sensing of Environment, 2016, 187, 320-331.	11.0	17
25	Can carbon emissions from tropical deforestation drop by 50% in 5Âyears?. Global Change Biology, 2016, 22, 1336-1347.	9.5	109
26	Fire in arctic tundra of Alaska: past fire activity, future fire potential, and significance for land management and ecology. International Journal of Wildland Fire, 2015, 24, 1045.	2.4	53
27	Long-Term Record of Sampled Disturbances in Northern Eurasian Boreal Forest from Pre-2000 Landsat Data. Remote Sensing, 2014, 6, 6020-6038.	4.0	7
28	Adaptation strategies to climate change in the Arctic: a global patchwork of reactive community-scale initiatives. Environmental Research Letters, 2014, 9, 111006.	5.2	6
29	Remote sensing estimates of stand-replacement fires in Russia, 2002–2011. Environmental Research Letters, 2014, 9, 105007.	5.2	70
30	Santa Ana winds and predictors of wildfire progression in southern California. International Journal of Wildland Fire, 2014, 23, 1119.	2.4	22
31	Development of Methods for Detection and Monitoring of Fire Disturbance in the Alaskan Tundra Using a Two-Decade Long Record of Synthetic Aperture Radar Satellite Images. Remote Sensing, 2014, 6, 6347-6364.	4.0	19
32	Analysis of the Impacts of armed conflict on the Eastern Afromontane forest region on the South Sudan — Uganda border using multitemporal Landsat imagery. Remote Sensing of Environment, 2012, 118, 10-20.	11.0	80
33	Quantifying burned area for North American forests: Implications for direct reduction of carbon stocks. Journal of Geophysical Research, 2011, 116, .	3.3	39
34	Comparison and assessment of coarse resolution land cover maps for Northern Eurasia. Remote Sensing of Environment, 2011, 115, 3539-3553.	11.0	75
35	Mapping burned area in Alaska using MODIS data: a data limitations-driven modification to the regional burned area algorithm. International Journal of Wildland Fire, 2011, 20, 487.	2.4	35
36	Modeling fire danger in data-poor regions: a case study from the Russian Far East. International Journal of Wildland Fire, 2009, 18, 19.	2.4	17

TATIANA V LOBODA

#	Article	IF	CITATIONS
37	The spatial and temporal distribution of crop residue burning in the contiguous United States. Science of the Total Environment, 2009, 407, 5701-5712.	8.0	115
38	An active-fire based burned area mapping algorithm for the MODIS sensor. Remote Sensing of Environment, 2009, 113, 408-420.	11.0	533
39	ASSESSING THE RISK OF IGNITION IN THE RUSSIAN FAR EAST WITHIN A MODELING FRAMEWORK OF FIRE THREAT. , 2007, 17, 791-805.		29
40	Global distribution of agricultural fires in croplands from 3 years of Moderate Resolution Imaging Spectroradiometer (MODIS) data. Global Biogeochemical Cycles, 2006, 20, n/a-n/a.	4.9	201
41	AVHRR-based mapping of fires in Russia: New products for fire management and carbon cycle studies. Remote Sensing of Environment, 2004, 93, 546-564.	11.0	224