
Paul Knekt

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3663637/publications.pdf Version: 2024-02-01



DALL KNEKT

#	Article	IF	CITATIONS
1	Ego Impairment Index (Ellâ€2) as a predictor of outcome in short―and longâ€ŧerm psychotherapy during a 5â€year followâ€up. Journal of Clinical Psychology, 2022, 78, 1739-1751.	1.9	1
2	Circulating free testosterone and risk of aggressive prostate cancer: Prospective and Mendelian randomisation analyses in international consortia. International Journal of Cancer, 2022, 151, 1033-1046.	5.1	18
3	Predicting improvement of work ability in modalities of short―and longâ€ŧerm psychotherapy: The differential impact of reflective ability and other aspects of patient suitability. Journal of Clinical Psychology, 2021, 77, 1905-1920.	1.9	8
4	Associations of Total Legume, Pulse, and Soy Consumption with Incident Type 2 Diabetes: Federated Meta-Analysis of 27 Studies from Diverse World Regions. Journal of Nutrition, 2021, 151, 1231-1240.	2.9	28
5	Heterogeneity of Associations between Total and Types of Fish Intake and the Incidence of Type 2 Diabetes: Federated Meta-Analysis of 28 Prospective Studies Including 956,122 Participants. Nutrients, 2021, 13, 1223.	4.1	8
6	Incidence trends and risk factors of lung cancer in never smokers: Pooled analyses of seven cohorts. International Journal of Cancer, 2021, 149, 2010-2019.	5.1	8
7	Poor suitability for psychotherapy – a risk factor for treatment non-attendance?. Journal of Affective Disorders, 2021, 295, 1432-1439.	4.1	4
8	Predicting sudden cardiac death in a general population using an electrocardiographic risk score. Heart, 2020, 106, 427-433.	2.9	35
9	Estimating expected life-years and risk factor associations with mortality in Finland: cohort study. BMJ Open, 2020, 10, e033741.	1.9	15
10	Pooling of Finnish population-based health studies: lifestyle risk factors of colorectal and lung cancer. Acta Oncológica, 2020, 59, 1338-1342.	1.8	7
11	Intentional weight loss as a predictor of type 2 diabetes occurrence in a general adult population. BMJ Open Diabetes Research and Care, 2020, 8, e001560.	2.8	2
12	Orthogonal P-wave morphology, conventional P-wave indices, and the risk of atrial fibrillation in the general population using data from the Finnish Hospital Discharge Register. Europace, 2020, 22, 1173-1181.	1.7	20
13	Electrocardiographic Risk Markers of Cardiac Death: Gender Differences in the General Population. Frontiers in Physiology, 2020, 11, 578059.	2.8	3
14	The Effectiveness of Three Psychotherapies of Different Type and Length in the Treatment of Patients Suffering from Anxiety Disorders. European Family Therapy Association Series, 2020, , 349-366.	0.3	2
15	Self-Report Dieters: Who Are They?. Nutrients, 2019, 11, 1789.	4.1	9
16	Dietary magnesium intake, serum high sensitivity C-reactive protein and the risk of incident knee osteoarthritis leading to hospitalization—A cohort study of 4,953 Finns. PLoS ONE, 2019, 14, e0214064.	2.5	19
17	Associations of Metabolic Syndrome and Its Components with the Risk of Incident Knee Osteoarthritis Leading to Hospitalization: A 32-Year Follow-up Study. Cartilage, 2019, , 194760351989473.	2.7	8
18	Airway obstruction, serum vitamin D and mortality in a 33-year follow-up study. European Journal of Clinical Nutrition, 2019, 73, 1024-1032.	2.9	6

Paul Knekt

#	Article	IF	CITATIONS
19	Effectiveness of psychoanalysis and long-term psychodynamic psychotherapy on personality and social functioning 10 years after start of treatment. Psychiatry Research, 2019, 272, 774-783.	3.3	9
20	Associations of early childhood adversities with mental disorders, psychological functioning, and suitability for psychotherapy in adulthood. Psychiatry Research, 2018, 264, 366-373.	3.3	14
21	Childhood adversities as predictors of improvement in psychiatric symptoms and global functioning in solution-focused and short- and long-term psychodynamic psychotherapy during a 5-year follow-up. Journal of Affective Disorders, 2018, 235, 525-534.	4.1	7
22	Airway obstruction and the risk of myocardial infarction and death from coronary heart disease: a national health examination survey with a 33-year follow-up period. European Journal of Epidemiology, 2018, 33, 89-98.	5.7	9
23	Follow-up of the effectiveness of long-term psychodynamic psychotherapy and psychoanalysis 5 years after the end of psychoanalysis: Minor differences in psychiatric symptoms and work ability. Neurology Psychiatry and Brain Research, 2018, 30, 163-166.	2.0	1
24	Genetic analysis of over 1 million people identifies 535 new loci associated with blood pressure traits. Nature Genetics, 2018, 50, 1412-1425.	21.4	924
25	The impact of psychotherapy treatments of different length and type on health behaviour during a five-year follow-up. Psychotherapy Research, 2017, 27, 397-409.	1.8	3
26	The prediction of the level of personality organization on reduction of psychiatric symptoms and improvement of work ability in short―versus longâ€ŧerm psychotherapies during a 5â€year followâ€up. Psychology and Psychotherapy: Theory, Research and Practice, 2017, 90, 353-376.	2.5	16
27	Dieting attempts modify the association between quality of diet and obesity. Nutrition Research, 2017, 45, 63-72.	2.9	7
28	Effects of Scheduled Waiting for Psychotherapy in Patients With Major Depression. Journal of Nervous and Mental Disease, 2017, 205, 611-617.	1.0	11
29	Cohort Profile: The Finnish Mobile Clinic Health Surveys FMC, FMCF and MFS. International Journal of Epidemiology, 2017, 46, 1760-1761i.	1.9	24
30	Dispositional optimism as predictor of outcome in short―and longâ€ŧerm psychotherapy. Psychology and Psychotherapy: Theory, Research and Practice, 2017, 90, 279-298.	2.5	12
31	Tuberculosis, Airway Obstruction and Mortality in a Finnish Population. COPD: Journal of Chronic Obstructive Pulmonary Disease, 2017, 14, 143-149.	1.6	7
32	Patients' pre-treatment interpersonal problems as predictors of therapeutic alliance in long-term psychodynamic psychotherapy. Psychiatry Research, 2016, 241, 110-117.	3.3	15
33	Diabetes, glucose tolerance, and the risk of sudden cardiac death. BMC Cardiovascular Disorders, 2016, 16, 51.	1.7	38
34	Estimating efficacy in the presence of non-ignorable non-trial interventions in the Helsinki Psychotherapy Study. Statistical Methods in Medical Research, 2016, 25, 885-901.	1.5	5
35	The cost-effectiveness of short-term and long-term psychotherapy in the treatment of depressive and anxiety disorders during a 5-year follow-up. Journal of Affective Disorders, 2016, 190, 254-263.	4.1	31
36	Higher serum 25-hydroxyvitamin D concentrations are related to a reduced risk of depression. British Journal of Nutrition, 2015, 113, 1418-1426.	2.3	47

PAUL KNEKT

#	Article	IF	CITATIONS
37	Prospective study on the components of metabolic syndrome and the incidence of Parkinson's disease. Parkinsonism and Related Disorders, 2015, 21, 1148-1155.	2.2	52
38	Association between all-cause and cause-specific mortality and the GOLD stages 1–4: A 30-year follow-up among Finnish adults. Respiratory Medicine, 2015, 109, 1012-1018.	2.9	23
39	Randomized trial on the effectiveness of long- and short-term psychotherapy on psychosocial functioning and quality of life during a 5-year follow-up. Psychiatry Research, 2015, 229, 381-388.	3.3	18
40	The effectiveness of short- and long-term psychotherapy on personality functioning during a 5-year follow-up. Journal of Affective Disorders, 2015, 173, 31-38.	4.1	35
41	Therapists' Professional and Personal Characteristics as Predictors of Working Alliance in Shortâ€Term and Longâ€Term Psychotherapies. Clinical Psychology and Psychotherapy, 2014, 21, 475-494.	2.7	40
42	Selfâ€reported immature defense style as a predictor of outcome in shortâ€term and longâ€term psychotherapy. Brain and Behavior, 2014, 4, 495-503.	2.2	9
43	Intelligence as a predictor of outcome in short- and long-term psychotherapy. Psychiatry Research, 2014, 220, 1019-1027.	3.3	9
44	Serum 25-Hydroxyvitamin D Concentration and Risk of Dementia. Epidemiology, 2014, 25, 799-804.	2.7	32
45	Social support as a predictor of the outcome of depressive and anxiety disorder in short-term and long-term psychotherapy. Psychiatry Research, 2014, 216, 44-51.	3.3	37
46	Serum 25-hydroxyvitamin D and the risk of knee and hip osteoarthritis leading to hospitalization: a cohort study of 5274 Finns. Rheumatology, 2014, 53, 1778-1782.	1.9	21
47	Reduced risk of Parkinson's disease associated with lower body mass index and heavy leisure-time physical activity. European Journal of Epidemiology, 2014, 29, 285-292.	5.7	60
48	Self-concept and quality of object relations as predictors of outcome in short- and long-term psychotherapy. Journal of Affective Disorders, 2014, 152-154, 202-211.	4.1	17
49	Vitamin D status is associated with sociodemographic factors, lifestyle and metabolic health. European Journal of Nutrition, 2013, 52, 513-525.	3.9	80
50	Fatty liver score and 15-year incidence of type 2 diabetes. Hepatology International, 2013, 7, 610-621.	4.2	11
51	Sagittal Abdominal Diameter as a New Predictor for Incident Diabetes. Diabetes Care, 2013, 36, 283-288.	8.6	43
52	Randomized trial on the effectiveness of long- and short-term psychotherapy on psychiatric symptoms and working ability during a 5-year follow-up. Nordic Journal of Psychiatry, 2013, 67, 59-68.	1.3	46
53	Psychological predictors of the recovery from mood or anxiety disorder in short-term and long-term psychotherapy during a 3-year follow-up. Psychiatry Research, 2013, 208, 162-173.	3.3	25
54	Concurrent Validity of the Quality of Object Relations Scale (QORS) in Relation to Proxy Assessment of the Theoretical Scale Constituents. Psychopathology, 2013, 46, 111-119.	1.5	7

PAUL KNEKT

#	Article	IF	CITATIONS
55	Quality of object relations modifies the effectiveness of short- and long-term psychotherapy on self-concept. Open Journal of Psychiatry, 2013, 03, 345-350.	0.6	6
56	The Effectiveness of Solution-Focused Therapy and Short- and Long-Term Psychodynamic Psychotherapy on Self-Concept During a 3-Year Follow-Up. Journal of Nervous and Mental Disease, 2012, 200, 946-953.	1.0	17
57	Association between the Rorschach Ego Impairment Index and the Level of Personality Organization interview assessment in depressive and anxiety disorder patients. Psychiatry Research, 2012, 200, 849-856.	3.3	14
58	Suitability for Psychotherapy Scale (SPS) and its reliability, validity, and prediction. British Journal of Clinical Psychology, 2012, 51, 351-375.	3.5	28
59	Therapists' professional and personal characteristics as predictors of outcome in short- and long-term psychotherapy. Journal of Affective Disorders, 2012, 138, 301-312.	4.1	74
60	The Helsinki Psychotherapy Study: Effectiveness, Sufficiency, and Suitability of Short- and Long-Term Psychotherapy. , 2012, , 71-94.		7
61	Quasi-experimental study on the effectiveness of psychoanalysis, long-term and short-term psychotherapy on psychiatric symptoms, work ability and functional capacity during a 5-year follow-up. Journal of Affective Disorders, 2011, 132, 37-47.	4.1	91
62	Use of auxiliary psychiatric treatment during a 5-year follow-up among patients receiving short- or long-term psychotherapy. Journal of Affective Disorders, 2011, 135, 221-230.	4.1	32
63	Self-Administered Questionnaire Is a Reliable Measure of Coffee Consumption. Journal of Epidemiology, 2010, 20, 363-369.	2.4	6
64	The relative importance of modifiable potential risk factors of type 2 diabetes: a meta-analysis of two cohorts. European Journal of Epidemiology, 2010, 25, 115-124.	5.7	76
65	Serum Vitamin D and the Risk of Parkinson Disease. Archives of Neurology, 2010, 67, 808-11.	4.5	267
66	Vitamin D Status and the Risk of Cardiovascular Disease Death. American Journal of Epidemiology, 2009, 170, 1032-1039.	3.4	250
67	Validity of Hospital Discharge Data in a Prospective Epidemiological Study on Stroke and Myocardial Infarction. Acta Medica Scandinavica, 2009, 216, 309-315.	0.0	92
68	Effectiveness of short-term and long-term psychotherapy on work ability and functional capacity — A randomized clinical trial on depressive and anxiety disorders. Journal of Affective Disorders, 2008, 107, 95-106.	4.1	89
69	Vitamin D Status and the Risk of Lung Cancer: A Cohort Study in Finland. Cancer Epidemiology Biomarkers and Prevention, 2008, 17, 3274-3278.	2.5	90
70	Serum Vitamin D and Subsequent Occurrence of Type 2 Diabetes. Epidemiology, 2008, 19, 666-671.	2.7	286
71	Pretreatment clinical and psychosocial predictors of remission from depression after short-term psychodynamic psychotherapy and solution-focused therapy: a 1-year follow-up study. Psychotherapy Research, 2008, 18, 191-199.	1.8	8
72	Serum 25-Hydroxyvitamin D Concentration and Subsequent Risk of Type 2 Diabetes. Diabetes Care, 2007, 30, 2569-2570.	8.6	283

PAUL KNEKT

#	Article	IF	CITATIONS
73	Validity of a food frequency questionnaire varied by age and body mass index. Journal of Clinical Epidemiology, 2006, 59, 994-1001.	5.0	157
74	Schizophrenia, neuroleptic medication and mortality. British Journal of Psychiatry, 2006, 188, 122-127.	2.8	317
75	Antioxidant vitamins and coronary heart disease risk: a pooled analysis of 9 cohorts. American Journal of Clinical Nutrition, 2004, 80, 1508-1520.	4.7	258
76	Whole-grain and fiber intake and the incidence of type 2 diabetes. American Journal of Clinical Nutrition, 2003, 77, 622-629.	4.7	511
77	Mental disorders and cause-specific mortality. British Journal of Psychiatry, 2001, 179, 498-502.	2.8	285
78	Risk Factors for Injurious Falls Leading to Hospitalization or Death in a Cohort of 19,500 Adults. American Journal of Epidemiology, 1993, 138, 384-394.	3.4	145
79	What Works for Whom: Patients' Psychological Resources and Vulnerabilities as Common and Specific Predictors of Working Alliance in Different Psychotherapies. Frontiers in Psychiatry, 0, 13, .	2.6	4