

Adam B Glick

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3676862/publications.pdf>

Version: 2024-02-01

26
papers

1,067
citations

471509

17
h-index

580821

25
g-index

26
all docs

26
docs citations

26
times ranked

1722
citing authors

#	ARTICLE	IF	CITATIONS
1	Conditional Gene Expression in the Epidermis of Transgenic Mice Using the Tetracycline-Regulated Transactivators tTA and rTA Linked to the Keratin 5 Promoter. <i>Journal of Investigative Dermatology</i> , 2000, 115, 788-794.	0.7	194
2	[1] Isolation and utilization of epidermal keratinocytes for oncogene research. <i>Methods in Enzymology</i> , 1995, 254, 3-20.	1.0	137
3	Defects in TGF β signaling overcome senescence of mouse keratinocytes expressing v-rasHa. <i>Oncogene</i> , 2000, 19, 1698-1709.	5.9	101
4	Genetic and Pharmacological Analysis Identifies a Physiological Role for the AHR in Epidermal Differentiation. <i>Journal of Investigative Dermatology</i> , 2015, 135, 1320-1328.	0.7	86
5	Smad3 regulates senescence and malignant conversion in a mouse multistage skin carcinogenesis model. <i>Cancer Research</i> , 2003, 63, 3447-52.	0.9	79
6	ER stress and distinct outputs of the IRE1 \pm RNase control proliferation and senescence in response to oncogenic Ras. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, 9900-9905.	7.1	58
7	Induction of p16 ^{ink4a} and p19 ^{ARF} by TGF β 1 contributes to growth arrest and senescence response in mouse keratinocytes. <i>Molecular Carcinogenesis</i> , 2009, 48, 181-186.	2.7	54
8	CD8+ T Cells Mediate RAS-Induced Psoriasis-Like Skin Inflammation through IFN- β . <i>Journal of Investigative Dermatology</i> , 2013, 133, 955-963.	0.7	43
9	The Nuclear Receptor Peroxisome Proliferator-activated Receptor- β (PPAR β) Promotes Oncogene-induced Cellular Senescence through Repression of Endoplasmic Reticulum Stress. <i>Journal of Biological Chemistry</i> , 2014, 289, 20102-20119.	3.4	39
10	Ultrasound-Guided Cytosolic Protein Delivery <i>via</i> Transient Fluorous Masks. <i>ACS Nano</i> , 2020, 14, 4061-4073.	14.6	36
11	Use of a TGF β type I receptor inhibitor in mouse skin carcinogenesis reveals a dual role for TGF β signaling in tumor promotion and progression. <i>Carcinogenesis</i> , 2010, 31, 2127-2135.	2.8	27
12	The TGF β 1 pathway is required for NF κ B dependent gene expression in mouse keratinocytes. <i>Cytokine</i> , 2013, 64, 652-659.	3.2	25
13	Editor's Highlight: Ah Receptor Activation Potentiates Neutrophil Chemoattractant (C-X-C Motif) Ligand 5 Expression in Keratinocytes and Skin. <i>Toxicological Sciences</i> , 2017, 160, 83-94.	3.1	25
14	Tumor suppressor and oncogene actions of TGF β 1 occur early in skin carcinogenesis and are mediated by Smad3. <i>Molecular Carcinogenesis</i> , 2009, 48, 441-453.	2.7	23
15	The high-risk benign tumor: Evidence from the two-stage skin cancer model and relevance for human cancer. <i>Molecular Carcinogenesis</i> , 2007, 46, 605-610.	2.7	21
16	TGF β 1-Induced Inflammation in Premalignant Epidermal Squamous Lesions Requires IL-17. <i>Journal of Investigative Dermatology</i> , 2010, 130, 2295-2303.	0.7	21
17	Transforming growth factor β 1 enhances tumor promotion in mouse skin carcinogenesis. <i>Carcinogenesis</i> , 2010, 31, 1116-1123.	2.8	20
18	Photocontrolled miR-148b nanoparticles cause apoptosis, inflammation and regression of Ras induced epidermal squamous cell carcinomas in mice. <i>Biomaterials</i> , 2020, 256, 120212.	11.4	16

#	ARTICLE	IF	CITATIONS
19	Tumor-promoting role of TGF β 1 signaling in ultraviolet B-induced skin carcinogenesis is associated with cutaneous inflammation and lymph node migration of dermal dendritic cells. <i>Carcinogenesis</i> , 2014, 35, 959-966.	2.8	15
20	The multiple roles of the unfolded protein response regulator IRE1 β in cancer. <i>Molecular Carcinogenesis</i> , 2019, 58, 1623-1630.	2.7	14
21	Context-dependent regulation of cutaneous immunological responses by TGF β 1 and its role in skin carcinogenesis. <i>Carcinogenesis</i> , 2007, 29, 9-14.	2.8	13
22	Pharmacologic Inhibition of ALK5 Causes Selective Induction of Terminal Differentiation in Mouse Keratinocytes Expressing Oncogenic <i>HRAS</i> . <i>Molecular Cancer Research</i> , 2011, 9, 746-756.	3.4	7
23	Differentiated State of Initiating Tumor Cells Is Key to Distinctive Immune Responses Seen in H-RasG12V-Induced Squamous Tumors. <i>Cancer Immunology Research</i> , 2017, 5, 198-210.	3.4	7
24	The Endoplasmic Reticulum Stress Sensor IRE1 β Regulates the UV DNA Repair Response through the Control of Intracellular Calcium Homeostasis. <i>Journal of Investigative Dermatology</i> , 2022, 142, 1682-1691.e7.	0.7	4
25	Targeted deletion of TGF β 1 in basal keratinocytes causes profound defects in stratified squamous epithelia and aberrant melanocyte migration. <i>Developmental Biology</i> , 2022, 485, 9-23.	2.0	2
26	A New Xenotransplantation Model Reveals Tumor-Initiating Cells in Cutaneous Squamous Cell Carcinoma. <i>Journal of Investigative Dermatology</i> , 2012, 132, 261-262.	0.7	0