

Katarzyna MiÄkus

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3696128/publications.pdf>

Version: 2024-02-01

29
papers

1,460
citations

516710

16
h-index

501196

28
g-index

29
all docs

29
docs citations

29
times ranked

2398
citing authors

#	ARTICLE	IF	CITATIONS
1	MCPIP1 regulates focal adhesion kinase and Rho GTPase-dependent migration in clear cell renal cell carcinoma. <i>European Journal of Pharmacology</i> , 2022, 922, 174804.	3.5	0
2	Fatty Acids and a High-Fat Diet Induce Epithelial-Mesenchymal Transition by Activating TGF β 2 and β -Catenin in Liver Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 1272.	4.1	9
3	Deletion of <i>Mcpip1</i> in <i>Mcpip1^{fl/fl}AlbCre</i> mice recapitulates the phenotype of human primary biliary cholangitis. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2021, 1867, 166086.	3.8	12
4	MCPIP1 inhibits Wnt/ β -catenin signaling pathway activity and modulates epithelial-mesenchymal transition during clear cell renal cell carcinoma progression by targeting miRNAs. <i>Oncogene</i> , 2021, 40, 6720-6735.	5.9	21
5	The anti-inflammatory protein MCPIP1 inhibits the development of ccRCC by maintaining high levels of tumour suppressors. <i>European Journal of Pharmacology</i> , 2020, 888, 173591.	3.5	10
6	New therapeutic strategies in nonalcoholic fatty liver disease: a focus on promising drugs for nonalcoholic steatohepatitis. <i>Pharmacological Reports</i> , 2020, 72, 1-12.	3.3	68
7	Activity of MCPIP1 RNase in tumor associated processes. <i>Journal of Experimental and Clinical Cancer Research</i> , 2019, 38, 421.	8.6	19
8	C-Met as a Key Factor Responsible for Sustaining Undifferentiated Phenotype and Therapy Resistance in Renal Carcinomas. <i>Cells</i> , 2019, 8, 272.	4.1	21
9	RNA sequencing reveals widespread transcriptome changes in a renal carcinoma cell line. <i>Oncotarget</i> , 2018, 9, 8597-8613.	1.8	22
10	MCPIP1 contributes to clear cell renal cell carcinomas development. <i>Angiogenesis</i> , 2017, 20, 325-340.	7.2	61
11	The Met tyrosine kinase receptor as a therapeutic target and a potential cancer stem cell factor responsible for therapy resistance. <i>Oncology Reports</i> , 2017, 37, 647-656.	2.6	18
12	MCPIP1 Downregulation in Clear Cell Renal Cell Carcinoma Promotes Vascularization and Metastatic Progression. <i>Cancer Research</i> , 2017, 77, 4905-4920.	0.9	60
13	MET receptor is a potential therapeutic target in high grade cervical cancer. <i>Oncotarget</i> , 2015, 6, 10086-10101.	1.8	15
14	Constitutive activation of MET signaling impairs myogenic differentiation of rhabdomyosarcoma and promotes its development and progression. <i>Oncotarget</i> , 2015, 6, 31378-31398.	1.8	25
15	Downregulation of the CXCR4 receptor inhibits cervical carcinoma metastatic behavior in vitro and in vivo. <i>International Journal of Oncology</i> , 2014, 44, 1853-1860.	3.3	10
16	Multifunctional protein APPL2 contributes to survival of human glioma cells. <i>Molecular Oncology</i> , 2013, 7, 67-84.	4.6	16
17	17AEP-GA, an HSP90 antagonist, is a potent inhibitor of glioblastoma cell proliferation, survival, migration and invasion. <i>Oncology Reports</i> , 2012, 28, 1903-1909.	2.6	15
18	Fenofibrate attenuates contact-stimulated cell motility and gap junctional coupling in DU-145 human prostate cancer cell populations. <i>Oncology Reports</i> , 2011, 26, 447-53.	2.6	24

#	ARTICLE	IF	CITATIONS
19	Genetically modified adipose tissue ^â derived mesenchymal stem cells overexpressing CXCR4 display increased motility, invasiveness, and homing to bone marrow of NOD/SCID mice. <i>Experimental Hematology</i> , 2011, 39, 686-696.e4.	0.4	85
20	Differential expression of Snail1 transcription factor and Snail1-related genes in alveolar and embryonal rhabdomyosarcoma subtypes.. <i>Folia Histochemica Et Cytobiologica</i> , 2011, 48, 671-7.	1.5	10
21	Role of I-TAC-binding receptors CXCR3 and CXCR7 in proliferation, activation of intracellular signaling pathways and migration of various tumor cell lines.. <i>Folia Histochemica Et Cytobiologica</i> , 2010, 48, 104-11.	1.5	21
22	Inhibition of rhabdomyosarcoma's metastatic behavior through downregulation of MET receptor signaling.. <i>Folia Histochemica Et Cytobiologica</i> , 2010, 47, 485-9.	1.5	17
23	Topographical control of prostate cancer cell migration. <i>Molecular Medicine Reports</i> , 2009, 2, 865-71.	2.4	8
24	Optimization of a synthetic siRNA delivery for the treatment of rhabdomyosarcoma. <i>Open Life Sciences</i> , 2008, 3, 371-379.	1.4	2
25	Leukemia Inhibitory Factor: A Newly Identified Metastatic Factor in Rhabdomyosarcomas. <i>Cancer Research</i> , 2007, 67, 2131-2140.	0.9	94
26	Genistein inhibits the contact-stimulated migration of prostate cancer cells. <i>Cellular and Molecular Biology Letters</i> , 2007, 12, 348-61.	7.0	26
27	Trafficking of Normal Stem Cells and Metastasis of Cancer Stem Cells Involve Similar Mechanisms: Pivotal Role of the SDF ^â â€“CXCR4 Axis. <i>Stem Cells</i> , 2005, 23, 879-894.	3.2	709
28	Contact stimulation of prostate cancer cell migration: the role of gap junctional coupling and migration stimulated by heterotypic cell-to-cell contacts in determination of the metastatic phenotype of Dunning rat prostate cancer cells. <i>Biology of the Cell</i> , 2005, 97, 893-903.	2.0	41
29	Contact-activated migration of melanoma B16 and sarcoma XC cells. <i>Biochemistry and Cell Biology</i> , 2001, 79, 425-440.	2.0	21