

Zhen-Yu He

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/371926/publications.pdf>

Version: 2024-02-01

132
papers

2,540
citations

257450

24
h-index

302126

39
g-index

137
all docs

137
docs citations

137
times ranked

4204
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | OUP accepted manuscript. BJS Open, 2022, 6, . | 1.7 | 0 |
| 2 | Prognostic significance of the skeletal muscle index and systemic inflammatory index in patients with lymph node-positive breast cancer after radical mastectomy. BMC Cancer, 2022, 22, 234. | 2.6 | 10 |
| 3 | The Predictive Effect of the 8th AJCC Pathological Prognostic Staging on the Benefit of Postmastectomy Radiotherapy in N2/N3 Breast Cancer. Breast Cancer: Targets and Therapy, 2022, Volume 14, 133-144. | 1.8 | 1 |
| 4 | Effect of Capecitabine Maintenance Therapy Using Lower Dosage and Higher Frequency vs Observation on Disease-Free Survival Among Patients With Early-Stage Triple-Negative Breast Cancer Who Had Received Standard Treatment. JAMA - Journal of the American Medical Association, 2021, 325, 50. | 7.4 | 113 |
| 5 | The prognostic and predictive value of the 8th American Joint Committee on Cancer (AJCC) staging system among early breast cancer patients aged <50 years. Gland Surgery, 2021, 10, 233-241. | 1.1 | 2 |
| 6 | Additional radiotherapy to breast-conserving surgery is an optional treatment for de novo stage IV breast cancer: A population-based analysis. Cancer Medicine, 2021, 10, 1634-1643. | 2.8 | 6 |
| 7 | Identification of MEC8/miR-378d/SOBP axis as a novel regulatory network and associated with immune infiltrates in ovarian carcinoma by integrated bioinformatics analysis. Cancer Medicine, 2021, 10, 2924-2939. | 2.8 | 9 |
| 8 | Triple-negative breast cancer outcomes: Does AJCC 8th staging improve chemotherapy decision-making. Breast, 2021, 59, 117-123. | 2.2 | 4 |
| 9 | Prognostic significance of the skeletal muscle index and an inflammation biomarker in patients with breast cancer who underwent postoperative adjuvant radiotherapy. Current Problems in Cancer, 2020, 44, 100513. | 2.0 | 12 |
| 10 | Should women with early breast cancer under 40 years of age have a routine 21-gene recurrence score testing: A SEER database study. Breast, 2020, 49, 233-241. | 2.2 | 8 |
| 11 | Prognostic significance of the Controlling Nutritional Status (CONUT) score in surgically treated breast cancer patients. Gland Surgery, 2020, 9, 1370-1379. | 1.1 | 13 |
| 12 | The effect of postmastectomy radiotherapy in node-positive triple-negative breast cancer. BMC Cancer, 2020, 20, 1146. | 2.6 | 7 |
| 13 | The Prognostic Prediction Value of Systemic Inflammation Score and the Development of a Nomogram for Patients With Surgically Treated Breast Cancer. Frontiers in Oncology, 2020, 10, 563731. | 2.8 | 15 |
| 14 | Incorporation of biologic factors for the staging of de novo stage IV breast cancer. Npj Breast Cancer, 2020, 6, 43. | 5.2 | 7 |
| 15 | Prognostic and Predictive Value of the American Joint Committee on Cancer Pathological Prognostic Staging System in Nodal Micrometastatic Breast Cancer. Frontiers in Oncology, 2020, 10, 570175. | 2.8 | 1 |
| 16 | Aggressive Local Treatment Improves Survival in Stage IV Breast Cancer With Synchronous Metastasis. Frontiers in Oncology, 2020, 10, 522580. | 2.8 | 7 |
| 17 | Real-world impact of postmastectomy radiotherapy in T1-2 breast cancer with one to three positive lymph nodes. Annals of Translational Medicine, 2020, 8, 489-489. | 1.7 | 6 |
| 18 | Evaluation of the 8th edition of the American joint committee on cancer's pathological staging system in prognosis assessment and treatment decision making for stage T1-2N1 breast cancer after mastectomy. Breast, 2020, 51, 2-10. | 2.2 | 13 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Prognostic validation and therapeutic decision-making of the AJCC eighth pathological prognostic staging for T3N0 breast cancer after mastectomy. <i>Clinical and Translational Medicine</i> , 2020, 10, 125-136. | 4.0 | 11 |
| 20 | The preoperative systemic inflammation response index (SIRI) independently predicts survival in postmenopausal women with breast cancer. <i>Current Problems in Cancer</i> , 2020, 44, 100560. | 2.0 | 34 |
| 21 | Chemotherapy and 21-gene recurrence score testing for older breast cancer patients: A competing-risks analysis. <i>Breast</i> , 2020, 54, 319-327. | 2.2 | 8 |
| 22 | The longitudinal risk of mortality between invasive ductal carcinoma and metaplastic breast carcinoma. <i>Scientific Reports</i> , 2020, 10, 22070. | 3.3 | 8 |
| 23 | Prognostic validation and treatment decision making of the 8th edition of the American Joint Committee on Cancer pathological staging system for elderly women with early-stage breast cancer. <i>Aging</i> , 2020, 12, 15077-15090. | 3.1 | 2 |
| 24 | The 1-year mortality after radiotherapy for nasopharyngeal carcinoma: a population-based analysis. <i>Future Oncology</i> , 2019, 15, 3357-3365. | 2.4 | 2 |
| 25 | Progesterone receptor status and tumor grade predict the 21-gene recurrence score of invasive lobular breast cancer. <i>Biomarkers in Medicine</i> , 2019, 13, 1005-1012. | 1.4 | 8 |
| 26 | Omission of adjuvant radiotherapy following breast-conserving surgery for elderly women with early-stage pure mucinous breast carcinoma. <i>Radiation Oncology</i> , 2019, 14, 190. | 2.7 | 9 |
| 27 | Real-World Impact of Survival by Period of Diagnosis in Epithelial Ovarian Cancer Between 1990 and 2014. <i>Frontiers in Oncology</i> , 2019, 9, 639. | 2.8 | 31 |
| 28 | The Effect of Post-mastectomy Radiotherapy in Patients With Metaplastic Breast Cancer: An Analysis of SEER Database. <i>Frontiers in Oncology</i> , 2019, 9, 747. | 2.8 | 11 |
| 29 | Lymph Node Status and Outcomes for Nasopharyngeal Carcinoma According to Histological Subtypes: A SEER Population-Based Retrospective Analysis. <i>Advances in Therapy</i> , 2019, 36, 3123-3133. | 2.9 | 18 |
| 30 | The Role of Axillary Lymph Node Dissection in Tubular Carcinoma of the Breast: A Population Database Study. <i>Medical Science Monitor</i> , 2019, 25, 880-887. | 1.1 | 5 |
| 31 | 21-Gene Recurrence Score Assay Could Not Predict Benefit of Post-mastectomy Radiotherapy in T1-2 N1mic ER-Positive HER2-Negative Breast Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 270. | 2.8 | 8 |
| 32 | Efficacy of controlled-release oxycodone for reducing pain due to oral mucositis in nasopharyngeal carcinoma patients treated with concurrent chemoradiotherapy: a prospective clinical trial. <i>Supportive Care in Cancer</i> , 2019, 27, 3759-3767. | 2.2 | 18 |
| 33 | Noninferior Outcome After Breast-Conserving Treatment Compared to Mastectomy in Breast Cancer Patients With Four or More Positive Lymph Nodes. <i>Frontiers in Oncology</i> , 2019, 9, 143. | 2.8 | 9 |
| 34 | The 21-gene recurrence score and effects of adjuvant radiotherapy after breast conserving surgery in early-stage breast cancer. <i>Future Oncology</i> , 2019, 15, 1629-1639. | 2.4 | 8 |
| 35 | <p></p>Effect of 21-gene recurrence score in decision-making for surgery in early stage breast cancer</p>. <i>OncoTargets and Therapy</i> , 2019, Volume 12, 2071-2078. | 2.0 | 2 |
| 36 | 21-Gene Recurrence Score Assay and Outcomes of Adjuvant Radiotherapy in Elderly Women With Early-Stage Breast Cancer After Breast-Conserving Surgery. <i>Frontiers in Oncology</i> , 2019, 9, 1. | 2.8 | 139 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | The effect of histological subtypes on survival outcome in nasopharyngeal carcinoma after extensive follow up. <i>Annals of Translational Medicine</i> , 2019, 7, 768-768. | 1.7 | 22 |
| 38 | Prognostic value of skeletal muscle index and monocyte-to-lymphocyte ratio for lymph node-positive breast cancer patients after mastectomy. <i>Annals of Translational Medicine</i> , 2019, 7, 775-775. | 1.7 | 12 |
| 39 | Inflammatory breast cancer outcomes by breast cancer subtype: a population-based study. <i>Future Oncology</i> , 2019, 15, 507-516. | 2.4 | 16 |
| 40 | 21-gene recurrence score and adjuvant chemotherapy decisions in patients with invasive lobular breast cancer. <i>Biomarkers in Medicine</i> , 2019, 13, 83-93. | 1.4 | 19 |
| 41 | The Value of Prognostic Nutritional Index (PNI) in Predicting Survival and Guiding Radiotherapy of Patients With T1-2N1 Breast Cancer. <i>Frontiers in Oncology</i> , 2019, 9, 1562. | 2.8 | 45 |
| 42 | Impact of the 21-gene recurrence score assay on chemotherapy decision making and outcomes for breast cancer patients with four or more positive lymph nodes. <i>Annals of Translational Medicine</i> , 2019, 7, 446-446. | 1.7 | 5 |
| 43 | KIF11 Functions as an Oncogene and Is Associated with Poor Outcomes from Breast Cancer. <i>Cancer Research and Treatment</i> , 2019, 51, 1207-1221. | 3.0 | 47 |
| 44 | Impact of 21-Gene Recurrence Score on Chemotherapy Decision in Invasive Ductal Carcinoma of Breast with Nodal Micrometastases. <i>Cancer Research and Treatment</i> , 2019, 51, 1437-1448. | 3.0 | 4 |
| 45 | Bioinformatics-Based Discovery of CKLF-Like MARVEL Transmembrane Member 5 as a Novel Biomarker for Breast Cancer. <i>Frontiers in Cell and Developmental Biology</i> , 2019, 7, 361. | 3.7 | 4 |
| 46 | The effect of lymphadenectomy in advanced ovarian cancer according to residual tumor status: A population-based study. <i>International Journal of Surgery</i> , 2018, 52, 11-15. | 2.7 | 19 |
| 47 | Comparison of the effects of local treatment strategies in non-metastatic Ewing sarcoma of bone. <i>Expert Review of Anticancer Therapy</i> , 2018, 18, 501-506. | 2.4 | 4 |
| 48 | Survival in signet ring cell carcinoma varies based on primary tumor location: a Surveillance, Epidemiology, and End Results database analysis. <i>Expert Review of Gastroenterology and Hepatology</i> , 2018, 12, 209-214. | 3.0 | 50 |
| 49 | The effects of postoperative radiotherapy on survival outcomes in patients under 65 with estrogen receptor positive tubular breast carcinoma. <i>Radiation Oncology</i> , 2018, 13, 226. | 2.7 | 7 |
| 50 | The Distribution and Outcomes of the 21-Gene Recurrence Score in T1-T2N0 Estrogen Receptor-Positive Breast Cancer With Different Histologic Subtypes. <i>Frontiers in Genetics</i> , 2018, 9, 638. | 2.3 | 23 |
| 51 | The Effect of Histological Subtypes on Outcomes of Stage IV Epithelial Ovarian Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 577. | 2.8 | 25 |
| 52 | Clinicopathologic characteristics and clinical outcomes of pure type and mixed type of tubular carcinoma of the breast: a single-institution cohort study. <i>Cancer Management and Research</i> , 2018, Volume 10, 4509-4515. | 1.9 | 6 |
| 53 | The Clinicopathological Features and Survival Outcomes of Different Histological Subtypes in Triple-negative Breast Cancer. <i>Journal of Cancer</i> , 2018, 9, 296-303. | 2.5 | 60 |
| 54 | Prognostic value of ductal carcinoma in situ component in invasive ductal carcinoma of the breast: a Surveillance, Epidemiology, and End Results database analysis. <i>Cancer Management and Research</i> , 2018, Volume 10, 527-534. | 1.9 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Trends and Outcomes of Sentinel Lymph Node Biopsy in Early-stage Vulvar Squamous Cell Carcinoma: A Population-based Study. <i>Journal of Cancer</i> , 2018, 9, 1951-1957. | 2.5 | 7 |
| 56 | The Effect of Marital Status on Nasopharyngeal Carcinoma Survival: A Surveillance, Epidemiology and End Results Study. <i>Journal of Cancer</i> , 2018, 9, 1870-1876. | 2.5 | 23 |
| 57 | Long-term survival effect of the interval between mastectomy and radiotherapy in locally advanced breast cancer. <i>Cancer Management and Research</i> , 2018, Volume 10, 2047-2054. | 1.9 | 11 |
| 58 | Comparison of survival outcomes of locally advanced breast cancer patients receiving post-mastectomy radiotherapy with and without immediate breast reconstruction: a population-based analysis. <i>Cancer Management and Research</i> , 2018, Volume 10, 1993-2002. | 1.9 | 9 |
| 59 | Downregulation of hsa_circ_0011946 suppresses the migration and invasion of the breast cancer cell line MCF-7 by targeting RFC3. <i>Cancer Management and Research</i> , 2018, Volume 10, 535-544. | 1.9 | 75 |
| 60 | Prognostic Value of the Number of Removed Lymph Nodes in Vulvar Squamous Cell Carcinoma Patients With Node-Positive Disease: A Population-Based Study. <i>Frontiers in Oncology</i> , 2018, 8, 184. | 2.8 | 1 |
| 61 | Omission of Postoperative Radiotherapy in Women Aged 65 Years or Older With Tubular Carcinoma of the Breast After Breast-Conserving Surgery. <i>Frontiers in Oncology</i> , 2018, 8, 190. | 2.8 | 9 |
| 62 | Tubular carcinomas of the breast: an epidemiologic study. <i>Future Oncology</i> , 2018, 14, 3037-3047. | 2.4 | 4 |
| 63 | Widowed status increases the risk of death in vulvar cancer. <i>Future Oncology</i> , 2018, 14, 2589-2598. | 2.4 | 8 |
| 64 | Patterns of Distant Metastasis Between Histological Types in Esophageal Cancer. <i>Frontiers in Oncology</i> , 2018, 8, 302. | 2.8 | 52 |
| 65 | Survival benefits with the addition of adjuvant hysterectomy to radiochemotherapy for treatment of stage I adenocarcinoma of the uterine cervix. <i>Journal of Surgical Oncology</i> , 2018, 118, 574-580. | 1.7 | 5 |
| 66 | Lymph node ratio has prognostic value related to the number of positive lymph nodes in patients with vulvar cancer. <i>Future Oncology</i> , 2018, 14, 2343-2351. | 2.4 | 3 |
| 67 | Clinical Features of Brain Metastases in Small Cell Lung Cancer: an Implication for Hippocampal Sparing Whole Brain Radiation Therapy. <i>Translational Oncology</i> , 2017, 10, 54-58. | 3.7 | 14 |
| 68 | Thymosin beta 10 is a key regulator of tumorigenesis and metastasis and a novel serum marker in breast cancer. <i>Breast Cancer Research</i> , 2017, 19, 15. | 5.0 | 89 |
| 69 | Up-Regulation of RFC3 Promotes Triple Negative Breast Cancer Metastasis and is Associated With Poor Prognosis Via EMT. <i>Translational Oncology</i> , 2017, 10, 1-9. | 3.7 | 46 |
| 70 | The prognostic value of histologic subtype in node-positive early-stage cervical cancer after hysterectomy and adjuvant radiotherapy. <i>International Journal of Surgery</i> , 2017, 44, 1-6. | 2.7 | 13 |
| 71 | Men and women show similar survival outcome in stage IV breast cancer. <i>Breast</i> , 2017, 34, 115-121. | 2.2 | 6 |
| 72 | The effect of local treatment modalities in patients with early-stage adenocarcinoma of the uterine cervix: A population-based analysis. <i>International Journal of Surgery</i> , 2017, 41, 16-22. | 2.7 | 14 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 73 | Adjuvant radiation therapy and survival for adenoid cystic carcinoma of the breast. <i>Breast</i> , 2017, 31, 214-218. | 2.2 | 21 |
| 74 | Multimodal treatment including hysterectomy improves survival in patients with locally advanced cervical cancer: A population-based, propensity score-matched analysis. <i>International Journal of Surgery</i> , 2017, 48, 122-127. | 2.7 | 7 |
| 75 | Demographic and clinicopathological characteristics of nasopharyngeal carcinoma and survival outcomes according to age at diagnosis: A population-based analysis. <i>Oral Oncology</i> , 2017, 73, 83-87. | 1.5 | 40 |
| 76 | Preoperative radiotherapy improves survival in rectal signet-ring cell carcinoma-a population-based study. <i>Radiation Oncology</i> , 2017, 12, 141. | 2.7 | 12 |
| 77 | The effect of distant metastases sites on survival in de novo stage-IV breast cancer: A SEER database analysis. <i>Tumor Biology</i> , 2017, 39, 101042831770508. | 1.8 | 56 |
| 78 | Therapeutic role of axillary lymph node dissection in patients with stage IV breast cancer: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 467-474. | 2.5 | 7 |
| 79 | Comparison of clinical outcomes of squamous cell carcinoma, adenocarcinoma, and adenosquamous carcinoma of the uterine cervix after definitive radiotherapy: a population-based analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017, 143, 115-122. | 2.5 | 59 |
| 80 | Sites of metastasis and overall survival in esophageal cancer: a population-based study. <i>Cancer Management and Research</i> , 2017, Volume 9, 781-788. | 1.9 | 68 |
| 81 | Postoperative radiotherapy for invasive micropapillary carcinoma of the breast: an analysis of Surveillance, Epidemiology, and End Results database. <i>Cancer Management and Research</i> , 2017, Volume 9, 453-459. | 1.9 | 14 |
| 82 | The impact of examined lymph node count on survival in squamous cell carcinoma and adenocarcinoma of the uterine cervix. <i>Cancer Management and Research</i> , 2017, Volume 9, 315-322. | 1.9 | 19 |
| 83 | Comparison of survival outcomes between radical hysterectomy and definitive radiochemotherapy in stage IB1 and IIA1 cervical cancer. <i>Cancer Management and Research</i> , 2017, Volume 9, 813-819. | 1.9 | 13 |
| 84 | Comparable Survival between Additional Radiotherapy and Local Surgery in Occult Breast Cancer after Axillary Lymph Node Dissection: A Population-based Analysis. <i>Journal of Cancer</i> , 2017, 8, 3849-3855. | 2.5 | 13 |
| 85 | The survival benefits of local surgery in stage IV breast cancer are not affected by breast cancer subtypes: a population-based analysis. <i>Oncotarget</i> , 2017, 8, 67851-67860. | 1.8 | 10 |
| 86 | Incorporation of the number of positive lymph nodes leads to better prognostic discrimination of node-positive early stage cervical cancer. <i>Oncotarget</i> , 2017, 8, 26057-26065. | 1.8 | 12 |
| 87 | Clinicopathological features of small cell carcinoma of the uterine cervix in the surveillance, epidemiology, and end results database. <i>Oncotarget</i> , 2017, 8, 40425-40433. | 1.8 | 17 |
| 88 | Clinicopathological characteristics, treatment, and survival outcomes of cystadenocarcinoma of the salivary gland: a population-based study. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 6569-6572. | 2.0 | 7 |
| 89 | Progesterone receptor loss identifies hormone receptor-positive and HER2-negative breast cancer subgroups at higher risk of relapse: a retrospective cohort study. <i>OncoTargets and Therapy</i> , 2016, 9, 1707. | 2.0 | 9 |
| 90 | Use of CEA and CA15-3 to Predict Axillary Lymph Node Metastasis in Patients with Breast Cancer. <i>Journal of Cancer</i> , 2016, 7, 37-41. | 2.5 | 23 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 91 | Patterns of distant metastasis in Chinese women according to breast cancer subtypes. <i>Oncotarget</i> , 2016, 7, 47975-47984. | 1.8 | 23 |
| 92 | Clinical features of brain metastases in breast cancer: an implication for hippocampal-sparing whole-brain radiation therapy. <i>Therapeutics and Clinical Risk Management</i> , 2016, Volume 12, 1849-1853. | 2.0 | 11 |
| 93 | Differences in esophageal cancer characteristics and survival between Chinese and Caucasian patients in the SEER database. <i>OncoTargets and Therapy</i> , 2016, Volume 9, 6435-6444. | 2.0 | 12 |
| 94 | A novel prognostic score model incorporating CDGSH iron sulfur domain2 (CISD2) predicts risk of disease progression in laryngeal squamous cell carcinoma. <i>Oncotarget</i> , 2016, 7, 22720-22732. | 1.8 | 25 |
| 95 | Impact of the number of resected lymph nodes on survival after preoperative radiotherapy for esophageal cancer. <i>Oncotarget</i> , 2016, 7, 22497-22507. | 1.8 | 14 |
| 96 | Prognostic value of lymph node ratio in stage IIIC epithelial ovarian cancer with node-positive in a SEER population-based study. <i>Oncotarget</i> , 2016, 7, 7952-7959. | 1.8 | 22 |
| 97 | Surgery Combined with Radiotherapy Improved Survival in Metastatic Esophageal Cancer in a Surveillance Epidemiology and End Results Population-based Study. <i>Scientific Reports</i> , 2016, 6, 28280. | 3.3 | 31 |
| 98 | Lymph node dissection improved survival in patients with metastatic thoracic esophageal cancer: An analysis of 220 patients from the SEER database. <i>International Journal of Surgery</i> , 2016, 35, 13-18. | 2.7 | 6 |
| 99 | Patterns of Regional Lymph Node Recurrence After Radical Surgery for Thoracic Esophageal Squamous Cell Carcinoma. <i>Annals of Thoracic Surgery</i> , 2016, 101, 551-557. | 1.3 | 22 |
| 100 | Risk factors for lymph node metastasis in ovarian cancer: Implications for systematic lymphadenectomy. <i>International Journal of Surgery</i> , 2016, 29, 123-127. | 2.7 | 20 |
| 101 | The local treatment modalities in FIGO stage I small cell carcinoma of the cervix are determined by disease stage and lymph node status. <i>Cancer Medicine</i> , 2016, 5, 1108-1115. | 2.8 | 17 |
| 102 | The Activation of ERK1/2 and JNK MAPK Signaling by Insulin/IGF-1 Is Responsible for the Development of Colon Cancer with Type 2 Diabetes Mellitus. <i>PLoS ONE</i> , 2016, 11, e0149822. | 2.5 | 38 |
| 103 | Effect of postoperative radiotherapy for squamous cell cancer of the breast in a surveillance epidemiology and end results population-based study. <i>Oncotarget</i> , 2016, 7, 10684-10693. | 1.8 | 4 |
| 104 | Upregulation of E2F8 promotes cell proliferation and tumorigenicity in breast cancer by modulating G1/S phase transition. <i>Oncotarget</i> , 2016, 7, 23757-23771. | 1.8 | 46 |
| 105 | Lymph node ratio may predict the benefit of postoperative radiotherapy in node-positive cervical cancer. <i>Oncotarget</i> , 2016, 7, 29420-29428. | 1.8 | 16 |
| 106 | Breast Cancer Subtype is Associated With Axillary Lymph Node Metastasis. <i>Medicine (United States)</i> , 2015, 94, e2213. | 1.0 | 32 |
| 107 | Number of Negative Lymph Nodes Can Predict Survival after Postmastectomy Radiotherapy According to Different Breast Cancer Subtypes. <i>Journal of Cancer</i> , 2015, 6, 261-269. | 2.5 | 3 |
| 108 | Tailoring Pelvic Lymphadenectomy for Patients with Stage IA2, IB1, and IIA1 Uterine Cervical Cancer. <i>Journal of Cancer</i> , 2015, 6, 377-381. | 2.5 | 18 |

| # | ARTICLE | IF | CITATIONS |
|-----|---|-----|-----------|
| 109 | Prognostic Impact of ABO Blood Group on the Survival in Patients with Ovarian Cancer. <i>Journal of Cancer</i> , 2015, 6, 970-975. | 2.5 | 17 |
| 110 | Prognostic value of lymph node ratio in patients with small-cell carcinoma of the cervix based on data from a large national registry. <i>OncoTargets and Therapy</i> , 2015, 9, 67. | 2.0 | 2 |
| 111 | Dosimetric Comparison of the Simultaneous Integrated Boost in Whole-Breast Irradiation after Breast-Conserving Surgery: IMRT, IMRT plus an Electron Boost and VMAT. <i>PLoS ONE</i> , 2015, 10, e0120811. | 2.5 | 15 |
| 112 | Clinicopathologic features and treatment of breast metastasis from nasopharyngeal carcinoma: A report of two cases and literature review. <i>Oncology Letters</i> , 2015, 10, 3675-3681. | 1.8 | 4 |
| 113 | Influence of different treatment modalities on survival of patients with low-grade endometrial stromal sarcoma: A retrospective cohort study. <i>International Journal of Surgery</i> , 2015, 23, 147-151. | 2.7 | 24 |
| 114 | Prognostic Value of Different Lymph Node Staging Methods in Esophageal Squamous Cell Carcinoma After Esophagectomy. <i>Annals of Thoracic Surgery</i> , 2015, 99, 284-290. | 1.3 | 19 |
| 115 | Number of negative lymph nodes is associated with disease-free survival in patients with breast cancer. <i>BMC Cancer</i> , 2015, 15, 43. | 2.6 | 10 |
| 116 | Postmastectomy Radiotherapy Improves Disease-Free Survival of High Risk of Locoregional Recurrence Breast Cancer Patients with T1-2 and 1 to 3 Positive Nodes. <i>PLoS ONE</i> , 2015, 10, e0119105. | 2.5 | 22 |
| 117 | Prognosis of patients with esophageal squamous cell carcinoma after esophagectomy using the log odds of positive lymph nodes. <i>Oncotarget</i> , 2015, 6, 36911-36922. | 1.8 | 26 |
| 118 | Distribution of metastatic disease in the brain in relation to the hippocampus: a retrospective single-center analysis of 6064 metastases in 632 patients. <i>Oncotarget</i> , 2015, 6, 44030-44036. | 1.8 | 25 |
| 119 | Using the Lymph Node Ratio to Evaluate the Prognosis of Stage II/III Breast Cancer Patients Who Received Neoadjuvant Chemotherapy and Mastectomy. <i>Cancer Research and Treatment</i> , 2015, 47, 757-764. | 3.0 | 20 |
| 120 | Number of negative lymph nodes should be considered for incorporation into staging for breast cancer. <i>American Journal of Cancer Research</i> , 2015, 5, 844-53. | 1.4 | 20 |
| 121 | Transdermal fentanyl for pain due to chemoradiotherapy-induced oral mucositis in nasopharyngeal cancer patients: evaluating efficacy, safety, and improvement in quality of life. <i>Drug Design, Development and Therapy</i> , 2014, 8, 497. | 4.3 | 16 |
| 122 | Prognostic Value of Ki-67 in Breast Cancer Patients with Positive Axillary Lymph Nodes: A Retrospective Cohort Study. <i>PLoS ONE</i> , 2014, 9, e87264. | 2.5 | 33 |
| 123 | The value of radiotherapy in breast cancer patients with isolated ipsilateral supraclavicular lymph node metastasis without distant metastases at diagnosis: a retrospective analysis of Chinese patients. <i>OncoTargets and Therapy</i> , 2014, 7, 281. | 2.0 | 6 |
| 124 | Accelerated Partial Breast Irradiation with Intensity-Modulated Radiotherapy Is Feasible for Chinese Breast Cancer Patients. <i>Journal of Breast Cancer</i> , 2014, 17, 256. | 1.9 | 1 |
| 125 | Post-mastectomy radiotherapy can improve survival in breast cancer patients aged 35 years or younger with four or more positive nodes but not in one to three positive nodes. <i>Therapeutics and Clinical Risk Management</i> , 2014, 10, 867. | 2.0 | 3 |
| 126 | Number of negative lymph nodes can predict survival of breast cancer patients with four or more positive lymph nodes after postmastectomy radiotherapy. <i>Radiation Oncology</i> , 2014, 9, 284. | 2.7 | 12 |

| # | ARTICLE | IF | CITATIONS |
|-----|--|-----|-----------|
| 127 | Dosimetric analysis of the brachial plexus among patients with breast cancer treated with post-mastectomy radiotherapy to the ipsilateral supraclavicular area: report of 3 cases of radiation-induced brachial plexus neuropathy. <i>Radiation Oncology</i> , 2014, 9, 292. | 2.7 | 16 |
| 128 | Tumor location is a prognostic factor for survival of Chinese women with T1-2N0M0 breast cancer. <i>International Journal of Surgery</i> , 2014, 12, 394-398. | 2.7 | 28 |
| 129 | Serum levels of CEA and CA15-3 in different molecular subtypes and prognostic value in Chinese breast cancer. <i>Breast</i> , 2014, 23, 88-93. | 2.2 | 90 |
| 130 | Benefit of Post-mastectomy Radiotherapy of the Supra-/infraclavicular Lymphatic Drainage Area in Breast Cancer Patients. <i>Asian Pacific Journal of Cancer Prevention</i> , 2014, 15, 5557-5563. | 1.2 | 1 |
| 131 | Use of the Metastatic Lymph Node Ratio to Evaluate the Prognosis of Esophageal Cancer Patients with Node Metastasis Following Radical Esophagectomy. <i>PLoS ONE</i> , 2013, 8, e73446. | 2.5 | 24 |
| 132 | Ovarian Ablation Using Goserelin Improves Survival of Premenopausal Patients with Stage II/III Hormone Receptor-Positive Breast Cancer without Chemotherapy-Induced Amenorrhea. <i>Cancer Research and Treatment</i> , 1970, 47, 55-63. | 3.0 | 8 |