Julie Bernhardt

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3730437/publications.pdf

Version: 2024-02-01

262 papers

14,534 citations

54 h-index 24982 109 g-index

278 all docs

278 docs citations

times ranked

278

11904 citing authors

#	Article	IF	CITATIONS
1	Stroke rehabilitation. Lancet, The, 2011, 377, 1693-1702.	13.7	1,961
2	Physical Activity and Exercise Recommendations for Stroke Survivors. Stroke, 2014, 45, 2532-2553.	2.0	1,009
3	Agreed definitions and a shared vision for new standards in stroke recovery research: The Stroke Recovery and Rehabilitation Roundtable taskforce. International Journal of Stroke, 2017, 12, 444-450.	5.9	624
4	Inactive and Alone. Stroke, 2004, 35, 1005-1009.	2.0	524
5	Standardized measurement of sensorimotor recovery in stroke trials: Consensus-based core recommendations from the Stroke Recovery and Rehabilitation Roundtable. International Journal of Stroke, 2017, 12, 451-461.	5.9	352
6	A Very Early Rehabilitation Trial for Stroke (AVERT). Stroke, 2008, 39, 390-396.	2.0	328
7	Extending thrombolysis to 4·5–9 h and wake-up stroke using perfusion imaging: a systematic review and meta-analysis of individual patient data. Lancet, The, 2019, 394, 139-147.	13.7	321
8	Motivational interviewing to increase physical activity in people with chronic health conditions: a systematic review and meta-analysis. Clinical Rehabilitation, 2014, 28, 1159-1171.	2.2	292
9	Very Early Mobilization After Stroke Fast-Tracks Return to Walking. Stroke, 2011, 42, 153-158.	2.0	257
10	Agreed Definitions and a Shared Vision for New Standards in Stroke Recovery Research: The Stroke Recovery and Rehabilitation Roundtable Taskforce. Neurorehabilitation and Neural Repair, 2017, 31, 793-799.	2.9	225
11	Stroke Survivors' Experiences of Physical Rehabilitation: A Systematic Review of Qualitative Studies. Archives of Physical Medicine and Rehabilitation, 2015, 96, 1698-1708.e10.	0.9	214
12	How Physically Active Are People Following Stroke? Systematic Review and Quantitative Synthesis. Physical Therapy, 2017, 97, 707-717.	2.4	209
13	Balance and mobility outcomes for stroke patients: a comprehensive audit. Australian Journal of Physiotherapy, 1997, 43, 173-180.	0.9	194
14	Physical Activity and Sedentary Behaviors in People With Stroke Living in the Community: A Systematic Review. Physical Therapy, 2014, 94, 185-196.	2.4	192
15	Prespecified dose-response analysis for A Very Early Rehabilitation Trial (AVERT). Neurology, 2016, 86, 2138-2145.	1.1	170
16	An enriched environment increases activity in stroke patients undergoing rehabilitation in a mixed rehabilitation unit: a pilot non-randomized controlled trial. Disability and Rehabilitation, 2014, 36, 255-262.	1.8	163
17	An International Standard Set of Patient-Centered Outcome Measures After Stroke. Stroke, 2016, 47, 180-186.	2.0	161
18	Loss of Skeletal Muscle Mass after Stroke: a Systematic Review. International Journal of Stroke, 2010, 5, 395-402.	5.9	151

#	Article	IF	CITATIONS
19	Sitting and Activity Time in People With Stroke. Physical Therapy, 2016, 96, 193-201.	2.4	149
20	Montreal Cognitive Assessment and Mini-Mental State Examination are both valid cognitive tools in stroke. Acta Neurologica Scandinavica, 2013, 128, 122-129.	2.1	139
21	Moving rehabilitation research forward: Developing consensus statements for rehabilitation and recovery research. International Journal of Stroke, 2016, 11, 454-458.	5.9	137
22	Standardized Measurement of Sensorimotor Recovery in Stroke Trials: Consensus-Based Core Recommendations from the Stroke Recovery and Rehabilitation Roundtable. Neurorehabilitation and Neural Repair, 2017, 31, 784-792.	2.9	135
23	Little therapy, little physical activity: Rehabilitation within the first 14 days of organized stroke unit care. Acta Dermato-Venereologica, 2007, 39, 43-48.	1.3	134
24	Physical Activity in Hospitalised Stroke Patients. Stroke Research and Treatment, 2012, 2012, 1-13.	0.8	131
25	The effect of physical activity on cognitive function after stroke: a systematic review. International Psychogeriatrics, 2012, 24, 557-567.	1.0	129
26	Very Early Rehabilitation or Intensive Telemetry after Stroke: A Pilot Randomised Trial. Cerebrovascular Diseases, 2010, 29, 352-360.	1.7	124
27	Effects of Physical Activity on Poststroke Cognitive Function. Stroke, 2017, 48, 3093-3100.	2.0	118
28	Early rehabilitation after stroke. Current Opinion in Neurology, 2017, 30, 48-54.	3.6	117
29	The importance of cognition to quality of life after stroke. Journal of Psychosomatic Research, 2014, 77, 374-379.	2.6	116
30	Not All Stroke Units Are the Same. Stroke, 2008, 39, 2059-2065.	2.0	111
31	A Very Early Rehabilitation Trial after stroke (AVERT): a Phase III, multicentre, randomised controlled trial. Health Technology Assessment, 2017, 21, 1-120.	2.8	109
32	Family-led rehabilitation after stroke in India (ATTEND): a randomised controlled trial. Lancet, The, 2017, 390, 588-599.	13.7	108
33	An Enriched Environment Improves Sensorimotor Function Post-Ischemic Stroke. Neurorehabilitation and Neural Repair, 2010, 24, 802-813.	2.9	106
34	Early Mobilization After Stroke. Stroke, 2015, 46, 1141-1146.	2.0	95
35	Energy Expenditure and Cost During Walking After Stroke: A Systematic Review. Archives of Physical Medicine and Rehabilitation, 2016, 97, 619-632.e1.	0.9	93
36	Exploring the Role of Accelerometers in the Measurement of Real World Upper-Limb Use After Stroke. Brain Impairment, 2016, 17, 16-33.	0.7	90

#	Article	IF	CITATIONS
37	Motor Impairment and Recovery in the Upper Limb After Stroke. Stroke, 2005, 36, 625-629.	2.0	89
38	Evolution of Brain Activation with Good and Poor Motor Recovery after Stroke. Neurorehabilitation and Neural Repair, 2006, 20, 24-41.	2.9	89
39	The Montreal Cognitive Assessment. Stroke, 2011, 42, 2642-2644.	2.0	89
40	How is physical activity monitored in people following stroke?. Disability and Rehabilitation, 2015, 37, 1717-1731.	1.8	83
41	Clinical feasibility of interactive motion-controlled games for stroke rehabilitation. Journal of NeuroEngineering and Rehabilitation, 2015, 12, 63.	4.6	82
42	A Very Early Rehabilitation Trial (AVERT). International Journal of Stroke, 2006, 1, 169-171.	5.9	74
43	Carers' Experiences, Needs, and Preferences During Inpatient Stroke Rehabilitation: A Systematic Review of Qualitative Studies. Archives of Physical Medicine and Rehabilitation, 2017, 98, 1852-1862.e13.	0.9	72
44	Physical Activity Early after Stroke and Its Association to Functional Outcome 3ÂMonths Later. Journal of Stroke and Cerebrovascular Diseases, 2014, 23, e305-e312.	1.6	70
45	Physical, cognitive and social activity levels of stroke patients undergoing rehabilitation within a mixed rehabilitation unit. Clinical Rehabilitation, 2014, 28, 91-101.	2.2	66
46	Very Early Mobilisation and Complications in the First 3 Months after Stroke: Further Results from Phase II of A Very Early Rehabilitation Trial (AVERT). Cerebrovascular Diseases, 2009, 28, 378-383.	1.7	65
47	A stroke recovery trial development framework: Consensus-based core recommendations from the Second Stroke Recovery and Rehabilitation Roundtable. International Journal of Stroke, 2019, 14, 792-802.	5.9	64
48	Mobilisation â€~in Bed' Is Not Mobilisation. Cerebrovascular Diseases, 2007, 24, 157-158.	1.7	63
49	Early Mobilization After Stroke. Stroke, 2010, 41, 2632-2636.	2.0	63
50	Very early versus delayed mobilisation after stroke. The Cochrane Library, 2009, , CD006187.	2.8	62
51	Physical Therapists' Guideline Adherence on Early Mobilization and Intensity of Practice at Dutch Acute Stroke Units. Stroke, 2012, 43, 2395-2401.	2.0	61
52	The effect of very early mobilisation after stroke on psychological well-being. Journal of Rehabilitation Medicine, 2008, 40, 609-614.	1.1	60
53	Stroke rehabilitation in low-income and middle-income countries: a call to action. Lancet, The, 2020, 396, 1452-1462.	13.7	59
54	Sitting time and physical activity after stroke: physical ability is only part of the story. Topics in Stroke Rehabilitation, 2016, 23, 36-42.	1.9	58

#	Article	IF	CITATIONS
55	Economic Evaluation alongside a Phase II, Multi-Centre, Randomised Controlled Trial of Very Early Rehabilitation after Stroke (AVERT). Cerebrovascular Diseases, 2008, 26, 475-481.	1.7	57
56	Reducing Sitting Time After Stroke: A Phase II Safety and Feasibility Randomized Controlled Trial. Archives of Physical Medicine and Rehabilitation, 2016, 97, 273-280.	0.9	57
57	Predictors of Poststroke Mobility: Systematic Review. International Journal of Stroke, 2011, 6, 321-327.	5.9	56
58	Circuit Class Therapy or Seven-Day Week Therapy for Increasing Rehabilitation Intensity of Therapy after Stroke (CIRCIT): A Randomized Controlled Trial. International Journal of Stroke, 2015, 10, 594-602.	5.9	56
59	Accuracy of Observational Kinematic Assessment of Upper-Limb Movements. Physical Therapy, 1998, 78, 259-270.	2.4	53
60	How Well Do Standard Stroke Outcome Measures Reflect Quality of Life?. Stroke, 2013, 44, 3161-3165.	2.0	52
61	Strapping the hemiplegic shoulder prevents development of pain during rehabilitation: a randomized controlled trial. Clinical Rehabilitation, 2006, 20, 287-295.	2.2	51
62	The NIH Stroke Scale Can Establish Cognitive Function after Stroke. Cerebrovascular Diseases, 2010, 30, 7-14.	1.7	51
63	A randomized control trial of intensive aphasia therapy after acute stroke: The Very Early Rehabilitation for SpEech (VERSE) study. International Journal of Stroke, 2021, 16, 556-572.	5.9	51
64	Exercise Preferences Are Different after Stroke. Stroke Research and Treatment, 2012, 2012, 1-9.	0.8	49
65	Translating the Use of An Enriched Environment Poststroke from Bench to Bedside: Study Design and Protocol Used to Test the Feasibility of Environmental Enrichment on Stroke Patients in Rehabilitation. International Journal of Stroke, 2012, 7, 521-526.	5.9	49
66	Ultrasound Is a Reliable Measure of Muscle Thickness in Acute Stroke Patients, for Some, but Not All Anatomical Sites: A Study of the Intra-Rater Reliability of Muscle Thickness Measures in Acute Stroke Patients. Ultrasound in Medicine and Biology, 2012, 38, 368-376.	1.5	48
67	Very early versus delayed mobilisation after stroke. The Cochrane Library, 2018, 2018, CD006187.	2.8	48
68	Efficacy and Safety of Individualized Coaching After Stroke: the LAST Study (Life After Stroke). Stroke, 2018, 49, 426-432.	2.0	47
69	Changes in balance and locomotion measures during rehabilitation following stroke. Physiotherapy Research International, 1998, 3, 109-122.	1.5	46
70	Prospective observation of physical activity in critically ill patients who were intubated for more than 48 hours. Journal of Critical Care, 2015, 30, 658-663.	2.2	46
71	Sedentary Behaviour and Physical Activity of People with Stroke in Rehabilitation Hospitals. Stroke Research and Treatment, 2014, 2014, 1-7.	0.8	44
72	Setting the scene for the Second Stroke Recovery and Rehabilitation Roundtable. International Journal of Stroke, 2019, 14, 450-456.	5.9	44

#	Article	IF	Citations
73	An Early Mobilization Protocol Successfully Delivers More and Earlier Therapy to Acute Stroke Patients. Neurorehabilitation and Neural Repair, 2012, 26, 20-26.	2.9	43
74	Exercise Reduces Infarct Volume and Facilitates Neurobehavioral Recovery. Neurorehabilitation and Neural Repair, 2014, 28, 800-812.	2.9	43
75	Poststroke Physical Activity Levels No Higher in Rehabilitation than in the Acute Hospital. Journal of Stroke and Cerebrovascular Diseases, 2016, 25, 938-945.	1.6	43
76	Changes in physical activity and related functional and disability levels in the first six months after stroke: A longitudinal follow-up study. Journal of Rehabilitation Medicine, 2013, 45, 423-428.	1.1	42
77	Hospital Differences in Motor Activity Early after Stroke: A Comparison of 11 Norwegian Stroke Units. Journal of Stroke and Cerebrovascular Diseases, 2015, 24, 1333-1340.	1.6	41
78	"Ward talk― Nurses' interaction with people with and without aphasia in the very early period poststroke. Aphasiology, 2016, 30, 609-628.	2.2	41
79	Patients' age as a determinant of care received following acute stroke: A systematic review. BMC Health Services Research, 2011, 11, 161.	2.2	40
80	Stroke management: updated recommendations for treatment along the care continuum. Internal Medicine Journal, 2012, 42, 562-569.	0.8	40
81	Measuring Activity Levels at an Acute Stroke Ward: Comparing Observations to a Device. BioMed Research International, 2013, 2013, 1-8.	1.9	40
82	Moving Rehabilitation Research Forward: Developing Consensus Statements for Rehabilitation and Recovery Research. Neurorehabilitation and Neural Repair, 2017, 31, 694-698.	2.9	40
83	More Outcomes than Trials: A Call for Consistent Data Collection across Stroke Rehabilitation Trials. International Journal of Stroke, 2013, 8, 18-24.	5.9	39
84	Advancing Stroke Recovery Through Improved Articulation of Nonpharmacological Intervention Dose. Stroke, 2021, 52, 761-769.	2.0	39
85	Stroke Patients Do Not Need to be Inactive in the First Two-Weeks after Stroke: Results from a Stroke Unit Focused on Early Rehabilitation. International Journal of Stroke, 2012, 7, 25-31.	5.9	37
86	When Should Rehabilitation Begin after Stroke?. International Journal of Stroke, 2013, 8, 5-7.	5.9	37
87	Frequent, short bouts of light-intensity exercises while standing decreases systolic blood pressure: Breaking Up Sitting Time after Stroke (BUST-Stroke) trial. International Journal of Stroke, 2018, 13, 932-940.	5.9	37
88	A comparative study of patients' activities and interactions in a stroke unit before and after reconstructionâ€"The significance of the built environment. PLoS ONE, 2017, 12, e0177477.	2.5	37
89	Selection for inpatient rehabilitation after severe stroke: What factors influence rehabilitation assessor decision-making?. Journal of Rehabilitation Medicine, 2013, 45, 24-31.	1.1	36
90	Early mobilization and quality of life after stroke. Neurology, 2019, 93, e717-e728.	1.1	34

#	Article	IF	Citations
91	Cutting a long story short: Reaction times in acute stroke are associated with longer term cognitive outcomes. Journal of the Neurological Sciences, 2012, 322, 102-106.	0.6	33
92	Stepping towards Prevention of Bone Loss after Stroke: A Systematic Review of the Skeletal Effects of Physical Activity after Stroke. International Journal of Stroke, 2012, 7, 330-335.	5.9	33
93	Boredom in patients with acquired brain injuries during inpatient rehabilitation: a scoping review. Disability and Rehabilitation, 2018, 40, 2713-2722.	1.8	33
94	The Modified Rankin Scale in Acute Stroke Has Good Inter-Rater-Reliability but Questionable Validity. Cerebrovascular Diseases, 2010, 29, 188-193.	1.7	32
95	Quality of life: An important outcome measure in a trial of very early mobilisation after stroke. Disability and Rehabilitation, 2010, 32, 875-884.	1.8	32
96	A mapping study on physical activity in stroke rehabilitation: Establishing the baseline. Journal of Rehabilitation Medicine, 2013, 45, 997-1003.	1.1	32
97	Stroke Rehabilitation in China: A Systematic Review and Meta-Analysis. International Journal of Stroke, 2014, 9, 494-502.	5.9	32
98	Additional weekend therapy may reduce length of rehabilitation stay after stroke: a meta-analysis of individual patient data. Journal of Physiotherapy, 2016, 62, 124-129.	1.7	31
99	Physical Fitness Training after Stroke, Time to Implement what we Know: More Research is Needed. International Journal of Stroke, 2011, 6, 506-508.	5.9	30
100	A qualitative exploration of discharge destination as an outcome or a driver of acute stroke care. BMC Health Services Research, 2014, 14, 193.	2.2	30
101	Are we armed with the right data? Pooled individual data review of biomarkers in people with severe upper limb impairment after stroke. NeuroImage: Clinical, 2017, 13, 310-319.	2.7	30
102	Can the physical environment itself influence neurological patient activity?. Disability and Rehabilitation, 2019, 41, 1177-1189.	1.8	30
103	†Better Wear Out Sheets than Shoes': A Survey of 202 Stroke Professionals' Early Mobilisation Practices and Concerns. International Journal of Stroke, 2011, 6, 10-15.	5.9	29
104	Prevalence of fatigue in patients 3Âmonths after stroke and association with early motor activity: a prospective study comparing stroke patients with a matched general population cohort. BMC Neurology, 2015, 15, 181.	1.8	29
105	The Scandinavian Stroke Scale is equally as good as The National Institutes of Health Stroke Scale in identifying 3-month outcome. Journal of Rehabilitation Medicine, 2016, 48, 909-912.	1.1	29
106	Best practice guidelines for the measurement of physical activity levels in stroke survivors: a secondary analysis of an observational study. International Journal of Rehabilitation Research, 2018, 41, 14-19.	1.3	29
107	An Observational Study of Acute Stroke Care in Four Countries: The European Registers of Stroke Study. Cerebrovascular Diseases, 2009, 28, 171-176.	1.7	28
108	A Phase 1 Exercise Dose Escalation Study for Stroke Survivors with Impaired Walking. International Journal of Stroke, 2015, 10, 1051-1056.	5.9	28

#	Article	IF	Citations
109	Implementing a complex rehabilitation intervention in a stroke trial: a qualitative process evaluation of AVERT. BMC Medical Research Methodology, 2016, 16, 52.	3.1	28
110	Could upright posture be harmful in the early stages of stroke? – Author's reply. Lancet, The, 2015, 386, 1734-1735.	13.7	27
111	Behavioral Mapping of Patient Activity to Explore the Built Environment During Rehabilitation. Herd, 2018, 11, 109-123.	1.5	25
112	Systematic Review of Process Indicators: Including Early Rehabilitation Interventions Used to Measure Quality of Acute Stroke Care. International Journal of Stroke, 2009, 4, 72-80.	5.9	24
113	Enhancing physical activity in older adults receiving hospital based rehabilitation: a phase II feasibility study. BMC Geriatrics, 2012, 12, 26.	2.7	24
114	A Stroke Recovery Trial Development Framework: Consensus-Based Core Recommendations from the Second Stroke Recovery and Rehabilitation Roundtable. Neurorehabilitation and Neural Repair, 2019, 33, 959-969.	2.9	24
115	Exploring post acute rehabilitation service use and outcomes for working age stroke survivors (â‰ 8 5) Tj ETQq1 e035850.	1 0.78431 1.9	.4 rgBT /Ove 24
116	Factors associated with time to independent walking recovery post-stroke. Journal of Neurology, Neurosurgery and Psychiatry, 2021, 92, 702-708.	1.9	24
117	Altering the rehabilitation environment to improve stroke survivor activity: A Phase II trial. International Journal of Stroke, 2022, 17, 299-307.	5.9	24
118	Physical Activity Patterns of Acute Stroke Patients Managed in a Rehabilitation Focused Stroke Unit. BioMed Research International, 2013, 2013, 1-8.	1.9	22
119	Approaches to Economic Evaluations of Stroke Rehabilitation. International Journal of Stroke, 2014, 9, 88-100.	5.9	22
120	Circuit Class Therapy and 7-Day-Week Therapy Increase Physiotherapy Time, But Not Patient Activity. Stroke, 2014, 45, 3002-3007.	2.0	22
121	Statistical Analysis Plan (SAP) for a Very Early Rehabilitation Trial (AVERT): An International Trial to Determine the Efficacy and Safety of Commencing out of Bed Standing and Walking Training (Very) Tj ETQq1 1 C Stroke, 2015, 10, 23-24.).784314 ı 5.9	rgBT /Overlo
122	Developing the Stroke Exercise Preference Inventory (SEPI). PLoS ONE, 2016, 11, e0164120.	2.5	22
123	A randomized controlled trial of very early rehabilitation in speech after stroke. International Journal of Stroke, 2016, 11, 586-592.	5.9	22
124	The personal and social experiences of community-dwelling younger adults after stroke in Australia: a qualitative interview study. BMJ Open, 2018, 8, e023525.	1.9	22
125	Telerehabilitation: Has Its Time Come?. Stroke, 2021, 52, 2694-2696.	2.0	22
126	Built environments for inpatient stroke rehabilitation services and care: a systematic literature review. BMJ Open, 2021, 11, e050247.	1.9	22

#	Article	IF	Citations
127	Timing and Dose of Upper Limb Motor Intervention After Stroke: A Systematic Review. Stroke, 2021, 52, 3706-3717.	2.0	22
128	Changes in Activity Levels in the First Month after Stroke. Journal of Physical Therapy Science, 2013, 25, 599-604.	0.6	21
129	Advances in Stroke 2017. Stroke, 2018, 49, e174-e199.	2.0	21
130	Rationale for Intervention and Dose Is Lacking in Stroke Recovery Trials: A Systematic Review. Stroke Research and Treatment, 2018, 2018, 1-9.	0.8	21
131	A systematic review protocol of timing, efficacy and cost effectiveness of upper limb therapy for motor recovery post-stroke. Systematic Reviews, 2019, 8, 187.	5.3	21
132	Robotic-assisted training after stroke: RATULS advances science. Lancet, The, 2019, 394, 6-8.	13.7	21
133	Low gait speed is associated with low physical activity and high sedentary time following stroke. Disability and Rehabilitation, 2021, 43, 2001-2008.	1.8	21
134	Changes in Fat Mass in Stroke Survivors: A Systematic Review. International Journal of Stroke, 2012, 7, 491-498.	5.9	20
135	Exploring the Efficacy of Constraint in Animal Models of Stroke. Neurorehabilitation and Neural Repair, 2013, 27, 3-12.	2.9	20
136	Two Days of Measurement Provides Reliable Estimates of Physical Activity Poststroke: An Observational Study. Archives of Physical Medicine and Rehabilitation, 2019, 100, 883-890.	0.9	20
137	How can stroke care be improved for younger service users? A qualitative study on the unmet needs of younger adults in inpatient and outpatient stroke care in Australia. Disability and Rehabilitation, 2020, 42, 1697-1704.	1.8	20
138	Hemispatial Neglect and Rehabilitation in Acute Stroke. Archives of Physical Medicine and Rehabilitation, 2009, 90, 1931-1936.	0.9	19
139	Representation of People with Aphasia in Randomized Controlled Trials of Acute Stroke Interventions. International Journal of Stroke, 2014, 9, 174-182.	5.9	19
140	Stand up and be counted: measuring time spent upright after hip fracture and comparison with community dwelling older people. Physiotherapy, 2005, 91, 215-222.	0.4	18
141	Accessing inpatient rehabilitation after acute severe stroke. International Journal of Rehabilitation Research, 2012, 35, 323-329.	1.3	18
142	Upright activity within the first week after stroke is associated with better functional outcome and health-related quality of life: A Norwegian multi-site study. Journal of Rehabilitation Medicine, 2016, 48, 280-286.	1.1	18
143	Bringing the single versus multi-patient room debate to vulnerable patient populations: a systematic review of the impact of room types on hospitalized older people and people with neurological disorders. Intelligent Buildings International, 2020, 12, 180-198.	2.3	18
144	Adherence to physical activity and cardiovascular recommendations during the 2 years after stroke rehabilitation discharge. Annals of Physical and Rehabilitation Medicine, 2021, 64, 101455.	2.3	18

#	Article	IF	CITATIONS
145	AVERT2(a very early rehabilitation trial, a very effective reproductive trigger): retrospective observational analysis of the number of babies born to trial staff. BMJ, The, 2015, 351, h6432.	6.0	17
146	Exploring threats to generalisability in a large international rehabilitation trial (AVERT). BMJ Open, 2015, 5, e008378.	1.9	17
147	What is spasticity? The discussion continues. International Journal of Therapy and Rehabilitation, 2007, 14, 391-395.	0.3	16
148	Measuring the Quality of Dysphagia Management Practices following Stroke: A Systematic Review. International Journal of Stroke, 2010, 5, 466-476.	5.9	16
149	Does evidence really matter? Professionals' opinions on the practice of early mobilization after stroke. Journal of Multidisciplinary Healthcare, 2011, 4, 367.	2.7	16
150	Circuit Class or Seven-Day Therapy for Increasing Intensity of Rehabilitation after Stroke: Protocol of the CIRCIT Trial. International Journal of Stroke, 2011, 6, 560-565.	5.9	16
151	Breaking up sitting time after stroke (BUST-Stroke). International Journal of Stroke, 2017, 12, 425-429.	5.9	16
152	How to Address Physical Activity Participation After Stroke in Research and Clinical Practice. Stroke, 2021, 52, e274-e277.	2.0	16
153	Why hospital design matters: A narrative review of built environments research relevant to stroke care. International Journal of Stroke, 2022, 17, 370-377.	5.9	16
154	Treatment and Outcomes of Working Aged Adults with Stroke: Results from a National Prospective Registry. Neuroepidemiology, 2017, 49, 113-120.	2.3	15
155	A Framework for Designing Inpatient Stroke Rehabilitation Facilities: A New Approach Using Interdisciplinary Value-Focused Thinking. Herd, 2019, 12, 142-158.	1.5	15
156	Early mobilisation post-stroke: a systematic review and meta-analysis of individual participant data. Disability and Rehabilitation, 2022, 44, 1156-1163.	1.8	15
157	A 2-Year Longitudinal Study of Physical Activity and Cardiovascular Risk in Survivors of Stroke. Physical Therapy, 2021, 101, .	2.4	15
158	Very Early Versus Delayed Mobilization After Stroke. Stroke, 2009, 40, .	2.0	14
159	Demographic and stroke-related factors as predictors of quality of acute stroke care provided by allied health professionals. Journal of Multidisciplinary Healthcare, 2011, 4, 247.	2.7	14
160	Additional structured physical activity does not improve walking in older people (> 60 years) undergoing inpatient rehabilitation: a randomised trial. Journal of Physiotherapy, 2018, 64, 237-244.	1.7	14
161	Breaking up sitting time after stroke (BUST-stroke). International Journal of Stroke, 2018, 13, 921-931.	5.9	14
162	Stroke: Physical Fitness, Exercise, and Fatigue. Stroke Research and Treatment, 2012, 2012, 1-2.	0.8	13

#	Article	IF	CITATIONS
163	Early Mobilization After Stroke Is Not Associated With Cognitive Outcome. Stroke, 2018, 49, 2147-2154.	2.0	13
164	Improving life after stroke needs global efforts to implement evidence-based physical activity pathways. International Journal of Stroke, 2019, 14, 457-459.	5.9	13
165	Cerebral haemodynamics with head position changes post-ischemic stroke: A systematic review and meta-analysis. Journal of Cerebral Blood Flow and Metabolism, 2020, 40, 1917-1933.	4.3	13
166	Look closer: The multidimensional patterns of post-stroke burden behind the modified Rankin Scale. International Journal of Stroke, 2021, 16, 420-428.	5.9	13
167	International stroke genetics consortium recommendations for studies of genetics of stroke outcome and recovery. International Journal of Stroke, 2022, 17, 260-268.	5.9	13
168	Relationship between pre-stroke physical activity and symptoms of post-stroke anxiety and depression: An observational study. Journal of Rehabilitation Medicine, 2019, 51, 755-760.	1.1	12
169	Does experience predict observational kinematic assessment accuracy?. Physiotherapy Theory and Practice, 2002, 18, 141-149.	1.3	11
170	More options and better job security required in career paths of physiotherapist researchers: an observational study. Australian Journal of Physiotherapy, 2008, 54, 135-140.	0.9	11
171	Bed Rest or Mobilization after rt-PA? A Case-Crossover Study of Factors Influencing Clinical Decision Making in Stroke Services. International Journal of Stroke, 2013, 8, 172-179.	5.9	11
172	An opportunistic study evaluating pharmacokinetics of sildenafil for the treatment of pulmonary hypertension in infants. Journal of Perinatology, 2016, 36, 744-747.	2.0	11
173	The Energy Cost of Steady State Physical Activity in Acute Stroke. Journal of Stroke and Cerebrovascular Diseases, 2018, 27, 1047-1054.	1.6	11
174	Economic evaluation of a phase III international randomised controlled trial of very early mobilisation after stroke (AVERT). BMJ Open, 2019, 9, e026230.	1.9	11
175	Early Mobilization After Stroke: Do Clinical Practice Guidelines Support Clinicians' Decision-Making?. Frontiers in Neurology, 2021, 12, 606525.	2.4	11
176	Are Patients with Intracerebral Haemorrhage Disadvantaged in Hospitals?. International Journal of Stroke, 2014, 9, 437-442.	5.9	10
177	Mobilization after thrombolysis (rtPA) within 24Âhours of acute stroke: what factors influence inclusion of patients in A Very Early Rehabilitation Trial (AVERT)?. BMC Neurology, 2014, 14, 163.	1.8	10
178	Is early rehabilitation a myth? Physical inactivity in the first week after myocardial infarction and stroke. Disability and Rehabilitation, 2016, 38, 1493-1499.	1.8	10
179	Safety and efficacy of recovery-promoting drugs for motor function after stroke: A systematic review of randomized controlled trials. Journal of Rehabilitation Medicine, 2019, 51, 319-330.	1.1	10
180	Utility-weighted modified Rankin Scale: Still too crude to be a truly patient-centric primary outcome measure?. International Journal of Stroke, 2020, 15, 268-277.	5.9	10

#	Article	IF	CITATIONS
181	Secondary Prevention of Stroke: Study Protocol for a Telehealth-Delivered Physical Activity and Diet Pilot Randomized Trial (ENAbLE-Pilot). Cerebrovascular Diseases, 2021, 50, 605-611.	1.7	10
182	"Can you hear me now?―Video conference coping strategies and experience during COVID-19 and beyond. Work, 2021, 70, 723-732.	1.1	10
183	The economic and health burden of stroke among younger adults in Australia from a societal perspective. BMC Public Health, 2022, 22, 218.	2.9	10
184	Physical inactivity, depression and anxiety in acute stroke. International Journal of Therapy and Rehabilitation, 2013, 20, 289-293.	0.3	9
185	The elephant in the single room debate: keeping patients active. BMJ, The, 2013, 347, f6333-f6333.	6.0	9
186	Changes to Volumetric Bone Mineral Density and Bone Strength after Stroke: A Prospective Study. International Journal of Stroke, 2015, 10, 396-399.	5.9	9
187	Economic Evaluation Plan (EEP) for A Very Early Rehabilitation Trial (AVERT): An international trial to compare the costs and cost-effectiveness of commencing out of bed standing and walking training (very early mobilization) within 24 h of stroke onset with usual stroke unit care. International lournal of Stroke, 2016, 11, 492-494.	5.9	9
188	Early Mobilization after Stroke: Changes in Clinical Opinion Despite an Unchanging Evidence Base. Journal of Stroke and Cerebrovascular Diseases, 2017, 26, 1-6.	1.6	9
189	The Effect of Tone-Reducing Orthotic Devices on Soleus Muscle Reflex Excitability while Standing in Patients with Spasticity Following Stroke. Prosthetics and Orthotics International, 2010, 34, 46-57.	1.0	8
190	Early Physical Activity and Discharge Destination after Stroke: A Comparison of Acute and Comprehensive Stroke Unit Care. Rehabilitation Research and Practice, 2013, 2013, 1-9.	0.6	8
191	How do comprehensive and acute stroke units differ? A critical review. International Journal of Therapy and Rehabilitation, 2013, 20, 41-53.	0.3	8
192	Upright activity and higher motor function may preserve bone mineral density within 6Âmonths of stroke: a longitudinal study. Archives of Osteoporosis, 2018, 13, 5.	2.4	8
193	The AVERT MoCA Data: Scoring Reliability in a Large Multicenter Trial. Assessment, 2020, 27, 976-981.	3.1	8
194	Exploring colour in context using Virtual Reality: Does a room change how you feel?. Virtual Reality, 2021, 25, 631-645.	6.1	8
195	Young Stroke Survivors' Preferred Methods of Meeting Their Unique Needs. Neurology, 2021, 96, e1701-e1710.	1.1	8
196	We only treat what it occurs to us to assess: the importance of knowledge-based assessment. , 2005, , $15-48$.		8
197	Stroke survivors' perceptions of the factors that influence engagement in activity outside dedicated therapy sessions in a rehabilitation unit: A qualitative study. Clinical Rehabilitation, 2022, 36, 822-830.	2.2	8
198	Giant Steps for the Science of Stroke Rehabilitation. International Journal of Stroke, 2013, 8, 1-2.	5.9	7

#	Article	IF	CITATIONS
199	The potential health and economic impact of improving stroke care standards for Australia. International Journal of Stroke, 2017, 12, 875-885.	5.9	7
200	A mixed-methods study to explore opinions of research translation held by researchers working in a Centre of Research Excellence in Australia. BMJ Open, 2018, 8, e022357.	1.9	7
201	Safety of Performing a Graded Exercise Test Early after Stroke and Transient Ischemic Attack. PM and R, 2020, 12, 445-453.	1.6	7
202	Fatal and Nonfatal Events Within 14 days After Early, Intensive Mobilization Poststroke. Neurology, 2021, 96, .	1.1	7
203	Distance-based Classification and Regression Trees for the analysis of complex predictors in health and medical research. Statistical Methods in Medical Research, 2021, 30, 2085-2104.	1.5	7
204	Training novice clinicians improves observation accuracy of the upper extremity after stroke. Archives of Physical Medicine and Rehabilitation, 2001, 82, 1611-1618.	0.9	6
205	Evaluating the effects of increasing physical activity to optimize rehabilitation outcomes in hospitalized older adults (MOVE Trial): study protocol for a randomized controlled trial. Trials, 2015, 16, 13.	1.6	6
206	How to do health services research in stroke: A focus on performance measurement and quality improvement. International Journal of Stroke, 2018, 13, 166-174.	5.9	6
207	Characterising Arm Recovery in People with Severe Stroke (CARPSS): protocol for a 12-month observational study of clinical, neuroimaging and neurophysiological biomarkers. BMJ Open, 2018, 8, e026435.	1.9	6
208	Implementing a protocol for a research impact assessment of the Centre for Research Excellence in Stroke Rehabilitation and Brain Recovery. Health Research Policy and Systems, 2018, 16, 71.	2.8	6
209	Reduced bone formation markers, and altered trabecular and cortical bone mineral densities of non-paretic femurs observed in rats with ischemic stroke: A randomized controlled pilot study. PLoS ONE, 2017, 12, e0172889.	2.5	6
210	Factors associated with paid employment 12 months after stroke in A Very Early Rehabilitation Trial (AVERT). Annals of Physical and Rehabilitation Medicine, 2022, 65, 101565.	2.3	6
211	The Therapy â€~Pill': Achieving Treatment Dose Within a Rehabilitation Trial. Brain Impairment, 2008, 9, 191-197.	0.7	5
212	An Investigation of the Neurophysiologic Effect of Tone-Reducing AFOs on Reflex Excitability in Subjects with Spasticity Following Stroke while Standing. Prosthetics and Orthotics International, 2010, 34, 154-165.	1.0	5
213	Age and gender as predictors of allied health quality stroke care. Journal of Multidisciplinary Healthcare, 2011, 4, 239.	2.7	5
214	Interventions for the uptake of evidence-based recommendations in acute stroke settings. The Cochrane Library, 0 , , .	2.8	5
215	Statistical analysis plan (SAP) for the Very Early Rehabilitation in Speech (VERSE) after stroke trial: an international 3-arm clinical trial to determine the effectiveness of early, intensive, prescribed, direct aphasia therapy. International Journal of Stroke, 2018, 13, 863-880.	5.9	5
216	Validity of Multisensor Array for Measuring Energy Expenditure of an Activity Bout in Early Stroke Survivors. Stroke Research and Treatment, 2018, 2018, 1-8.	0.8	5

#	Article	IF	Citations
217	What's in a Building? A Descriptive Survey of Adult Inpatient Rehabilitation Facility Buildings in Victoria, Australia. Archives of Rehabilitation Research and Clinical Translation, 2020, 2, 100040.	0.9	5
218	Clustering clinical and health care processes using a novel measure of dissimilarity for variable-length sequences of ordinal states. Statistical Methods in Medical Research, 2020, 29, 3059-3075.	1.5	5
219	Using the Barthel Index and modified Rankin Scale as Outcome Measures for Stroke Rehabilitation Trials; A Comparison of Minimum Sample Size Requirements. Journal of Stroke and Cerebrovascular Diseases, 2022, 31, 106229.	1.6	5
220	Co-Designing a New Yoga-Based Mindfulness Intervention for Survivors of Stroke: A Formative Evaluation. Neurology International, 2022, 14, 1-10.	2.8	5
221	Gender and being born overseas influences the amount of acute stroke therapy. Journal of Rehabilitation Medicine, 2013, 45, 130-136.	1.1	4
222	Reducing sedentary time and fat mass may improve glucose tolerance and insulin sensitivity in adults surviving 6 months after stroke: A phase I pilot study. European Stroke Journal, 2017, 2, 144-153.	5.5	4
223	Prestroke physical activity to reduce stroke severity. Neurology, 2018, 91, 727-728.	1.1	4
224	Increased Relative Functional Gain and Improved Stroke Outcomes: A Linked Registry Study of the Impact of Rehabilitation. Journal of Stroke and Cerebrovascular Diseases, 2021, 30, 106015.	1.6	4
225	Acute Hospital Admission for Stroke Is Characterised by Inactivity. Stroke Research and Treatment, 2020, 2020, 1-8.	0.8	4
226	Impairments, and physical design and culture of a rehabilitation unit influence stroke survivor activity: qualitative analysis of rehabilitation staff perceptions. Disability and Rehabilitation, 2022, 44, 8436-8441.	1.8	4
227	Response to Letter by Freeman et al Regarding Article, "Very Early Mobilization After Stroke Fast-Tracks Return to Walking: Further Results From the Phase II AVERT Randomized Controlled Trialâ€. Stroke, 2011, 42, e585.	2.0	3
228	Carers' experiences, needs and preferences during inpatient stroke rehabilitation: a protocol for a systematic review of qualitative studies. Systematic Reviews, 2015, 4, 108.	5.3	3
229	Application of Theory in Studies of Healthcare Built Environment Research. Herd, 2020, 13, 154-170.	1.5	3
230	Factors associated with improved walking in older people during hospital rehabilitation: secondary analysis of a randomized controlled trial. BMC Geriatrics, 2021, 21, 90.	2.7	3
231	What is the relationship between physical activity and cardiovascular risk factors in stroke survivors post completion of rehabilitation? Protocol for a longitudinal study. BMJ Open, 2017, 7, e019193.	1.9	3
232	How Many Hours of Device Wear Time Are Required to Accurately Measure Physical Activity Post Stroke?. International Journal of Environmental Research and Public Health, 2022, 19, 1191.	2.6	3
233	Depression and a lack of socialization are associated with high levels of boredom during stroke rehabilitation: An exploratory study using a new conceptual framework. Neuropsychological Rehabilitation, 2023, 33, 497-527.	1.6	3
234	Types of physical activity performed pre and post stroke. Brazilian Journal of Physical Therapy, 2022, 26, 100412.	2.5	3

#	Article	IF	CITATIONS
235	Changing practice for acute hemiplegic shoulder care: A best practice model. International Journal of Therapy and Rehabilitation, 2007, 14, 266-273.	0.3	2
236	Breaking up sitting time after stroke – How much less sitting is needed to improve blood pressure after stroke (BUST-BP-Dose): Protocol for a dose-finding study. Contemporary Clinical Trials Communications, 2019, 13, 100310.	1.1	2
237	Commentary on Arias M & Description of acute stroke patients. ⟨i⟩Journal of Clinical Nursing⟨/i⟩ 16, 282–288. Journal of Clinical Nursing, 2008, 17, 1957-1958.	3.0	1
238	Physiotherapists should ensure their representation in measures of the quality of patient care. Journal of Physiotherapy, 2010, 56, 219-220.	1.7	1
239	Early Mobilization Testing in Patients With Acute Stroke. Chest, 2012, 141, 1641-1642.	0.8	1
240	Response to letter: And yet it moves – AVERT enlightens translations stroke research. International Journal of Stroke, 2017, 12, NP14-NP15.	5.9	1
241	Editorial. International Journal of Stroke, 2017, 12, 443-443.	5.9	1
242	Determining Maximal Tolerable Aerobic Training Intensity in the Acute Phase after Stroke: a Novel Dose Ranging Trial Protocol. Journal of Stroke and Cerebrovascular Diseases, 2020, 29, 105359.	1.6	1
243	Occlusive Disease and Upright Activity in Acute Ischemic Stroke. Journal of Stroke and Cerebrovascular Diseases, 2021, 30, 105604.	1.6	1
244	What Is Next After This Well-Conducted, but Neutral, Multisite Trial Testing Self-Rehabilitation Approaches?. Stroke, 2021, 52, 1948-1950.	2.0	1
245	Gender Equity in Leadership and Conferences of the Stroke Society of Australasia. Cerebrovascular Diseases, 2021, , 1-6.	1.7	1
246	Ambulatory activity in stroke survivors associated with functional outcome and quality of life: An observational cohort study. Annals of Physical and Rehabilitation Medicine, 2022, 65, 101540.	2.3	1
247	Stroke population–specific neuroanatomical CT-MRI brain atlas. Neuroradiology, 2022, , 1.	2.2	1
248	Abstract 153: Shorter, More Frequent Out of Bed Activity Very Early After Stroke (AVERT) is Associated With a Favorable Outcome at 3 Months. Stroke, 2016, 47, .	2.0	1
249	Abstract 76: Exploring Efficacy and Safety of Very Early Mobilization Within 24 Hours of Stroke Onset Versus Usual Stroke Unit Care (A Very Early Rehabilitation Trial, AVERT): Pre-specified Subgroup Analysis. Stroke, 2016, 47, .	2.0	1
250	Self-evaluation of personal needs by community-living young stroke survivors using an online English language questionnaire. Disability and Rehabilitation, 2023, 45, 1830-1835.	1.8	1
251	Advancing rehabilitation clinical trials design. International Journal of Therapy and Rehabilitation, 2007, 14, 52-52.	0.3	0
252	Harnessing experience-dependent plasticity for CNS repair and regeneration. Future Neurology, 2012, 7, 523-525.	0.5	0

#	Article	IF	CITATIONS
253	Exercise after stroke - introduction. World Stroke Academy, 2013, 1, 11-11.	0.1	O
254	Interdisciplinary Team Interactions in Stroke Units: Can Team Dynamics Influence Patient Outcomes from a Clinician's Perspective. International Journal of Physical Medicine & Rehabilitation, 2014, s3, .	0.5	0
255	Very early mobilisation within 24 hours of stroke results in a less favourable outcome at 3 months [Author's response]. Journal of Physiotherapy, 2015, 61, 220-221.	1.7	O
256	30â€Exploring opinions about research translation held by leading australian stroke researchers. , 2018, , .		0
257	Authors' response to Letter to the Editor: Divergence among researchers regarding the stratification of time after stroke is still a concern. International Journal of Stroke, 2018, 13, NP13-NP13.	5.9	O
258	Critically appraised paper: Combined aerobic and resistance training is superior to usual care for improving some aspects of mobility after stroke [commentary]. Journal of Physiotherapy, 2020, 66, 131.	1.7	0
259	Organization of Care., 2012,, 93-105.		O
260	Abstract TP438: Reaction Time In Acute Stroke Predicts Long-Term Quality Of Life. Stroke, 2013, 44, .	2.0	0
261	Abstract TMP27: How Generalizable are Rehabilitation Trials? The Results of a Large International Rehabilitation Trial (AVERT). Stroke, 2016, 47, .	2.0	0
262	Early mobilisation and rehabilitation in intensive care unitâ€"ready for implementation?. Annals of Translational Medicine, 2017, 5, 57-57.	1.7	0