

Masami Ishibashi

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3754911/publications.pdf>

Version: 2024-02-01

365
papers

10,684
citations

38742

50
h-index

76900

74
g-index

437
all docs

437
docs citations

437
times ranked

7197
citing authors

#	ARTICLE	IF	CITATIONS
1	Bioactive metabolites of symbiotic marine microorganisms. <i>Chemical Reviews</i> , 1993, 93, 1753-1769.	47.7	344
2	Scytophycins, cytotoxic and antimycotic agents from the cyanophyte <i>Scytonema pseudohofmanni</i> . <i>Journal of Organic Chemistry</i> , 1986, 51, 5300-5306.	3.2	210
3	Purealidins B and C, new bromotyrosine alkaloids from the okinawan marine sponge <i>psammaphysilla purea</i> . <i>Tetrahedron</i> , 1991, 47, 6617-6622.	1.9	193
4	Ircinalins A and B from the Okinawan marine sponge <i>Ircinia</i> sp.: plausible biogenetic precursors of manzamine alkaloids. <i>Journal of Organic Chemistry</i> , 1992, 57, 2480-2483.	3.2	161
5	Separation of <i>Leucas aspera</i> , a Medicinal Plant of Bangladesh, Guided by Prostaglandin Inhibitory and Antioxidant Activities. <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 595-598.	1.3	144
6	Manzacidins A-C, novel tetrahydropyrimidine alkaloids from the Okinawan marine sponge <i>Hymeniacidon</i> sp. <i>Journal of Organic Chemistry</i> , 1991, 56, 4574-4576.	3.2	136
7	Ageliferins, potent actomyosin ATPase activators from the Okinawan marine sponge sp. <i>Tetrahedron</i> , 1990, 46, 5579-5586.	1.9	122
8	Analgesic Components of <i>Saposhnikovia</i> Root (<i>Saposhnikovia divaricata</i>).. <i>Chemical and Pharmaceutical Bulletin</i> , 2001, 49, 154-160.	1.3	121
9	Naturally Occurring Small Molecule Inhibitors of Hedgehog/Gli Mediated Transcription. <i>ChemBioChem</i> , 2008, 9, 1082-1092.	2.6	121
10	Penaresidin A and B, two novel azetidone alkaloids with potent actomyosin ATPase-activating activity from the Okinawan marine sponge <i>Penares</i> sp.. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 1135-1137.	0.9	114
11	Prianosin A, a novel antileukemic alkaloid from the okinawan marine sponge <i>Prianos melanos</i> . <i>Tetrahedron Letters</i> , 1987, 28, 4939-4942.	1.4	102
12	Amphidinolide C: the first twenty-five membered macrocyclic lactone with potent antineoplastic activity from the cultured dinoflagellate <i>Amphidinium</i> sp. <i>Journal of the American Chemical Society</i> , 1988, 110, 490-494.	13.7	101
13	Amphidinolides: Unique Macrolides from Marine Dinoflagellates. <i>Heterocycles</i> , 1997, 44, 543.	0.7	95
14	Hyrminosins A and B, new indole alkaloids from the Okinawan marine sponge <i>Hyrminos erecta</i> . <i>Tetrahedron</i> , 1990, 46, 7699-7702.	1.9	88
15	Amphidinolides G and H: new potent cytotoxic macrolides from the cultured symbiotic dinoflagellate <i>Amphidinium</i> sp. <i>Journal of Organic Chemistry</i> , 1991, 56, 5221-5224.	3.2	87
16	Studies on search for bioactive natural products targeting TRAIL signaling leading to tumor cell apoptosis. <i>Medicinal Research Reviews</i> , 2008, 28, 688-714.	10.5	84
17	Amphidinolide-A, a novel antineoplastic macrolide from the marine dinoflagellate <i>Amphidinium</i> sp.. <i>Tetrahedron Letters</i> , 1986, 27, 5755-5758.	1.4	83
18	Catalytic Asymmetric Synthesis of Mixed 3,3'-Bisindoles and Their Evaluation as Wnt Signaling Inhibitors. <i>Angewandte Chemie - International Edition</i> , 2013, 52, 2486-2490.	13.8	81

#	ARTICLE	IF	CITATIONS
19	Six Immunosuppressive Features from an Ascomycete, <i>Zopfiella ion gicaudata</i> , Found in a Screening Study Monitored by Immunomodulatory Activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 1005-1008.	1.3	80
20	Pseudodistomins A and B, novel antineoplastic piperidine alkaloids with calmodulin antagonistic activity from the Okinawan tunicate <i>Pseudodistoma kanoko</i> . <i>Journal of Organic Chemistry</i> , 1987, 52, 450-453.	3.2	79
21	Izumiphenazines A-C: Isolation and Structure Elucidation of Phenazine Derivatives from <i>Streptomyces</i> sp. IFM 11204. <i>Journal of Natural Products</i> , 2010, 73, 1999-2002.	3.0	75
22	Immunomodulatory Constituents from an Ascomycete, <i>Chaetomium seminudum</i> . <i>Journal of Natural Products</i> , 2004, 67, 98-102.	3.0	72
23	Keramamides B, D, novel peptides from the Okinawan marine sponge <i>Theonella</i> sp. <i>Journal of the American Chemical Society</i> , 1991, 113, 7812-7813.	13.7	70
24	Luteophanol A, a New Polyhydroxyl Compound from Symbiotic Marine Dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of Organic Chemistry</i> , 1997, 62, 3820-3823.	3.2	70
25	Keramamide F, a new thiazole-containing peptide from the Okinawan marine sponge <i>Theonella</i> sp. <i>Journal of Organic Chemistry</i> , 1992, 57, 5540-5542.	3.2	69
26	Rigidin, a novel alkaloid with calmodulin antagonistic activity from the okinawan marine tunicate <i>Eudistoma cf. rigida</i> . <i>Tetrahedron Letters</i> , 1990, 31, 4617-4620.	1.4	68
27	Amphidinolide F, a new cytotoxic macrolide from the marine dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of Antibiotics</i> , 1991, 44, 1259-1261.	2.0	68
28	Taurospongins A, a Novel Acetylenic Fatty Acid Derivative Inhibiting DNA Polymerase β and HIV Reverse Transcriptase from Sponge <i>Hippospongiasp.</i> . <i>Journal of Organic Chemistry</i> , 1997, 62, 3831-3836.	3.2	67
29	Iejimalides C and D, new antineoplastic 24-membered macrolide sulfates from the okinawan marine tunicate <i>Eudistoma cf. rigida</i> . <i>Tetrahedron Letters</i> , 1991, 32, 797-798.	1.4	65
30	Convolutamides A-F, novel β -lactam alkaloids from the marine bryozoan <i>Amathia convoluta</i> . <i>Tetrahedron</i> , 1994, 50, 10201-10206.	1.9	65
31	Dihydroflavonol BB-1, an extract of natural plant <i>Blumea balsamifera</i> , abrogates TRAIL resistance in leukemia cells. <i>Blood</i> , 2006, 107, 679-688.	1.4	65
32	Lignan glycosides and flavonoids from <i>Saraca asoca</i> with antioxidant activity. <i>Journal of Natural Medicines</i> , 2007, 61, 480-482.	2.3	65
33	Prostaglandin inhibitory and antioxidant components of <i>Cistus laurifolius</i> , a Turkish medicinal plant. <i>Journal of Ethnopharmacology</i> , 2006, 108, 371-378.	4.1	63
34	Bioactive products from marine micro- and macro-organisms. <i>Pure and Applied Chemistry</i> , 1999, 71, 1123-1126.	1.9	62
35	Novel antibiotics, furaquinocins A and B. Taxonomy, fermentation, isolation and physico-chemical and biological characteristics.. <i>Journal of Antibiotics</i> , 1990, 43, 247-252.	2.0	61
36	Novel antibiotics, furaquinocins C, D, E, F, G and H.. <i>Journal of Antibiotics</i> , 1991, 44, 390-395.	2.0	60

#	ARTICLE	IF	CITATIONS
37	Biosynthesis of furaquinocins A and B. <i>Journal of Organic Chemistry</i> , 1990, 55, 1132-1133.	3.2	59
38	Luteophanols B and C, new polyhydroxyl compounds from marine dinoflagellate <i>Amphidinium</i> sp.. <i>Tetrahedron</i> , 1998, 54, 14455-14464.	1.9	59
39	Terpenoids and a Flavonoid Glycoside from <i>Acacia pennata</i> Leaves as Hedgehog/GLI-Mediated Transcriptional Inhibitors. <i>Journal of Natural Products</i> , 2010, 73, 995-997.	3.0	59
40	New Wnt/Catenin Signaling Inhibitors Isolated from <i>Eleutherine palmifolia</i> . <i>Chemistry - an Asian Journal</i> , 2009, 4, 540-547.	3.3	57
41	Stelliferins A-F, new antineoplastic isomalabaricane triterpenes from the Okinawan marine sponge <i>Jaspis stellifera</i> . <i>Tetrahedron</i> , 1991, 47, 2181-2194.	1.9	56
42	Laboratory Culture of the Myxomycetes: Formation of Fruiting Bodies of <i>Didymium bahiense</i> and Its Plasmodial Production of Makaluvamine A. <i>Journal of Natural Products</i> , 2001, 64, 108-110.	3.0	56
43	Hedgehog/GLI-mediated transcriptional inhibitors from <i>Zizyphus cambodiana</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2008, 16, 9420-9424.	3.0	56
44	Metachromins A and B, novel antineoplastic sesquiterpenoids from the Okinawan sponge <i>Hippospongia</i> cf. <i>metachromia</i> . <i>Journal of Organic Chemistry</i> , 1988, 53, 2855-2858.	3.2	55
45	Keramamide A, a novel peptide from the Okinawan marine sponge <i>Theonella</i> sp.. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 2609.	0.9	55
46	Agelasine G, a New Antileukemic Alkaloid from the Okinawan Marine Sponge <i>Agelas</i> sp.. <i>Chemical and Pharmaceutical Bulletin</i> , 1992, 40, 766-767.	1.3	55
47	A novel natural compound, a cycloanthranilylproline derivative (Fuligocandin B), sensitizes leukemia cells to apoptosis induced by tumor necrosis factor-related apoptosis-inducing ligand (TRAIL) through 15-deoxy- $\Delta^{12,14}$ prostaglandin J2 production. <i>Blood</i> , 2007, 110, 1664-1674.	1.4	55
48	Konbamide, a novel peptide with calmodulin antagonistic activity from the Okinawan marine sponge <i>Theonella</i> sp.. <i>Journal of the Chemical Society Chemical Communications</i> , 1991, , 1050-1052.	2.0	54
49	Amphidinolide E, a novel antileukemic 19-membered macrolide from the cultured symbiotic dinoflagellate <i>Amphidinium</i> sp. <i>Journal of Organic Chemistry</i> , 1990, 55, 3421-3423.	3.2	52
50	Cystodytins D-I, New Cytotoxic Tetracyclic Aromatic Alkaloids from the Okinawan Marine Tunicate <i>Cystodytes dellechiaiei</i> . <i>Journal of Natural Products</i> , 1991, 54, 1634-1638.	3.0	52
51	Symbioramide, a novel Ca ²⁺ -ATPase activator from the cultured dinoflagellate <i>Symbiodinium</i> sp.. <i>Experientia</i> , 1988, 44, 800-802.	1.2	51
52	Cytotoxic Macrolides from a Cultured Marine Dinoflagellate of the Genus <i>Amphidinium</i> . <i>Journal of Natural Products</i> , 1989, 52, 1036-1041.	3.0	50
53	Two New Tryptophan-Derived Alkaloids from the Okinawan Marine Sponge <i>Aplysina</i> Sp.. <i>Journal of Natural Products</i> , 1994, 57, 1008-1011.	3.0	50
54	Immunomodulatory Constituents from an Ascomycete, <i>Eupenicillium crustaceum</i> , and Revised Absolute Structure of Macrophorin D. <i>Journal of Natural Products</i> , 2001, 64, 1234-1237.	3.0	50

#	ARTICLE	IF	CITATIONS
55	Synthesis, determination of stereochemistry, and evaluation of new bisindole alkaloids from the myxomycete <i>Arcyria ferruginea</i> : An approach for Wnt signal inhibitor. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2007, 17, 4254-4257.	2.2	50
56	Revised structures of prianosins C and D, antineoplastic alkaloids from the Okinawan marine sponge <i>Prianos melanos</i> . <i>Tetrahedron Letters</i> , 1991, 32, 1227-1228.	1.4	49
57	Pseudodistomin C, a new piperidine alkaloid with unusual absolute configuration from the Okinawan tunicate <i>Pseudodistoma kanoko</i> . <i>Journal of Organic Chemistry</i> , 1995, 60, 6941-6945.	3.2	49
58	Flavonoids from <i>Sonneratia caseolaris</i> . <i>Journal of Natural Medicines</i> , 2006, 60, 264-265.	2.3	49
59	Manzamenones A-F from the Okinawan marine sponges <i>Plakortis</i> Sp.: novel dimeric fatty acid derivatives possessing a bicyclo[4.3.0]nonane skeleton. <i>Journal of Organic Chemistry</i> , 1992, 57, 5255-5260.	3.2	48
60	Theonezolid a: a novel polyketide natural product from the Okinawan marine sponge <i>Theonella</i> sp. <i>Journal of the American Chemical Society</i> , 1993, 115, 6661-6665.	13.7	47
61	Sterol and Triterpenoid Constituents of <i>Verbenalittoralis</i> with NGF-Potentiating Activity. <i>Journal of Natural Products</i> , 2003, 66, 696-698.	3.0	47
62	New chlorogenin hexasaccharide isolated from <i>Agave fourcroydes</i> with cytotoxic and cell cycle inhibitory activities. <i>Bioorganic and Medicinal Chemistry</i> , 2004, 12, 3841-3845.	3.0	47
63	Collagenase Inhibitory Quinic Acid Esters from <i>Pomoea caprae</i> . <i>Journal of Natural Products</i> , 2005, 68, 794-796.	3.0	46
64	Sphingosine-related Marine Alkaloids: Cyclic Amino Alcohols. <i>Heterocycles</i> , 1996, 42, 943.	0.7	46
65	Amphidinolide-B, a novel macrolide with potent antineoplastic activity from the marine dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of the Chemical Society Chemical Communications</i> , 1987, , 1127.	2.0	45
66	Littoralisone, a Novel Neuritogenic Iridolactone Having an Unprecedented Heptacyclic Skeleton Including Four- and Nine-Membered Rings Consisting of Glucose from <i>Verbena littoralis</i> . <i>Journal of Organic Chemistry</i> , 2001, 66, 2165-2167.	3.2	45
67	Isolation of new carbamate- or pyridine-containing natural products, fuzanins A, B, C, and D from <i>Kitasatospora</i> sp. IFM10917. <i>Tetrahedron</i> , 2009, 65, 369-373.	1.9	45
68	Luffariolides A-E, new cytotoxic sesterterpenes from the Okinawan marine sponge <i>Luffariella</i> sp. <i>Journal of Organic Chemistry</i> , 1992, 57, 3503-3507.	3.2	44
69	Untenone A, a new cyclopentenone from the Okinawan marine sponge <i>Plakortis</i> sp. corresponding to the dienophile in the biosynthesis of manzamenones. <i>Tetrahedron Letters</i> , 1993, 34, 3749-3750.	1.4	44
70	Absolute stereochemistry of penaresidins A and B. <i>Tetrahedron Letters</i> , 1996, 37, 6775-6776.	1.4	44
71	Kehokorins A-C, novel cytotoxic dibenzofurans isolated from the myxomycete <i>Trichia favoginea</i> var. <i>persimilis</i> . <i>Tetrahedron Letters</i> , 2006, 47, 1505-1508.	1.4	44
72	Calotropin: A Cardenolide from <i>Calotropis gigantea</i> that Inhibits Wnt Signaling by Increasing Casein Kinase 1 β in Colon Cancer Cells. <i>ChemBioChem</i> , 2014, 15, 872-878.	2.6	44

#	ARTICLE	IF	CITATIONS
73	Niphatesines Aâ€“D, new antineoplastic pyridine alkaloids from the okinawan marine sponge Niphates sp.. Journal of the Chemical Society Perkin Transactions 1, 1990, , 3301-3303.	0.9	43
74	Amphidinolides O and P, novel 15-membered macrolides from the dinoflagellate Amphidinium sp.: Analysis of the relative stereochemistry and stable solution conformation. Journal of Organic Chemistry, 1995, 60, 6062-6066.	3.2	43
75	Dendocarbins Aâ€“N, New Drimane Sesquiterpenes from the Nudibranch Dendrodoris carbunculosa. Journal of Natural Products, 2001, 64, 726-731.	3.0	43
76	Eudistomidins E and F, new $\hat{1}^2$ -carboline alkaloids from the okinawan marine tunicate Eudistoma glaucus. Tetrahedron Letters, 1991, 32, 3539-3542.	1.4	42
77	Absolute stereochemistry of amphidinolide B. Tetrahedron Letters, 1994, 35, 8241-8242.	1.4	42
78	Colopsinol A, a Novel Polyhydroxyl Metabolite from Marine Dinoflagellate Amphidinium sp.. Journal of Organic Chemistry, 1999, 64, 1478-1482.	3.2	41
79	Five New Chromones Possessing Monoamine Oxidase Inhibitor Activity from an Ascomycete, Chaetomium quadrangulatum.. Chemical and Pharmaceutical Bulletin, 2002, 50, 330-336.	1.3	41
80	New Onoceranoid Triterpene Constituents from Lansium domesticum. Journal of Natural Products, 2002, 65, 1709-1711.	3.0	41
81	Isolation of bisindole alkaloids that inhibit the cell cycle from Myxomycetes Arcyria ferruginea and Tubifera casparyi. Bioorganic and Medicinal Chemistry Letters, 2003, 13, 2879-2881.	2.2	41
82	Sesquiterpenoids and Plasmin-Inhibitory Flavonoids from Blumea balsamifera. Journal of Natural Products, 2005, 68, 447-449.	3.0	41
83	New Hedgehog/GLI signaling inhibitors from Excoecaria agallocha. Bioorganic and Medicinal Chemistry Letters, 2011, 21, 718-722.	2.2	41
84	Structures of New Quassinoid Glycosides, Yadanzioids A, B, C, D, E, G, H, and New Quassinoids, Dehydrobrusatol and Dehydrobruceantinol from Brucea javanica(L.) MERR. Bulletin of the Chemical Society of Japan, 1985, 58, 2680-2686.	3.2	39
85	Niphatesines Eâ€“H, new pyridine alkaloids from the Okinawan marine sponge Niphates sp.. Journal of the Chemical Society Perkin Transactions 1, 1992, , 1291-1294.	0.9	39
86	Amphidinolide J: A cytotoxic macrolide from the marine dinoflagellate Amphidinium sp. Determination of the absolute stereochemistry. Journal of Organic Chemistry, 1993, 58, 2645-2646.	3.2	39
87	Nine Constituents Including Six Xanthone-Related Compounds Isolated from Two Ascomycetes, Gelasinospora santi-florii and Emericella quadrilineata, Found in a Screening Study Focused on Immunomodulatory Activity. Chemical and Pharmaceutical Bulletin, 2006, 54, 550-553.	1.3	39
88	A novel sesquiterpenoid dimer parviflorene F induces apoptosis by up-regulating the expression of TRAIL-R2 and a caspase-dependent mechanism. Bioorganic and Medicinal Chemistry, 2008, 16, 1756-1763.	3.0	39
89	Cycloartane Triterpenes Isolated from <i>Combretum quadrangulare</i> in a Screening Program for Death-Receptor Expression Enhancing Activity. Journal of Natural Products, 2011, 74, 249-255.	3.0	39
90	Constituents of the Root Bark of Ailanthus altissima SWINGLE. Isolation and X-Ray Crystal Structures of Shinjudilactone and Shinjulactone C and Conversion of Ailanthone into Shinjudilactone. Bulletin of the Chemical Society of Japan, 1983, 56, 3683-3693.	3.2	38

#	ARTICLE	IF	CITATIONS
91	Purealidin A, a new cytotoxic bromotyrosine-derived alkaloid from the Okinawan marine sponge <i>Psammaplysilla porea</i> . <i>Experientia</i> , 1991, 47, 299-300.	1.2	38
92	Cycloanthranilylproline-Derived Constituents from a Myxomycete <i>Fuligo candida</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2004, 52, 368-370.	1.3	38
93	Brandisianins A-F, Isoflavonoids Isolated from <i>Millettia brandisiana</i> in a Screening Program for Death-Receptor Expression Enhancement Activity. <i>Journal of Natural Products</i> , 2007, 70, 1910-1914.	3.0	38
94	9-Hydroxycanthin-6-one, a β -Carboline Alkaloid from <i>Eurycoma longifolia</i> , Is the First Wnt Signal Inhibitor through Activation of Glycogen Synthase Kinase 3 β without Depending on Casein Kinase 1 α . <i>Journal of Natural Products</i> , 2015, 78, 1139-1146.	3.0	38
95	New congeners of swinholides from the okinawan marine sponge <i>Theonella</i> sp.. <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 3185.	0.9	37
96	Diterpenes from <i>Leucasaspera</i> Inhibiting Prostaglandin-Induced Contractions. <i>Journal of Natural Products</i> , 2006, 69, 988-994.	3.0	37
97	Studies on the macrolides from marine dinoflagellate <i>Amphidinium</i> sp.: Structures of amphidinolides R and S and a succinate feeding experiment. <i>Tetrahedron</i> , 1997, 53, 7827-7832.	1.9	36
98	A new lignan from <i>Aphanamixis polystachya</i> . <i>Journal of Natural Medicines</i> , 2006, 60, 258-260.	2.3	36
99	Constituents of Seeds of <i>Brucea javanica</i> . Structures of New Bitter Principles, Yadanziolides A, B, C, Yadanziolides F, I, J, and L.. <i>Bulletin of the Chemical Society of Japan</i> , 1985, 58, 2673-2679.	3.2	35
100	Amphidinolide N, a novel 26-membered macrolide with remarkably potent cytotoxicity from the cultured marine dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of the Chemical Society Chemical Communications</i> , 1994, , 1455.	2.0	35
101	Constituents of <i>Amoora cucullata</i> with TRAIL resistance-overcoming activity. <i>Organic and Biomolecular Chemistry</i> , 2010, 8, 3696.	2.8	35
102	Tyrosine Derivatives Isolated from <i>Streptomyces</i> sp. IFM 10937 in a Screening Program for TRAIL-Resistance-Overcoming Activity. <i>Journal of Natural Products</i> , 2008, 71, 1963-1966.	3.0	34
103	Structures of novel antibiotics, furaquinocins A and B. <i>Tetrahedron Letters</i> , 1989, 30, 7427-7430.	1.4	33
104	Plakoridine A, a new tyramine-containing pyrrolidine alkaloid from the Okinawan marine sponge <i>Plakortis</i> sp.. <i>Journal of Organic Chemistry</i> , 1994, 59, 3712-3713.	3.2	33
105	Immunomodulatory Constituents from an Ascomycete, <i>Emericella aurantio-brunnea</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 2000, 48, 1436-1441.	1.3	33
106	New hedgehog/GLI-signaling inhibitors from <i>Adenium obesum</i> . <i>Organic and Biomolecular Chemistry</i> , 2011, 9, 1133-1139.	2.8	33
107	Natural compounds with Wnt signal modulating activity. <i>Natural Product Reports</i> , 2015, 32, 1622-1628.	10.3	33
108	Shimofuridin A, a nucleoside derivative embracing an acylfucopyranoside unit isolated from the Okinawan marine tunicate <i>Aplidium multiplicatum</i> . <i>Journal of Organic Chemistry</i> , 1994, 59, 255-257.	3.2	32

#	ARTICLE	IF	CITATIONS
109	Amphidinolide Q, a novel 12-membered macrolide from the cultured marine dinoflagellate <i>Amphidinium</i> sp. <i>Tetrahedron Letters</i> , 1996, 37, 1449-1450.	1.4	32
110	New cytotoxic bisindole alkaloids with protein tyrosine kinase inhibitory activity from a myxomycete <i>Lycogala epidendrum</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2005, 15, 2776-2780.	2.2	32
111	Death receptor 5 promoter-enhancing compounds isolated from <i>Catimbiium speciosum</i> and their enhancement effect on TRAIL-induced apoptosis. <i>Bioorganic and Medicinal Chemistry</i> , 2009, 17, 6748-6754.	3.0	32
112	Isolation and structure elucidation of izuminosides A-C: a rare phenazine glycosides from <i>Streptomyces</i> sp. IFM 11260. <i>Journal of Antibiotics</i> , 2011, 64, 271-275.	2.0	32
113	Cycloartane Triterpenes and Ingot Diterpenes Isolated from <i>Euphorbia neriifolia</i> in a Screening Program for Death-Receptor Expression-Enhancing Activity. <i>Planta Medica</i> , 2012, 78, 1370-1377.	1.3	32
114	Xylogranin B: A Potent Wnt Signal Inhibitory Limonoid from <i>Xylocarpus granatum</i> . <i>Organic Letters</i> , 2013, 15, 6106-6109.	4.6	32
115	9-Methyl-7-bromoedistomin D, a potent inducer of calcium release from sarcoplasmic reticulum of skeletal muscle. <i>Experientia</i> , 1989, 45, 782-783.	1.2	31
116	Three new oxylipins related to 3,6-dioxo-4-docosenoic acid from Okinawan marine sponges, <i>Plakortis</i> spp. <i>Tetrahedron</i> , 1995, 51, 5979-5986.	1.9	31
117	Studies on the Stereochemistry of Amphidinolides: Synthesis of a Diastereomer of the C-1-C-9 Fragment of Amphidinolide C.. <i>Chemical and Pharmaceutical Bulletin</i> , 1996, 44, 1819-1822.	1.3	31
118	Parviflorenes B-F, novel cytotoxic unsymmetrical sesquiterpene-dimers with three backbone skeletons from <i>Curcuma parviflora</i> . <i>Tetrahedron</i> , 2004, 60, 10817-10824.	1.9	31
119	Cryptolepine, isolated from <i>Sida acuta</i> , sensitizes human gastric adenocarcinoma cells to TRAIL-induced apoptosis. <i>Phytotherapy Research</i> , 2011, 25, 147-150.	5.8	31
120	Yoropyrazone, a new naphthopyridazone alkaloid isolated from <i>Streptomyces</i> sp. IFM 11307 and evaluation of its TRAIL resistance-overcoming activity. <i>Journal of Antibiotics</i> , 2012, 65, 245-248.	2.0	31
121	Structure Determination of Bitter Principles in <i>Ailanthus altissima</i> . Structure of Shinjulactone A and Revised Structure of Ailanthone. <i>Bulletin of the Chemical Society of Japan</i> , 1983, 56, 3694-3698.	3.2	30
122	Theoneberine: the first brominated benzyltetrahydroprotoberberine alkaloid from the Okinawan marine sponge <i>Theonella</i> sp. <i>Journal of Organic Chemistry</i> , 1992, 57, 6680-6682.	3.2	30
123	Purealidin D, a new pyridine alkaloid from the okinawan marine sponge <i>psammaplysilla pura</i> . <i>Tetrahedron Letters</i> , 1992, 33, 2597-2598.	1.4	30
124	Two new polycyclic aromatic alkaloids from the Okinawan marine sponge <i>Biemna</i> sp.. <i>Tetrahedron</i> , 1993, 49, 8337-8342.	1.9	30
125	New pyranonaphthoquinones and a phenazine alkaloid isolated from <i>Streptomyces</i> sp. IFM 11307 with TRAIL resistance-overcoming activity. <i>Journal of Antibiotics</i> , 2011, 64, 729-734.	2.0	30
126	New Pentacyclic Compounds from the Okinawan Marine Sponge <i>Xestospongia sapra</i> . <i>Journal of Natural Products</i> , 1992, 55, 994-998.	3.0	29

#	ARTICLE	IF	CITATIONS
127	Pseudodistomin B: Revised structure and first total synthesis. <i>Tetrahedron Letters</i> , 1992, 33, 7389-7390.	1.4	29
128	Amphidinin A, a novel amphidinolide-related metabolite from the cultured marine dinoflagellate <i>Amphidinium</i> sp.. <i>Tetrahedron Letters</i> , 1994, 35, 7049-7050.	1.4	29
129	Littorachalcone, a New Enhancer of NGF-Mediated Neurite Outgrowth, from <i>Verbena littoralis</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 872-874.	1.3	29
130	Katorazone, a new yellow pigment with a 2-azaquinone-phenylhydrazone structure produced by <i>Streptomyces</i> sp. IFM 11299. <i>Tetrahedron Letters</i> , 2012, 53, 3346-3348.	1.4	29
131	Amphidinolide K, a new 19-membered macrolide from the cultured symbiotic dinoflagellate <i>Amphidinium</i> sp. <i>Journal of Organic Chemistry</i> , 1993, 58, 6928-6929.	3.2	28
132	Isolation of Apo-9'-fucoxanthinone from the Cultured Marine Dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of Natural Products</i> , 1995, 58, 1097-1099.	3.0	28
133	Total synthesis of pseudodistomin C, a sphingosine-related piperidine alkaloid from tunicate <i>Pseudodistoma kanoko</i> . <i>Tetrahedron</i> , 1996, 52, 4573-4580.	1.9	28
134	Prenylated Flavonoids and Resveratrol Derivatives Isolated from <i>Artocarpus communis</i> with the Ability to Overcome TRAIL Resistance. <i>Journal of Natural Products</i> , 2015, 78, 103-110.	3.0	28
135	Anti-proliferative effect of <i>Moringa oleifera</i> Lam (Moringaceae) leaf extract on human colon cancer HCT116 cell line. <i>Tropical Journal of Pharmaceutical Research</i> , 2017, 16, 371.	0.3	28
136	Plakotenin, a new cytotoxic carboxylic acid from the okinawan marine sponge <i>plakortis</i> Sp.. <i>Tetrahedron Letters</i> , 1992, 33, 2579-2580.	1.4	27
137	Isolation and structures of Theonezolides B and C from the Okinawan marine sponge <i>Theonella</i> sp.. <i>Tetrahedron</i> , 1994, 50, 8355-8362.	1.9	27
138	Revised Structure of Pseudodistomin A, a Piperidine Alkaloid Isolated from the Okinawan Tunicate <i>Pseudodistoma kanoko</i> . <i>Journal of Natural Products</i> , 1995, 58, 804-806.	3.0	27
139	Biosynthetic studies of amphidinolide J: explanation of the generation of the unusual odd-numbered macrocyclic lactone. <i>Journal of the Chemical Society Chemical Communications</i> , 1995, , 1639.	2.0	27
140	Topsentinols A-J, New Sterols with Highly Branched Side Chains from Marine Sponge <i>Topsentia</i> sp.. <i>Chemical and Pharmaceutical Bulletin</i> , 1997, 45, 1435-1438.	1.3	27
141	Marine Natural Products and Marine Chemical Ecology. , 1999, , 415-649.		27
142	Prenylated Xanthenes from <i>Garcinia xanthochymus</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 1332-1334.	1.3	27
143	Bisindole Alkaloids from <i>Myxomycetes Arcyriadenudata</i> and <i>Arcyria obvelata</i> . <i>Journal of Natural Products</i> , 2006, 69, 1252-1254.	3.0	27
144	Physalin H from <i>Solanum nigrum</i> as an Hh signaling inhibitor blocks GLI1-DNA-complex formation. <i>Beilstein Journal of Organic Chemistry</i> , 2014, 10, 134-140.	2.2	27

#	ARTICLE	IF	CITATIONS
145	1H- and 13C-nmr Spectral Investigation on Amphidinolide A, an Antileukemic Marine Macrolide. Journal of Natural Products, 1991, 54, 1435-1439.	3.0	26
146	Keruffaride: Structure Revision and Isolation from Plural Genera of Okinawan Marine Sponges. Journal of Natural Products, 1993, 56, 1856-1860.	3.0	26
147	Luffariolides F and G, New Manoalide Derivatives from the Okinawan Marine Sponge Luffariella Sp.. Journal of Natural Products, 1993, 56, 436-439.	3.0	26
148	A New Iridoid Glycoside with Nerve Growth Factor-Potentiating Activity, Gelsemiol 6'-trans-Caffeoyl-1-glucoside, from Verbena littoralis. Chemical and Pharmaceutical Bulletin, 2003, 51, 1103-1105.	1.3	26
149	Six New Constituents from an Ascomycete, Chaetomium quadrangulatum, Found in a Screening Study Focused on Monoamine Oxidase Inhibitory Activity.. Chemical and Pharmaceutical Bulletin, 2003, 51, 247-251.	1.3	26
150	A Bisindole Alkaloid with Hedgehog Signal Inhibitory Activity from the Myxomycete Perichaena chrysosperma. Journal of Natural Products, 2010, 73, 1711-1713.	3.0	26
151	Constituents of Seeds of Ailanthus altissima SWINGLE. Isolation and Structures of Shinjuglycosides A, B, C, and D. Bulletin of the Chemical Society of Japan, 1984, 57, 2496-2501.	3.2	25
152	Structure Determination of Bitter Principles of Ailanthus altissima. Structures of Shinjulactones B, D, and E. Bulletin of the Chemical Society of Japan, 1984, 57, 2484-2489.	3.2	25
153	Lindbladione and Related Naphthoquinone Pigments from a Myxomycete Lindbladia tubulina. Chemical and Pharmaceutical Bulletin, 2002, 50, 1126-1127.	1.3	25
154	Cribrarione A, a new antimicrobial naphthoquinone pigment from a myxomycete Cribraria purpurea. Tetrahedron, 2003, 59, 3433-3435.	1.9	25
155	Melleumin A, a novel peptide lactone isolated from the cultured myxomycete Physarum melleum. Tetrahedron Letters, 2005, 46, 267-271.	1.4	25
156	Cassaine diterpenoid dimers isolated from Erythropheum succirubrum with TRAIL-resistance overcoming activity. Tetrahedron Letters, 2009, 50, 4658-4662.	1.4	25
157	Ricinine: A pyridone alkaloid from Ricinus communis that activates the Wnt signaling pathway through casein kinase 1 α . Bioorganic and Medicinal Chemistry, 2014, 22, 4597-4601.	3.0	25
158	Hes1 inhibitor isolated by target protein oriented natural products isolation (TPO-NAPI) of differentiation activators of neural stem cells. Chemical Science, 2016, 7, 1514-1520.	7.4	25
159	Hymenosulphate, a novel sterol sulphate with Ca-releasing activity from the cultured marine haptophyte Hymenomonas sp. Journal of the Chemical Society Perkin Transactions 1, 1989, , 101.	0.9	24
160	Colopsinols B and C, new long chain polyhydroxy compounds from cultured marine dinoflagellate Amphidinium sp.. Journal of the Chemical Society Perkin Transactions 1, 1999, , 3483-3487.	0.9	24
161	Steroidal saponins from Calamus insignis, and their cell growth and cell cycle inhibitory activities. Bioorganic and Medicinal Chemistry, 2006, 14, 659-665.	3.0	24
162	Secoiridoid components from Jasminum grandiflorum. Phytochemistry, 2007, 68, 1718-1721.	2.9	24

#	ARTICLE	IF	CITATIONS
163	Boesenberols, Pimarane Diterpenes with TRAIL-Resistance-Overcoming Activity from <i>Boesenbergia pandurata</i> . <i>Journal of Natural Products</i> , 2016, 79, 2075-2082.	3.0	24
164	Bio-active Natural Products with TRAIL-Resistance Overcoming Activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2016, 64, 119-127.	1.3	24
165	Nabscessins A and B, Aminocyclitol Derivatives from <i>Nocardia abscessus</i> IFM 10029 ^T . <i>Journal of Natural Products</i> , 2017, 80, 565-568.	3.0	24
166	Purealidins E-G, New Bromotyrosine Alkaloids from the Okinawan Marine Sponge <i>Psammaplysilla pura</i> . <i>Journal of Natural Products</i> , 1992, 55, 1325-1327.	3.0	23
167	Isolation and structure of shimofuridins B ¹ / ₄ G from the Okinawan marine tunicate <i>Aplidium multiplicatum</i> . <i>Tetrahedron</i> , 1994, 50, 8651-8656.	1.9	23
168	Cribrarione B, a New Naphthoquinone Pigment from the Myxomycete <i>Cribraria cancellata</i> . <i>Journal of Natural Products</i> , 2003, 66, 1611-1612.	3.0	23
169	New Bisindole Alkaloids Isolated from Myxomycetes <i>Arcyria cinerea</i> and <i>Lycogala epidendrum</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 594-597.	1.3	23
170	First isolation of sesquiterpenes and flavonoids from <i>Zingiber spectabile</i> and identification of zerumbone as the major cell growth inhibitory component. <i>Natural Product Research</i> , 2007, 21, 1242-1247.	1.8	23
171	The first Hes1 dimer inhibitors from natural products. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2009, 19, 5778-5781.	2.2	23
172	2-Methoxy-1,4-naphthoquinone isolated from <i>Impatiens balsamina</i> in a screening program for activity to inhibit Wnt signaling. <i>Journal of Natural Medicines</i> , 2011, 65, 234-236.	2.3	23
173	Hh signaling inhibitors from <i>Vitex negundo</i> ; naturally occurring inhibitors of the GLI1-DNA complex. <i>Molecular BioSystems</i> , 2013, 9, 1012.	2.9	23
174	Bioactive Secondary Metabolites with Unique Aromatic and Heterocyclic Structures Obtained from Terrestrial Actinomycetes Species. <i>Chemical and Pharmaceutical Bulletin</i> , 2016, 64, 668-675.	1.3	23
175	Elmenols C-H, new angucycline derivatives isolated from a culture of <i>Streptomyces</i> sp. IFM 11490. <i>Journal of Antibiotics</i> , 2017, 70, 601-606.	2.0	23
176	SO-75R1, a new mutactimycin derivative produced by <i>Nocardia brasiliensis</i> . <i>Journal of Antibiotics</i> , 1992, 45, 995-997.	2.0	22
177	Aplysidine, a new nucleoside from the Okinawan marine sponge <i>Aplysina</i> Sp.. <i>Tetrahedron</i> , 1992, 48, 7145-7148.	1.9	22
178	Colopsinols D and E, New Polyhydroxyl Linear Carbon Chain Compounds from Marine Dinoflagellate <i>Amphidinium</i> sp.. <i>Chemical and Pharmaceutical Bulletin</i> , 2000, 48, 1447-1451.	1.3	22
179	Death receptor 5 targeting activity-guided isolation of isoflavones from <i>Milletia brandisiana</i> and <i>Ardisia colorata</i> and evaluation of ability to induce TRAIL-mediated apoptosis. <i>Bioorganic and Medicinal Chemistry</i> , 2009, 17, 1181-1186.	3.0	22
180	Chromomycins A2 and A3 from Marine Actinomycetes with TRAIL Resistance-Overcoming and Wnt Signal Inhibitory Activities. <i>Marine Drugs</i> , 2014, 12, 3466-3476.	4.6	22

#	ARTICLE	IF	CITATIONS
181	Hedgehog inhibitors from <i>Withania somnifera</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015, 25, 3541-3544.	2.2	22
182	New quassinoid glycosides, yadanziosides A-H, from <i>Brucea javanica</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1984, 32, 4702-4705.	1.3	21
183	Manzamenones G and H, new dimeric fatty-acid derivatives from the Okinawan marine sponge <i>Plakortis</i> sp.. <i>Tetrahedron</i> , 1993, 49, 5955-5960.	1.9	21
184	Quinic acid esters from <i>Pluchea indica</i> with collagenase, MMP-2 and MMP-9 inhibitory activities. <i>Phytotherapy Research</i> , 2008, 22, 264-266.	5.8	21
185	Acylated Triterpenoid Saponins from <i>Schima noronhae</i> and Their Cell Growth Inhibitory Activity. <i>Journal of Natural Products</i> , 2008, 71, 918-921.	3.0	21
186	Izumiphenazine D, a New Phenazoquinoline N-Oxide from <i>Streptomyces</i> sp. IFM 11204. <i>Chemical and Pharmaceutical Bulletin</i> , 2011, 59, 508-510.	1.3	21
187	Scopadulciol, Isolated from <i>Scoparia dulcis</i> , Induces β -Catenin Degradation and Overcomes Tumor Necrosis Factor-Related Apoptosis Ligand Resistance in AGS Human Gastric Adenocarcinoma Cells. <i>Journal of Natural Products</i> , 2015, 78, 864-872.	3.0	21
188	Sesquiterpenes with TRAIL-resistance overcoming activity from <i>Xanthium strumarium</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2015, 23, 4746-4754.	3.0	21
189	Notch Inhibitors from <i>Calotropis gigantea</i> That Induce Neuronal Differentiation of Neural Stem Cells. <i>Journal of Natural Products</i> , 2017, 80, 2453-2461.	3.0	21
190	Structures of yadanziolides A, B, and C, new bitter principles from <i>Brucea javanica</i> .. <i>Chemical and Pharmaceutical Bulletin</i> , 1984, 32, 4698-4701.	1.3	20
191	New congeners of bistheonellides from Okinawan marine sponges of the genus <i>Theonella</i> . <i>Journal of the Chemical Society Perkin Transactions 1</i> , 1991, , 2379.	0.9	20
192	New Sesterterpenes with Nerve Growth Factor Synthesis-Stimulating Activity from the Okinawan Marine Sponge <i>Hirtios</i> sp.. <i>Chemical and Pharmaceutical Bulletin</i> , 1993, 41, 2190-2191.	1.3	20
193	Nepheliosyne A, New C47 Acetylenic Acid from the Okinawan Marine Sponge <i>Xestospongia</i> sp.. <i>Journal of Natural Products</i> , 1994, 57, 1300-1303.	3.0	20
194	New polyketide peroxides from Okinawan marine sponge <i>Plakortis</i> sp.. <i>Tetrahedron</i> , 1998, 54, 2041-2048.	1.9	20
195	Tubiferal A, a backbone-rearranged triterpenoid lactone isolated from the myxomycete <i>Tubifera dimorphotheca</i> , possessing reversal of drug resistance activity. <i>Tetrahedron</i> , 2004, 60, 9835-9839.	1.9	20
196	Search for Bioactive Natural Products Targeting Cancer-Related Signaling Pathways. Yuki Gosei Kagaku Kyokaiishi/ <i>Journal of Synthetic Organic Chemistry</i> , 2009, 67, 1094-1104.	0.1	20
197	A Method for the Rapid Discovery of Naturally Occurring Products by Proteins Immobilized on Magnetic Beads and Reverse Affinity Chromatography. <i>Chemistry - an Asian Journal</i> , 2009, 4, 1802-1808.	3.3	20
198	Naturally occurring Ngn2 promoter activators from <i>Butea superba</i> . <i>Molecular BioSystems</i> , 2013, 9, 2489.	2.9	20

#	ARTICLE	IF	CITATIONS
199	Two new coumarins and a new xanthone from the leaves of <i>Rhizophora mucronata</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2018, 28, 1063-1066.	2.2	20
200	Shinjulactones G and H, New Bitter Principles of <i>Ailanthus altissima</i> SWINGLE. <i>Bulletin of the Chemical Society of Japan</i> , 1984, 57, 2013-2014.	3.2	19
201	New Multiflorane-Type Triterpenoid Acids from <i>Sandoricum indicum</i> . <i>Journal of Natural Products</i> , 2001, 64, 1243-1245.	3.0	19
202	Parviflorene A, a novel cytotoxic unsymmetrical sesquiterpene dimer constituent from <i>Curcuma parviflora</i> . <i>Tetrahedron Letters</i> , 2003, 44, 2327-2329.	1.4	19
203	Physarigins A-C, three new yellow pigments from a cultured myxomycete <i>Physarum rigidum</i> . <i>Tetrahedron Letters</i> , 2003, 44, 4479-4481.	1.4	19
204	Dihydroindbladiones, Three New Naphthoquinone Pigments from a Myxomycete <i>Lindbladia tubulina</i> . <i>Journal of Natural Products</i> , 2003, 66, 999-1001.	3.0	19
205	Determination of absolute stereochemistry, total synthesis, and evaluation of peptides from the myxomycete <i>Physarum melleum</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 2008, 18, 95-98.	2.2	19
206	Cerasoidine, a Bis-aporphine Alkaloid Isolated from <i>Polyalthia cerasoides</i> during Screening for Wnt Signal Inhibitors. <i>Journal of Natural Products</i> , 2016, 79, 2083-2088.	3.0	19
207	The Notch inhibitor cowanin accelerates nicastrin degradation. <i>Scientific Reports</i> , 2018, 8, 5376.	3.3	19
208	SHINJUDILACTONE, A NEW BITTER PRINCIPLE FROM <i>AILANTHUS ALTISSIMA</i> SWINGLE. <i>Chemistry Letters</i> , 1981, 10, 1597-1598.	1.3	18
209	Shinjulactone C, a new quassinoid with a 11 \pm , 12 \pm :5 \pm , 13 \pm -dicyclo-9 \pm -picrasane skeleton from SWINGLE. <i>Tetrahedron Letters</i> , 1982, 23, 1205-1206.	1.4	18
210	Structure Determination of Bitter Principles of <i>Ailanthus altissima</i> . Structures of Shinjulactones F, I, J, and K. <i>Bulletin of the Chemical Society of Japan</i> , 1984, 57, 2885-2892.	3.2	18
211	Mangiferin Identified in a Screening Study Guided by Neuraminidase Inhibitory Activity. <i>Planta Medica</i> , 2007, 73, 1195-1196.	1.3	18
212	Cardenolide Glycosides of <i>Thevetia peruviana</i> and Triterpenoid Saponins of <i>Sapindus emarginatus</i> as TRAIL Resistance-Overcoming Compounds. <i>Journal of Natural Products</i> , 2009, 72, 1507-1511.	3.0	18
213	Structure of New Monoterpene Glycoside from <i>Sibiraea angustata</i> RCHD. and Its Anti-obestic Effect. <i>Chemical and Pharmaceutical Bulletin</i> , 2009, 57, 294-297.	1.3	18
214	Activity of Mangosteen Xanthenes and Teleocidin A-2 in Death Receptor Expression Enhancement and Tumor Necrosis Factor Related Apoptosis-Inducing Ligand Assays. <i>Journal of Natural Products</i> , 2010, 73, 452-455.	3.0	18
215	1-(1 \pm -Diazobenzyl)pyrene: A Reagent for Photolabile and Fluorescent Protection of Carboxyl Groups of Amino Acids and Peptides. <i>Synlett</i> , 1991, 1991, 35-36.	1.8	17
216	Biemnasterol, a New Cytotoxic Sterol with the Rare 22,25-Diene Side Chain, Isolated from the Marine Sponge <i>Biemna</i> sp.. <i>Journal of Natural Products</i> , 1993, 56, 2016-2018.	3.0	17

#	ARTICLE	IF	CITATIONS
217	The mode of rabbit platelet shape change and aggregation induced by theonezolid-A, a novel polyketide macrolide, isolated from the Okinawan marine sponge <i>Theonella</i> sp.. Canadian Journal of Physiology and Pharmacology, 1996, 74, 193-199.	1.4	17
218	Absolute stereochemistry of TT-1 (rasfonin), an $\hat{1}\pm$ -pyrone-containing natural product from a fungus, <i>Trichurus terrophilus</i> . Tetrahedron Letters, 2003, 44, 8427-8431.	1.4	17
219	Isolation of Bioactive Natural Products from Myxomycetes. Medicinal Chemistry, 2005, 1, 575-590.	1.5	17
220	Search for bioactive natural products from medicinal plants of Bangladesh. Journal of Natural Medicines, 2010, 64, 393-401.	2.3	17
221	Acoschimperoside P, 2 $\hat{2}$ -acetate: a Hedgehog signaling inhibitory constituent from <i>Vallisneria spiralis</i> . Journal of Natural Medicines, 2011, 65, 629-632.	2.3	17
222	Hedgehog/GLI-mediated transcriptional activity inhibitors from <i>Crinum asiaticum</i> . Journal of Natural Medicines, 2015, 69, 538-542.	2.3	17
223	Novel cytotoxic antibiotics, glucopiericidinols A1 and A2 Taxonomy, fermentation, isolation, structure elucidation and biological characteristics.. Journal of Antibiotics, 1989, 42, 1734-1740.	2.0	16
224	Wrightiamines A and B, Two New Cytotoxic Pregnane Alkaloids from <i>Wrightia javanica</i> . Chemical and Pharmaceutical Bulletin, 2003, 51, 737-739.	1.3	16
225	Isolation of Austroinulin Possessing Cell Cycle Inhibition Activity from <i>Blumea glomerata</i> and Revision of Its Absolute Configuration. Planta Medica, 2004, 70, 1170-1173.	1.3	16
226	Total synthesis of TT-1 (rasfonin), an $\hat{1}\pm$ -pyrone-containing natural product from a fungus <i>Trichurus terrophilus</i> . Tetrahedron, 2005, 61, 1827-1833.	1.9	16
227	Cytotoxic dimeric sesquiterpenoids from <i>Curcuma parviflora</i> : isolation of three new parviflorenes and absolute stereochemistry of parviflorenes A, B, D, F, and G. Tetrahedron, 2005, 61, 6700-6706.	1.9	16
228	Fuligoic acid, a new yellow pigment with a chlorinated polyene-pyrone acid structure isolated from the myxomycete <i>Fuligo septica</i> f. <i>flava</i> . Tetrahedron Letters, 2009, 50, 3189-3190.	1.4	16
229	Eudesmane-Type Sesquiterpenoid and Guaianolides from <i>Kandelia candel</i> in a Screening Program for Compounds to Overcome TRAIL Resistance. Journal of Natural Products, 2012, 75, 1431-1435.	3.0	16
230	Inubosins A, B, and C Are Acridine Alkaloids Isolated from a Culture of <i>Streptomyces</i> sp. IFM 11440 with Ngn2 Promoter Activity. Journal of Natural Products, 2015, 78, 311-314.	3.0	16
231	Hes1-Binding Compounds Isolated by Target Protein Oriented Natural Products Isolation (TPO-NAPI). Journal of Natural Products, 2017, 80, 538-543.	3.0	16
232	GLI1 Inhibitors Identified by Target Protein Oriented Natural Products Isolation (TPO-NAPI) with Hedgehog Inhibition. ACS Chemical Biology, 2018, 13, 2551-2559.	3.4	16
233	ON THE STRUCTURE OF AILANTHONE, A BITTER PRINCIPLE FROM <i>AILANTHUS ALTISSIMA</i> . Chemistry Letters, 1982, 11, 661-662.	1.3	15
234	Structure Determination of a New Bitter Principle, Shinjulactone L, from <i>Ailanthus altissima</i> . Bulletin of the Chemical Society of Japan, 1985, 58, 2723-2724.	3.2	15

#	ARTICLE	IF	CITATIONS
235	The Stereochemistry of Variabilin, an Antimicrobial Sesterterpene Isolated from Marine Sponges. <i>Natural Product Research</i> , 1993, 3, 189-192.	0.4	15
236	Amphidinolide M, a Novel 29-Membered Macrolide from the Cultured Marine Dinoflagellate <i>Amphidinium</i> sp.. <i>Journal of Organic Chemistry</i> , 1994, 59, 4698-4700.	3.2	15
237	Structure-activity relationship of bromoeudistomin D, a powerful Ca ²⁺ releaser in skeletal muscle sarcoplasmic reticulum. <i>European Journal of Pharmacology</i> , 1995, 288, 285-293.	2.6	15
238	Bahiensol, a New Glycerolipid from a Cultured Myxomycete <i>Didymium bahiense</i> var. <i>bahiense</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2003, 51, 612-613.	1.3	15
239	Efficient Synthesis of Chromone and Flavonoid Derivatives with Diverse Heterocyclic Units. <i>Chemistry - an Asian Journal</i> , 2008, 3, 2056-2064.	3.3	15
240	Glycosides from <i>Vallisneria spiralis</i> with TRAIL-Resistance-Overcoming Activity. <i>Heterocycles</i> , 2010, 80, 477.	0.7	15
241	Bioactive Natural Products from Myxomycetes Having Effects on Signaling Pathways. <i>Heterocycles</i> , 2012, 85, 1299.	0.7	15
242	Novel cytotoxic isobenzofuran derivatives from <i>Streptomyces</i> sp. IFM 11490. <i>Tetrahedron Letters</i> , 2015, 56, 6345-6347.	1.4	15
243	Coronaridine, an iboga type alkaloid from <i>Tabernaemontana divaricata</i> , inhibits the Wnt signaling pathway by decreasing β -catenin mRNA expression. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015, 25, 3937-3940.	2.2	15
244	Stereochemical Assignment and Biological Evaluation of BE-14106 Unveils the Importance of One Acetate Unit for the Antifungal Activity of Polyene Macrolactams. <i>Journal of Natural Products</i> , 2016, 79, 1877-1880.	3.0	15
245	Total synthesis of agalloside, isolated from <i>Aquilaria agallocha</i> , by the 5-O-glycosylation of flavan. <i>Organic and Biomolecular Chemistry</i> , 2017, 15, 5025-5032.	2.8	15
246	Coculture of a Pathogenic Actinomycete and Animal Cells To Produce Nocarjamide, a Cyclic Nonapeptide with Wnt Signal-Activating Effect. <i>Organic Letters</i> , 2018, 20, 5831-5834.	4.6	15
247	Parviflorene J, a Cytotoxic Sesquiterpene Dimer with a New Rearranged Skeleton from <i>Curcuma parviflora</i> . <i>Heterocycles</i> , 2007, 72, 649.	0.7	15
248	Eleven New 2-Pyrones from a Fungi Imperfecti, <i>Trichurus terrophilus</i> , Found in a Screening Study Guided by Immunomodulatory Activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2005, 53, 923-929.	1.3	14
249	New Kehokorins and Trichiols from the Myxomycete <i>Trichia favoginea</i> . <i>Heterocycles</i> , 2007, 71, 1807.	0.7	14
250	Amphidinolide B, a powerful activator of actomyosin ATPase enhances skeletal muscle contraction. <i>Biochimica Et Biophysica Acta - General Subjects</i> , 1999, 1427, 24-32.	2.4	13
251	Constituents from <i>Hoya parasitica</i> and their Cell Growth Inhibitory Activity. <i>Planta Medica</i> , 2008, 74, 760-763.	1.3	13
252	Valosinâ€containing Protein is a Target of 5â€ and Enhances TRAIL Resistance in Cancer Cells. <i>ChemistryOpen</i> , 2016, 5, 574-579.	1.9	13

#	ARTICLE	IF	CITATIONS
253	Sharkquinone, a new ana-quinonoid tetracene derivative from marine-derived <i>Streptomyces</i> sp. EGY1 with TRAIL resistance-overcoming activity. <i>Journal of Natural Medicines</i> , 2017, 71, 564-569.	2.3	13
254	Dehydropropylpantothenamide isolated by a co-culture of <i>Nocardia tenerifensis</i> IFM 10554T in the presence of animal cells. <i>Journal of Natural Medicines</i> , 2018, 72, 280-289.	2.3	13
255	The Notch Inhibitors Isolated from <i>Nerium indicum</i> . <i>Journal of Natural Products</i> , 2018, 81, 1235-1240.	3.0	13
256	Flavonoids from <i>Eupatorium odoratum</i> with Death Receptor 5 Promoter Enhancing Activity. <i>Heterocycles</i> , 2009, 77, 1379.	0.7	13
257	A new antibiotic, okicenone. II. Physico-chemical properties and structure elucidation.. <i>Journal of Antibiotics</i> , 1991, 44, 819-823.	2.0	12
258	Stereochemistry of the Modified Tryptophan Residues Contained in Konbamide and Keramamide a Isolated from Marine Sponges of the Genus <i>Theonella</i> . <i>Natural Product Research</i> , 1994, 4, 293-296.	0.4	12
259	Pyrroloiminoquinone pigments from <i>Didymium iridis</i> . <i>Biochemical Systematics and Ecology</i> , 2005, 33, 323-325.	1.3	12
260	Cribrarione C, a Naphthoquinone Pigment from the Myxomycete <i>Cribraria meylanii</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2009, 57, 894-895.	1.3	12
261	Cucullamide, a New Putrescine Bisamide from <i>Amoora cucullata</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2010, 58, 1116-1118.	1.3	12
262	Phenolic compounds from the bark of <i>Oroxylum indicum</i> activate the Ngn2 promoter. <i>Journal of Natural Medicines</i> , 2015, 69, 589-594.	2.3	12
263	Synthesis of rocaglamide derivatives and evaluation of their Wnt signal inhibitory activities. <i>Organic and Biomolecular Chemistry</i> , 2016, 14, 3061-3068.	2.8	12
264	New phenazine analogues from <i>Streptomyces</i> sp. IFM 11694 with TRAIL resistance-overcoming activities. <i>Journal of Antibiotics</i> , 2016, 69, 446-450.	2.0	12
265	Screening for natural products that affect Wnt signaling activity. <i>Journal of Natural Medicines</i> , 2019, 73, 697-705.	2.3	12
266	New Cyclic Polyketide Peroxides from Okinawan Marine Sponge <i>Plakortis</i> sp.. <i>Journal of Natural Products</i> , 1998, 61, 1427-1429.	3.0	11
267	Constituents of <i>Pongamia pinnata</i> Isolated in a Screening for Activity to Overcome Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Resistance. <i>Chemical and Pharmaceutical Bulletin</i> , 2010, 58, 1549-1551.	1.3	11
268	Naturally occurring FANCFâ€“Hes1 complex inhibitors from <i>Wrightia religiosa</i> . <i>MedChemComm</i> , 2015, 6, 455-460.	3.4	11
269	Boesenberols 1â€“K, new isopimarane diterpenes from <i>Boesenbergia pandurata</i> with TRAIL-resistance overcoming activity. <i>Tetrahedron Letters</i> , 2017, 58, 3838-3841.	1.4	11
270	Sucupiranins Aâ€“L, Furanocassane Diterpenoids from the Seeds of <i>Bowdichia virgilioides</i> . <i>Journal of Natural Products</i> , 2017, 80, 3120-3127.	3.0	11

#	ARTICLE	IF	CITATIONS
271	Conversion of ailanthone into shinjulactone C through an ionic [4+2] cycloaddition reaction. <i>Tetrahedron Letters</i> , 1983, 24, 4843-4846.	1.4	10
272	Chapter 2 Marine Alkaloids II. <i>Alkaloids: Chemistry and Pharmacology</i> , 1992, , 41-124.	0.2	10
273	Inhibition of Mammalian Topoisomerase I by Xestoquinone and Halenaquinone. <i>Bioscience, Biotechnology and Biochemistry</i> , 1993, 57, 330-331.	1.3	10
274	4, 6-Dibromo-3-hydroxycarbazole (an analogue of caffeine-like Ca ²⁺ releaser), a novel type of inhibitor of Ca ²⁺ -induced Ca ²⁺ release in skeletal muscle sarcoplasmic reticulum. <i>British Journal of Pharmacology</i> , 1995, 114, 941-948.	5.4	10
275	Casein kinase II inhibitors isolated from two Brazilian plants <i>Hymenaea parvifolia</i> and <i>Wulffia baccata</i> . <i>Bioorganic and Medicinal Chemistry Letters</i> , 1999, 9, 2157-2160.	2.2	10
276	Practical Total Synthesis of Fuligocandins A and B. <i>Synlett</i> , 2010, 2010, 2498-2502.	1.8	10
277	Prenylflavonoids isolated from <i>Artocarpus champeden</i> with TRAIL-resistance overcoming activity. <i>Phytochemistry</i> , 2013, 96, 299-304.	2.9	10
278	Identification of BMI1 Promoter Inhibitors from <i>Beaumontia murtonii</i> and <i>Eugenia operculata</i> . <i>Journal of Natural Products</i> , 2017, 80, 1853-1859.	3.0	10
279	SHINJULACTONE F, A NEW BITTER PRINCIPLE WITH A 5 ^H -PICRASANE SKELETON FROM <i>AILANTHUS ALTISSIMASWINGLE</i> . <i>Chemistry Letters</i> , 1984, 13, 555-556.	1.3	9
280	Bitter Principles of <i>Ailanthus altissima</i> SWINGLE. Conversion of Ailanthone into Shinjulactone C. <i>Bulletin of the Chemical Society of Japan</i> , 1985, 58, 2357-2360.	3.2	9
281	A new antibiotic, okicenone. I. Taxonomy, fermentation, isolation and biological characteristics.. <i>Journal of Antibiotics</i> , 1991, 44, 814-818.	2.0	9
282	Monoamine Oxidase Inhibitory Naphthoquinone and/or Naphthalene Dimers from <i>Lemuni Hitam</i> (<i>Diospyros</i> sp.), a Malaysian Herbal Medicine.. <i>Chemical and Pharmaceutical Bulletin</i> , 1999, 47, 1473-1476.	1.3	9
283	5 ^H , 8 ^H -Epidioxysterol sulfate from a diatom <i>Odontella aurita</i> . <i>Phytochemistry</i> , 2002, 61, 359-360.	2.9	9
284	Synthesis and Evaluation of Isomers of Melleumin B from the Myxomycete <i>Physarum melleum</i> ; an Approach for Wnt Signal Inhibitor Trusted in Nature. <i>Heterocycles</i> , 2008, 76, 1425.	0.7	9
285	Isolation and structure elucidation of flavonoid glycosides from <i>Solanum verbascifolium</i> . <i>Phytochemistry Letters</i> , 2010, 3, 88-92.	1.2	9
286	Nonactin and Related Compounds Found in a Screening Program for Wnt Signal Inhibitory Activity. <i>Heterocycles</i> , 2012, 84, 1245.	0.7	9
287	Î ² -Sitosterol and flavonoids isolated from <i>Bauhinia malabarica</i> found during screening for Wnt signaling inhibitory activity. <i>Journal of Natural Medicines</i> , 2014, 68, 242-245.	2.3	9
288	Hedgehog inhibitors from <i>Artocarpus communis</i> and <i>Hyptis suaveolens</i> . <i>Bioorganic and Medicinal Chemistry</i> , 2015, 23, 4150-4154.	3.0	9

#	ARTICLE	IF	CITATIONS
289	Sulfotanone, a new alkyl sulfonic acid derivative from <i>Streptomyces</i> sp. IFM 11694 with TRAIL resistance-overcoming activity. <i>Journal of Natural Medicines</i> , 2016, 70, 266-270.	2.3	9
290	Two new alkaloids from <i>Crinum asiaticum</i> var. <i>japonicum</i> . <i>Journal of Natural Medicines</i> , 2019, 73, 648-652.	2.3	9
291	Conversion of aianthone into shinjudilactone, a backbone-rearranged picrasane.. <i>Chemical and Pharmaceutical Bulletin</i> , 1982, 30, 1917-1920.	1.3	8
292	Cadinane Sesquiterpenes from <i>Curcuma parviflora</i> . <i>Journal of Natural Products</i> , 2009, 72, 782-783.	3.0	8
293	Isolation of new fuzanins, carbamate-containing natural products, from <i>Kitasatospora</i> sp. IFM10917. <i>Journal of Antibiotics</i> , 2010, 63, 385-388.	2.0	8
294	Dehydrofuligoic Acid, a New Yellow Pigment Isolated from the Myxomycete <i>Fuligo septica</i> f. <i>flava</i> . <i>Heterocycles</i> , 2010, 82, 839.	0.7	8
295	Limonoids with Wnt signal inhibitory activity isolated from the fruits of <i>Azadirachta excelsa</i> . <i>Phytochemistry Letters</i> , 2015, 11, 280-285.	1.2	8
296	Structures of shinjulactones D and E, new bitter principles of <i>Ailanthus altissima</i> Swingle.. <i>Chemical and Pharmaceutical Bulletin</i> , 1983, 31, 2179-2182.	1.3	7
297	Keruffaride, a new all-cis-cyclopentanepentol-containing metabolite from the okinawan marine sponge <i>Iuffariella</i> sp.. <i>Journal of the Chemical Society Chemical Communications</i> , 1993, , 79.	2.0	7
298	Studies on stereochemistry of theonezolides A $\frac{1}{4}$ C: Determination of absolute stereochemistry of the C-4 $\frac{1}{4}$ C-17 fragment. <i>Tetrahedron</i> , 1998, 54, 4819-4826.	1.9	7
299	First isolation of three diterpenes as naturally-occurring compounds from <i>Sindora siamensis</i> . <i>Journal of Natural Medicines</i> , 2006, 61, 77-79.	2.3	7
300	Novel tryptophan metabolites, chromoazepinone A, B and C, produced by a blocked mutant of <i>Chromobacterium violaceum</i> , the biosynthetic implications and the biological activity of chromoazepinone A and B. <i>Organic and Biomolecular Chemistry</i> , 2010, 8, 3157.	2.8	7
301	Bioactive Heterocyclic Natural Products from Actinomycetes Having Effects on Cancer-Related Signaling Pathways. <i>Progress in the Chemistry of Organic Natural Products</i> , 2014, 99, 147-198.	1.1	7
302	Boehmenan, a lignan from <i>Hibiscus ficulneus</i> , showed Wnt signal inhibitory activity. <i>Bioorganic and Medicinal Chemistry Letters</i> , 2015, 25, 2735-2738.	2.2	7
303	Identification of BMI1 promoter inhibitors from <i>Streptomyces</i> sp. IFM-11958. <i>Bioorganic and Medicinal Chemistry</i> , 2019, 27, 2998-3003.	3.0	7
304	Cadinane sesquiterpenoids isolated from <i>Santalum album</i> using a screening program for Wnt signal inhibitory activity. <i>Journal of Natural Medicines</i> , 2020, 74, 476-481.	2.3	7
305	Isolation of nocobactin NAs as Notch signal inhibitors from <i>Nocardia farcinica</i> , a possibility of invasive evolution. <i>Journal of Antibiotics</i> , 2021, 74, 255-259.	2.0	7
306	Practical Stereoselective Synthesis of C3 α -Spirooxindole β - and C2 α -Spiropseudoindoxyl β -Pyrrolidines via Organocatalyzed Pictet-Spengler Reaction/Oxidative Rearrangement Sequence. <i>Advanced Synthesis and Catalysis</i> , 2021, 363, 2648-2663.	4.3	7

#	ARTICLE	IF	CITATIONS
307	Tryptophan 4-Aminocinnamamides from a Myxomycete <i>Fuligo aurea</i> . <i>Heterocycles</i> , 2006, 69, 463.	0.7	7
308	Studies on the Stereochemistry of Theonezolides A-C: Elucidation of the Relative Configurations of 1,3-Diol Moieties of the C-4...C-17 Fragment. <i>Heterocycles</i> , 1998, 49, 39.	0.7	7
309	Development of Unexploited Microbial Resources: Natural Products from Slime Molds.. Yuki Gosei Kagaku Kyokaiishi/Journal of Synthetic Organic Chemistry, 2003, 61, 152-163.	0.1	6
310	Tubiferic Acid, a New 9,10-Secocycloartane Triterpenoid Acid Isolated from the Myxomycete <i>Tubulifera arachnoidea</i> . <i>Chemical and Pharmaceutical Bulletin</i> , 2011, 59, 279-281.	1.3	6
311	Isolation of Nabscessin C from <i>Nocardia abscessus</i> ; IFM 10029 and a Study on Biosynthetic Pathway for Nabscessins. <i>Chemical and Pharmaceutical Bulletin</i> , 2018, 66, 976-982.	1.3	6
312	Isolation and evaluation of cardenolides from <i>Lansium domesticum</i> as Notch inhibitors. <i>Journal of Natural Medicines</i> , 2020, 74, 758-766.	2.3	6
313	9-Methyl-7-bromoedistomin D induces Ca ²⁺ release from cardiac sarcoplasmic reticulum. <i>European Journal of Pharmacology</i> , 1998, 357, 261-265.	3.5	5
314	Bioactive secondary metabolites from Okinawan sponges and tunicates. <i>Studies in Natural Products Chemistry</i> , 2000, 23, 185-231.	1.8	5
315	Alkenylphenol and alkenylsalicylic acid from <i>Mangifera caesia</i> . <i>Biochemical Systematics and Ecology</i> , 2002, 30, 475-478.	1.3	5
316	Flavonoids and sesquiterpenoids, constituents from <i>Eupatorium capillifolium</i> , found in a screening study guided by cell growth inhibitory activity. <i>Journal of Natural Medicines</i> , 2006, 60, 325-328.	2.3	5
317	Total synthesis and evaluation of Wnt signal inhibition of melleumin A and B, and their derivatives. <i>Organic and Biomolecular Chemistry</i> , 2011, 8, 5285-93.	2.8	5
318	Total syntheses of schizandriside, saracoside and (±)-isolariciresinol with antioxidant activities. <i>Journal of Natural Medicines</i> , 2018, 72, 651-654.	2.3	5
319	Characterization of Nocardithiocin Derivatives Produced by Amino Acid Substitution of Precursor Peptide notG. <i>International Journal of Peptide Research and Therapeutics</i> , 2020, 26, 281-290.	1.9	5
320	Target protein-oriented isolation of Hes1 dimer inhibitors using protein based methods. <i>Scientific Reports</i> , 2020, 10, 1381.	3.3	5
321	Thannilignan glucoside and 2-(β -glucopyranosyl)-3-isoxazolin-5-one derivative, two new compounds isolated from <i>Terminalia bellirica</i> . <i>Journal of Natural Medicines</i> , 2022, 76, 482-489.	2.3	5
322	Two Bioactive Compounds, Uniformides A and B, Isolated from a Culture of <i>Nocardia uniformis</i> IFM0856 in the Presence of Animal Cells. <i>Organic Letters</i> , 2022, 24, 4998-5002.	4.6	5
323	Bitter principles of <i>Ailanthus altissima</i> Swingle. Studies on oxidative degradation of the A-ring of ailanthone derivatives.. <i>Chemical and Pharmaceutical Bulletin</i> , 1987, 35, 3011-3015.	1.3	4
324	Total synthesis of lindbladione, a Hes1 dimerization inhibitor and neural stem cell activator isolated from <i>Lindbladia tubulina</i> . <i>Scientific Reports</i> , 2020, 10, 21433.	3.3	4

#	ARTICLE	IF	CITATIONS
325	Evaluation of Naturally Occurring HIF α Inhibitors for Pulmonary Arterial Hypertension. <i>ChemBioChem</i> , 2021, 22, 2799-2804.	2.6	4
326	Constituents from the Rhizomes of <i>Curcuma comosa</i> and Their Wnt Signal Inhibitory Activities. <i>Heterocycles</i> , 2014, 88, 1501.	0.7	4
327	Trichiol and 3-epitrichiol acetate, novel cytotoxic sterols with an unprecedented 2,6-dioxabicyclo[2.2.2]octan-3-one ring system from the myxomycete <i>Trichia favoginea</i> var. <i>persimilis</i> . <i>Tetrahedron Letters</i> , 2006, 47, 4351-4354.	1.4	3
328	GRISEOVIRIDIN AND CYCLIC HYDROXAMATES FOUND IN A SCREENING PROGRAM FOR Wnt SIGNAL INHIBITOR. <i>Heterocycles</i> , 2012, 86, 1517.	0.7	3
329	Synthesis and Evaluation of Fuligocandin B Derivatives with Activity for Overcoming TRAIL Resistance. <i>Chemical and Pharmaceutical Bulletin</i> , 2018, 66, 810-817.	1.3	3
330	Isolation of Inohanalactone, a β -Butyrolactone, from <i>Nocardia inohanensis</i> IFM0092T. <i>Heterocycles</i> , 2020, 101, 312.	0.7	3
331	Isolation of Peptidolipin NA Derivatives from the Culture of <i>Nocardia arthritidis</i> IFM10035T in the Presence of Mouse Macrophage Cells. <i>Heterocycles</i> , 2022, 104, 185.	0.7	3
332	Is 18 β -Glycyrrhizin a real natural product? Improved preparation of 18 β -Glycyrrhizin from 18 β -Glycyrrhizin as a positive standard for HPLC analysis of licorice extracts. <i>Journal of Natural Medicines</i> , 2022, 76, 367-378.	2.3	3
333	Phorbol esters with wnt signal-augmenting effects isolated from <i>Excoecaria indica</i> . <i>Natural Product Communications</i> , 2012, 7, 475-7.	0.5	3
334	Structure-Activity Relationships for the Ca ²⁺ -releasing Activity of 6-Hydroxy- β -carboline Analogues in Skeletal Muscle Sarcoplasmic Reticulum—The Effects of Halogen Substitution at C-5 and C-7. <i>Journal of Pharmacy and Pharmacology</i> , 2010, 52, 517-521.	2.4	2
335	A New Resin Glycoside from <i>Ipomoea Maxima</i> . <i>Natural Product Communications</i> , 2012, 7, 1934578X1200700.	0.5	2
336	Isolation of β -Indomycinone Guided by Cytotoxicity Tests from <i>Streptomyces</i> sp. IFM11607 and Revision of its Double Bond Geometry. <i>Natural Product Communications</i> , 2014, 9, 1934578X1400900.	0.5	2
337	Efficient Synthesis of Heterocyclic Flavonoids with Hedgehog Signal Inhibitory Activity. <i>Chemical and Pharmaceutical Bulletin</i> , 2017, 65, 784-795.	1.3	2
338	Nonactic Acid Derivatives Isolated From <i>Streptomyces werraensis</i> IFM12104 in a Screening Program for BMI1 Promoter Inhibitory Activity. <i>Natural Product Communications</i> , 2019, 14, 1934578X1986658.	0.5	2
339	Isolation of Adenosine and Cordysin B from <i>Anredera cordifolia</i> that Stimulates CRE-Mediated Transcription in PC12 Cells. <i>Planta Medica International Open</i> , 2021, 8, e19-e24.	0.5	2
340	Screening study of cancer-related cellular signals from microbial natural products. <i>Journal of Antibiotics</i> , 2021, 74, 629-638.	2.0	2
341	Isolation and Evaluation of Hedgehog Inhibitors from Christmas Grass (<i>Themeda arguens</i>). <i>Heterocycles</i> , 2017, 95, 210.	0.7	2
342	Isolation of β -indomycinone guided by cytotoxicity tests from <i>Streptomyces</i> sp. IFM11607 and revision of its double bond geometry. <i>Natural Product Communications</i> , 2014, 9, 1327-8.	0.5	2

#	ARTICLE	IF	CITATIONS
343	Bioactivity-guided Isolation of TRAIL-Resistance-Overcoming Activity Compounds From the Leaves of <i>Murraya exotica</i> . Natural Product Communications, 2021, 16, 1934578X2110658.	0.5	2
344	Phorbol Esters with Wnt Signal-Augmenting Effects Isolated from <i>Excoecaria Indica</i> . Natural Product Communications, 2012, 7, 1934578X1200700.	0.5	1
345	Isolation of Alkamides with Death Receptor-Enhancing Activities from <i>Piper chaba</i> . Heterocycles, 2015, 90, 1317.	0.7	1
346	Bioassay-Guided Isolation of Compounds from <i>Datura Stramonium</i> with TRAIL-Resistance Overcoming Activity. Natural Product Communications, 2016, 11, 1934578X1601100.	0.5	1
347	Constituents from <i>Entada scandens</i> with TRAIL-resistance Overcoming Activity. Natural Product Communications, 2016, 11, 1934578X1601100.	0.5	1
348	New Flavanol Dimers from the Bark of <i>Celtis tetrandra</i> and Their TRAIL Resistance-Overcoming Activity. Natural Product Communications, 2018, 13, 1934578X1801300.	0.5	1
349	Identification of 1 ² ,2 ¹ -epoxytagitinin C as a Notch inhibitor, oxidative stress mechanism and its anti-leukemia activity. Journal of Natural Medicines, 2021, , .	2.3	1
350	9-Methyl-7-bromoedistomin D, a powerful caffeine-like Ca ²⁺ releaser, acts on Ca ²⁺ -induced Ca ²⁺ release channels of cardiac sarcoplasmic reticulum. The Japanese Journal of Pharmacology, 1998, 76, 204.	1.2	0
351	Development of Unexploited Microbial Resources: Natural Products from Slime Molds. ChemInform, 2003, 34, no.	0.0	0
352	Cribrarione A, a New Antimicrobial Naphthoquinone Pigment from a Myxomycete <i>Cribraria purpurea</i> . ChemInform, 2003, 34, no.	0.0	0
353	Physarigins A-C, Three New Yellow Pigments from a Cultured Myxomycete <i>Physarum rigidum</i> . ChemInform, 2003, 34, no.	0.0	0
354	Discovery of New Macrolides from Marine Organisms. , 2003, , 57-98.		0
355	Germacranolides from <i>Enhydra fluctuans</i> with TRAIL-resistance Abrogating Activity. Natural Product Communications, 2017, 12, 1934578X1701200.	0.5	0
356	Greetings from the chief editor. Journal of Natural Medicines, 2018, 72, 595-595.	2.3	0
357	Sucupiranins Q, five new furanocassane-type diterpenoids from the seeds of <i>Bowdichia virgilioides</i> . Tetrahedron, 2019, 75, 130511.	1.9	0
358	Acacienone, a terpenoid-like natural product having an unprecedented C20 framework isolated from <i>Acacia mangium</i> leaves. Journal of Natural Medicines, 2021, 75, 99-104.	2.3	0
359	Target Protein-Oriented Isolations for Chemical Biology based on Natural Products. Yuki Gosei Kagaku Kyokaiishi/Journal of Synthetic Organic Chemistry, 2021, 79, 684-693.	0.1	0
360	Dihydroflavonol BB-1, an Extract of Natural Plant <i>Blumea Balsamifera</i> , Abrogates TRAIL Resistance in Adult T-Cell Leukemia/Lymphoma.. Blood, 2004, 104, 2290-2290.	1.4	0

#	ARTICLE	IF	CITATIONS
361	Cell Growth and Cell Cycle Inhibitory Activities of 20-Epidiosgenyl Saponin from <i>Calamus insignis</i> . <i>Heterocycles</i> , 2007, 74, 931.	0.7	0
362	Bioactive Substances from Marine Organisms. <i>Journal of Fiber Science and Technology</i> , 1990, 46, P283-P287.	0.0	0
363	Target Protein-Oriented Natural Product Isolation Methods. , 2020, , 457-474.		0
364	The mode of rabbit platelet shape change and aggregation induced by theonezolid-A, a novel polyketide macrolide, isolated from the Okinawan marine sponge <l>Theonella</l> sp.. <i>Canadian Journal of Physiology and Pharmacology</i> , 1996, 74, 193-199.	1.4	0
365	Isolation of Ikahonone, 4-Methyl-2,4-dihydroxy-3-pentanone from <i>Bacillus cereus</i> IFM12235. <i>Heterocycles</i> , 2022, 105, 523.	0.7	0