
List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3754911/publications.pdf Version: 2024-02-01



#	Article	lF	CITATIONS
1	Thannilignan glucoside and 2-(β-glucopyranosyl)-3-isoxazolin-5-one derivative, two new compounds isolated from Terminalia bellirica. Journal of Natural Medicines, 2022, 76, 482-489.	2.3	5
2	Isolation of Peptidolipin NA Derivatives from the Culture of Nocardia arthritidis IFM10035T in the Presence of Mouse Macrophage Cells. Heterocycles, 2022, 104, 185.	0.7	3
3	Is 18α-Glycyrrhizin a real natural product? Improved preparation of 18α-Glycyrrhizin from 18Î2-Glycyrrhizin as a positive standard for HPLC analysis of licorice extracts. Journal of Natural Medicines, 2022, 76, 367-378.	2.3	3
4	Isolation of Ikahonone, 4-Methyl-2,4-dihydroxy-3-pentanone from Bacillus cereus IFM12235. Heterocycles, 2022, 105, 523.	0.7	0
5	Two Bioactive Compounds, Uniformides A and B, Isolated from a Culture of <i>Nocardia uniformis</i> IFM0856 ^T in the Presence of Animal Cells. Organic Letters, 2022, 24, 4998-5002.	4.6	5
6	Acacienone, a terpenoid-like natural product having an unprecedented C20 framework isolated from Acacia mangium leaves. Journal of Natural Medicines, 2021, 75, 99-104.	2.3	0
7	Isolation of nocobactin NAs as Notch signal inhibitors from Nocardia farcinica, a possibility of invasive evolution. Journal of Antibiotics, 2021, 74, 255-259.	2.0	7
8	Practical Stereoselective Synthesis of C3â€Spirooxindole―and C2â€Spiropseudoindoxylâ€Pyrrolidines <i>via</i> Organocatalyzed Pictetâ€Spengler Reaction/Oxidative Rearrangement Sequence. Advanced Synthesis and Catalysis, 2021, 363, 2648-2663.	4.3	7
9	Isolation of Adenosine and Cordysinin B from Anredera cordifolia that Stimulates CRE-Mediated Transcription in PC12 Cells. Planta Medica International Open, 2021, 8, e19-e24.	0.5	2
10	Screening study of cancer-related cellular signals from microbial natural products. Journal of Antibiotics, 2021, 74, 629-638.	2.0	2
11	Evaluation of Naturally Occurring HIFâ€1 Inhibitors for Pulmonary Arterial Hypertension. ChemBioChem, 2021, 22, 2799-2804.	2.6	4
12	Target Protein-Oriented Isolations for Chemical Biology based on Natural Products. Yuki Gosei Kagaku Kyokaishi/Journal of Synthetic Organic Chemistry, 2021, 79, 684-693.	0.1	0
13	Identification of $1\hat{l}^2,2\hat{l}\pm$ -epoxytagitinin C as a Notch inhibitor, oxidative stress mechanism and its anti-leukemia activity. Journal of Natural Medicines, 2021, , .	2.3	1
14	Bioactivity-guided Isolation of TRAIL-Resistance-Overcoming Activity Compounds From the Leaves of <i>Murraya exotica</i> . Natural Product Communications, 2021, 16, 1934578X2110658.	0.5	2
15	Characterization of Nocardithiocin Derivatives Produced by Amino Acid Substitution of Precursor Peptide notG. International Journal of Peptide Research and Therapeutics, 2020, 26, 281-290.	1.9	5
16	Cadinane sesquiterpenoids isolated from Santalum album using a screening program for Wnt signal inhibitory activity. Journal of Natural Medicines, 2020, 74, 476-481.	2.3	7
17	Total synthesis of lindbladione, a Hes1 dimerization inhibitor and neural stem cell activator isolated from Lindbladia tubulina. Scientific Reports, 2020, 10, 21433.	3.3	4
18	Isolation and evaluation of cardenolides from Lansium domesticum as Notch inhibitors. Journal of Natural Medicines, 2020, 74, 758-766.	2.3	6

#	Article	IF	CITATIONS
19	Target protein-oriented isolation of Hes1 dimer inhibitors using protein based methods. Scientific Reports, 2020, 10, 1381.	3.3	5
20	Isolation of Inohanalactone, a Î ³ -Butyrolactone, from Nocardia inohanensis IFM0092T. Heterocycles, 2020, 101, 312.	0.7	3
21	Target Protein-Oriented Natural Product Isolation Methods. , 2020, , 457-474.		0
22	Sucupiranins M–Q, five new furanocassane-type diterpenoids from the seeds of Bowdichia virgilioides. Tetrahedron, 2019, 75, 130511.	1.9	0
23	Nonactic Acid Derivatives Isolated From Streptomyces werraensis IFM12104 in a Screening Program for BMI1 Promoter Inhibitory Activity. Natural Product Communications, 2019, 14, 1934578X1986658.	0.5	2
24	Screening for natural products that affect Wnt signaling activity. Journal of Natural Medicines, 2019, 73, 697-705.	2.3	12
25	Identification of BMI1 promoter inhibitors from Streptomyces sp. IFM-11958. Bioorganic and Medicinal Chemistry, 2019, 27, 2998-3003.	3.0	7
26	Two new alkaloids from Crinum asiaticum var. japonicum. Journal of Natural Medicines, 2019, 73, 648-652.	2.3	9
27	Dehydropropylpantothenamide isolated by a co-culture of Nocardia tenerifensis IFM 10554T in the presence of animal cells. Journal of Natural Medicines, 2018, 72, 280-289.	2.3	13
28	Total syntheses of schizandriside, saracoside and (±)-isolariciresinol with antioxidant activities. Journal of Natural Medicines, 2018, 72, 651-654.	2.3	5
29	The Notch Inhibitors Isolated from <i>Nerium indicum</i> . Journal of Natural Products, 2018, 81, 1235-1240.	3.0	13
30	Two new coumarins and a new xanthone from the leaves of Rhizophora mucronata. Bioorganic and Medicinal Chemistry Letters, 2018, 28, 1063-1066.	2.2	20
31	Greetings from the chief editor. Journal of Natural Medicines, 2018, 72, 595-595.	2.3	Ο
32	The Notch inhibitor cowanin accelerates nicastrin degradation. Scientific Reports, 2018, 8, 5376.	3.3	19
33	New Flavanol Dimers from the Bark of <i>Celtis tetrandra</i> and Their TRAIL Resistance-Overcoming Activity. Natural Product Communications, 2018, 13, 1934578X1801300.	0.5	1
34	lsolation of Nabscessin C from <i>Nocardia abscessus</i> IFM 10029 ^T and a Study on Biosynthetic Pathway for Nabscessins. Chemical and Pharmaceutical Bulletin, 2018, 66, 976-982.	1.3	6
35	Coculture of a Pathogenic Actinomycete and Animal Cells To Produce Nocarjamide, a Cyclic Nonapeptide with Wnt Signal-Activating Effect. Organic Letters, 2018, 20, 5831-5834.	4.6	15
36	GLI1 Inhibitors Identified by Target Protein Oriented Natural Products Isolation (TPO-NAPI) with Hedgehog Inhibition. ACS Chemical Biology, 2018, 13, 2551-2559.	3.4	16

#	Article	IF	CITATIONS
37	Synthesis and Evaluation of Fuligocandin B Derivatives with Activity for Overcoming TRAIL Resistance. Chemical and Pharmaceutical Bulletin, 2018, 66, 810-817.	1.3	3
38	Nabscessins A and B, Aminocyclitol Derivatives from <i>Nocardia abscessus</i> IFM 10029 ^T . Journal of Natural Products, 2017, 80, 565-568.	3.0	24
39	Elmenols C-H, new angucycline derivatives isolated from a culture of Streptomyces sp. IFM 11490. Journal of Antibiotics, 2017, 70, 601-606.	2.0	23
40	Hes1-Binding Compounds Isolated by Target Protein Oriented Natural Products Isolation (TPO-NAPI). Journal of Natural Products, 2017, 80, 538-543.	3.0	16
41	Identification of BMI1 Promoter Inhibitors from <i>Beaumontia murtonii</i> and <i>Eugenia operculata</i> . Journal of Natural Products, 2017, 80, 1853-1859.	3.0	10
42	Total synthesis of agalloside, isolated from Aquilaria agallocha, by the 5-O-glycosylation of flavan. Organic and Biomolecular Chemistry, 2017, 15, 5025-5032.	2.8	15
43	Sharkquinone, a new ana-quinonoid tetracene derivative from marine-derived Streptomyces sp. EGY1 with TRAIL resistance-overcoming activity. Journal of Natural Medicines, 2017, 71, 564-569.	2.3	13
44	Boesenberols l–K, new isopimarane diterpenes from Boesenbergia pandurata with TRAIL-resistance overcoming activity. Tetrahedron Letters, 2017, 58, 3838-3841.	1.4	11
45	Notch Inhibitors from <i>Calotropis gigantea</i> That Induce Neuronal Differentiation of Neural Stem Cells. Journal of Natural Products, 2017, 80, 2453-2461.	3.0	21
46	Sucupiranins A–L, Furanocassane Diterpenoids from the Seeds of <i>Bowdichia virgilioides</i> . Journal of Natural Products, 2017, 80, 3120-3127.	3.0	11
47	Efficient Synthesis of Heterocyclic Flavonoids with Hedgehog Signal Inhibitory Activity. Chemical and Pharmaceutical Bulletin, 2017, 65, 784-795.	1.3	2
48	Germacranolides from Enhydra fluctuans with TRAIL-resistance Abrogating Activity. Natural Product Communications, 2017, 12, 1934578X1701200.	0.5	0
49	Anti-proliferative effect of Moringa oleifera Lam (Moringaceae) leaf extract on human colon cancer HCT116 cell line. Tropical Journal of Pharmaceutical Research, 2017, 16, 371.	0.3	28
50	Isolation and Evaluation of Hedgehog Inhibitors from Christmas Grass (Themeda arguens). Heterocycles, 2017, 95, 210.	0.7	2
51	Bioassay-Guided Isolation of Compounds from <i>Datura Stramonium</i> with TRAIL-Resistance Overcoming Activity. Natural Product Communications, 2016, 11, 1934578X1601100.	0.5	1
52	Constituents from Entada scandens with TRAIL-resistance Overcoming Activity. Natural Product Communications, 2016, 11, 1934578X1601100.	0.5	1
53	Valosinâ€containing Protein is a Target of 5′â€l Fuligocandin B and Enhances TRAIL Resistance in Cancer Cells. ChemistryOpen, 2016, 5, 574-579.	1.9	13
54	Cerasoidine, a Bis-aporphine Alkaloid Isolated from <i>Polyalthia cerasoides</i> during Screening for Wnt Signal Inhibitors. Journal of Natural Products, 2016, 79, 2083-2088.	3.0	19

#	Article	IF	CITATIONS
55	Boesenberols, Pimarane Diterpenes with TRAIL-Resistance-Overcoming Activity from <i>Boesenbergia pandurata</i> . Journal of Natural Products, 2016, 79, 2075-2082.	3.0	24
56	Bio-active Natural Products with TRAIL-Resistance Overcoming Activity. Chemical and Pharmaceutical Bulletin, 2016, 64, 119-127.	1.3	24
57	Bioactive Secondary Metabolites with Unique Aromatic and Heterocyclic Structures Obtained from Terrestrial Actinomycetes Species. Chemical and Pharmaceutical Bulletin, 2016, 64, 668-675.	1.3	23
58	Stereochemical Assignment and Biological Evaluation of BE-14106 Unveils the Importance of One Acetate Unit for the Antifungal Activity of Polyene Macrolactams. Journal of Natural Products, 2016, 79, 1877-1880.	3.0	15
59	Hes1 inhibitor isolated by target protein oriented natural products isolation (TPO-NAPI) of differentiation activators of neural stem cells. Chemical Science, 2016, 7, 1514-1520.	7.4	25
60	Synthesis of rocaglamide derivatives and evaluation of their Wnt signal inhibitory activities. Organic and Biomolecular Chemistry, 2016, 14, 3061-3068.	2.8	12
61	New phenazine analogues from Streptomyces sp. IFM 11694 with TRAIL resistance-overcoming activities. Journal of Antibiotics, 2016, 69, 446-450.	2.0	12
62	Sulfotanone, a new alkyl sulfonic acid derivative from Streptomyces sp. IFM 11694 with TRAIL resistance-overcoming activity. Journal of Natural Medicines, 2016, 70, 266-270.	2.3	9
63	Isolation of Alkamides with Death Receptor-Enhancing Activities from Piper chaba. Heterocycles, 2015, 90, 1317.	0.7	1
64	Scopadulciol, Isolated from <i>Scoparia dulcis</i> , Induces β-Catenin Degradation and Overcomes Tumor Necrosis Factor-Related Apoptosis Ligand Resistance in AGS Human Gastric Adenocarcinoma Cells. Journal of Natural Products, 2015, 78, 864-872.	3.0	21
65	Naturally occurring FANCF–Hes1 complex inhibitors from Wrightia religiosa. MedChemComm, 2015, 6, 455-460.	3.4	11
66	Hedgehog/GLI-mediated transcriptional activity inhibitors from Crinum asiaticum. Journal of Natural Medicines, 2015, 69, 538-542.	2.3	17
67	Boehmenan, a lignan from Hibiscus ficulneus, showed Wnt signal inhibitory activity. Bioorganic and Medicinal Chemistry Letters, 2015, 25, 2735-2738.	2.2	7
68	Novel cytotoxic isobenzofuran derivatives from Streptomyces sp. IFM 11490. Tetrahedron Letters, 2015, 56, 6345-6347.	1.4	15
69	Inubosins A, B, and C Are Acridine Alkaloids Isolated from a Culture of <i>Streptomyces</i> sp. IFM 11440 with Ngn2 Promoter Activity. Journal of Natural Products, 2015, 78, 311-314.	3.0	16
70	Limonoids with Wnt signal inhibitory activity isolated from the fruits of Azadirachta excelsa. Phytochemistry Letters, 2015, 11, 280-285.	1.2	8
71	Hedgehog inhibitors from Withania somnifera. Bioorganic and Medicinal Chemistry Letters, 2015, 25, 3541-3544.	2.2	22
72	Coronaridine, an iboga type alkaloid from Tabernaemontana divaricata, inhibits the Wnt signaling pathway by decreasing l²-catenin mRNA expression. Bioorganic and Medicinal Chemistry Letters, 2015, 25, 3937-3940.	2.2	15

#	Article	IF	CITATIONS
73	Phenolic compounds from the bark of Oroxylum indicum activate the Ngn2 promoter. Journal of Natural Medicines, 2015, 69, 589-594.	2.3	12
74	Sesquiterpenes with TRAIL-resistance overcoming activity from Xanthium strumarium. Bioorganic and Medicinal Chemistry, 2015, 23, 4746-4754.	3.0	21
75	Hedgehog inhibitors from Artocarpus communis and Hyptis suaveolens. Bioorganic and Medicinal Chemistry, 2015, 23, 4150-4154.	3.0	9
76	9-Hydroxycanthin-6-one, a β-Carboline Alkaloid from <i>Eurycoma longifolia</i> , Is the First Wnt Signal Inhibitor through Activation of Glycogen Synthase Kinase 3Ĩ² without Depending on Casein Kinase 1α. Journal of Natural Products, 2015, 78, 1139-1146.	3.0	38
77	Natural compounds with Wnt signal modulating activity. Natural Product Reports, 2015, 32, 1622-1628.	10.3	33
78	Prenylated Flavonoids and Resveratrol Derivatives Isolated from <i>Artocarpus communis</i> with the Ability to Overcome TRAIL Resistance. Journal of Natural Products, 2015, 78, 103-110.	3.0	28
79	Chromomycins A2 and A3 from Marine Actinomycetes with TRAIL Resistance-Overcoming and Wnt Signal Inhibitory Activities. Marine Drugs, 2014, 12, 3466-3476.	4.6	22
80	Physalin H from <i>Solanum nigrum</i> as an Hh signaling inhibitor blocks GLI1–DNA-complex formation. Beilstein Journal of Organic Chemistry, 2014, 10, 134-140.	2.2	27
81	Isolation of β-Indomycinone Guided by Cytotoxicity Tests from Streptomyces sp. IFM11607 and Revision of its Double Bond Geometry. Natural Product Communications, 2014, 9, 1934578X1400900.	0.5	2
82	Bioactive Heterocyclic Natural Products from Actinomycetes Having Effects on Cancer-Related Signaling Pathways. Progress in the Chemistry of Organic Natural Products, 2014, 99, 147-198.	1.1	7
83	β-Sitosterol and flavonoids isolated from Bauhinia malabarica found during screening for Wnt signaling inhibitory activity. Journal of Natural Medicines, 2014, 68, 242-245.	2.3	9
84	Calotropin: A Cardenolide from <i>Calotropis gigantea</i> that Inhibits Wnt Signaling by Increasing Casein Kinase 11± in Colon Cancer Cells. ChemBioChem, 2014, 15, 872-878.	2.6	44
85	Ricinine: A pyridone alkaloid from Ricinus communis that activates the Wnt signaling pathway through casein kinase 11±. Bioorganic and Medicinal Chemistry, 2014, 22, 4597-4601.	3.0	25
86	Constituents from the Rhizomes of Curcuma comosa and Their Wnt Signal Inhibitory Activities. Heterocycles, 2014, 88, 1501.	0.7	4
87	Isolation of β-indomycinone guided by cytotoxicity tests from Streptomyces sp. IFM11607 and revision of its double bond geometry. Natural Product Communications, 2014, 9, 1327-8.	0.5	2
88	Naturally occurring Ngn2 promoter activators from Butea superba. Molecular BioSystems, 2013, 9, 2489.	2.9	20
89	Prenylflavonoids isolated from Artocarpus champeden with TRAIL-resistance overcoming activity. Phytochemistry, 2013, 96, 299-304.	2.9	10
90	Hh signaling inhibitors from Vitex negundo; naturally occurring inhibitors of the GLI1–DNA complex. Molecular BioSystems, 2013, 9, 1012.	2.9	23

#	Article	IF	CITATIONS
91	Catalytic Asymmetric Synthesis of Mixed 3,3′â€Bisindoles and Their Evaluation as Wnt Signaling Inhibitors. Angewandte Chemie - International Edition, 2013, 52, 2486-2490.	13.8	81
92	Xylogranin B: A Potent Wnt Signal Inhibitory Limonoid from <i>Xylocarpus granatum</i> . Organic Letters, 2013, 15, 6106-6109.	4.6	32
93	Cycloartane Triterpenes and Ingol Diterpenes Isolated from Euphorbia neriifolia in a Screening Program for Death-Receptor Expression-Enhancing Activity. Planta Medica, 2012, 78, 1370-1377.	1.3	32
94	GRISEOVIRIDIN AND CYCLIC HYDROXAMATES FOUND IN A SCREENING PROGRAM FOR Wnt SIGNAL INHIBITOR. Heterocycles, 2012, 86, 1517.	0.7	3
95	Nonactin and Related Compounds Found in a Screening Program for Wnt Signal Inhibitory Activity. Heterocycles, 2012, 84, 1245.	0.7	9
96	Eudesmane-Type Sesquiterpenoid and Guaianolides from <i>Kandelia candel</i> in a Screening Program for Compounds to Overcome TRAIL Resistance. Journal of Natural Products, 2012, 75, 1431-1435.	3.0	16
97	Yoropyrazone, a new naphthopyridazone alkaloid isolated from Streptomyces sp. IFM 11307 and evaluation of its TRAIL resistance-overcoming activity. Journal of Antibiotics, 2012, 65, 245-248.	2.0	31
98	Phorbol Esters with Wnt Signal-Augmenting Effects Isolated from <i>Excoecaria Indica</i> . Natural Product Communications, 2012, 7, 1934578X1200700.	0.5	1
99	A New Resin Glycoside from Ipomoea Maxima. Natural Product Communications, 2012, 7, 1934578X1200700.	0.5	2
100	Bioactive Natural Products from Myxomycetes Having Effects on Signaling Pathways. Heterocycles, 2012, 85, 1299.	0.7	15
101	Katorazone, a new yellow pigment with a 2-azaquinone-phenylhydrazone structure produced by Streptomyces sp. IFM 11299. Tetrahedron Letters, 2012, 53, 3346-3348.	1.4	29
102	Phorbol esters with wnt signal-augmenting effects isolated from Excoecaria indica. Natural Product Communications, 2012, 7, 475-7.	0.5	3
103	Cycloartane Triterpenes Isolated from <i>Combretum quadrangulare</i> in a Screening Program for Death-Receptor Expression Enhancing Activity. Journal of Natural Products, 2011, 74, 249-255.	3.0	39
104	New pyranonaphthoquinones and a phenazine alkaloid isolated from Streptomyces sp. IFM 11307 with TRAIL resistance-overcoming activity. Journal of Antibiotics, 2011, 64, 729-734.	2.0	30
105	New hedgehog/GLI-signaling inhibitors from Adenium obesum. Organic and Biomolecular Chemistry, 2011, 9, 1133-1139.	2.8	33
106	Total synthesis and evaluation of Wnt signal inhibition of melleumin A and B, and their derivatives. Organic and Biomolecular Chemistry, 2011, 8, 5285-93.	2.8	5
107	Tubiferic Acid, a New 9,10-Secocycloartane Triterpenoid Acid Isolated from the Myxomycete Tubulifera arachnoidea. Chemical and Pharmaceutical Bulletin, 2011, 59, 279-281.	1.3	6
108	Izumiphenazine D, a New Phenazoquinoline N-Oxide from Streptomyces sp. IFM 11204. Chemical and Pharmaceutical Bulletin, 2011, 59, 508-510.	1.3	21

#	Article	IF	CITATIONS
109	Isolation and structure elucidation of izuminosides A–C: a rare phenazine glycosides from Streptomyces sp. IFM 11260. Journal of Antibiotics, 2011, 64, 271-275.	2.0	32
110	2-Methoxy-1,4-naphthoquinone isolated from Impatiens balsamina in a screening program for activity to inhibit Wnt signaling. Journal of Natural Medicines, 2011, 65, 234-236.	2.3	23
111	Acoschimperoside P, 2′-acetate: a Hedgehog signaling inhibitory constituent from Vallaris glabra. Journal of Natural Medicines, 2011, 65, 629-632.	2.3	17
112	Cryptolepine, isolated from <i>Sida acuta</i> , sensitizes human gastric adenocarcinoma cells to TRAILâ€induced apoptosis. Phytotherapy Research, 2011, 25, 147-150.	5.8	31
113	New Hedgehog/GLI signaling inhibitors from Excoecaria agallocha. Bioorganic and Medicinal Chemistry Letters, 2011, 21, 718-722.	2.2	41
114	Cucullamide, a New Putrescine Bisamide from Amoora cucullata. Chemical and Pharmaceutical Bulletin, 2010, 58, 1116-1118.	1.3	12
115	Glycosides from Vallaris solanaceae with TRAIL-Resistance-Overcoming Activity. Heterocycles, 2010, 80, 477.	0.7	15
116	Constituents of Pongamia pinnata Isolated in a Screening for Activity to Overcome Tumor Necrosis Factor-Related Apoptosis-Inducing Ligand-Resistance. Chemical and Pharmaceutical Bulletin, 2010, 58, 1549-1551.	1.3	11
117	Structure-Activity Relationships for the Ca2+-releasing Activity of 6-Hydroxy-β-carboline Analogues in Skeletal Muscle Sarcoplasmic Reticulum—The Effects of Halogen Substitution at C-5 and C-7. Journal of Pharmacy and Pharmacology, 2010, 52, 517-521.	2.4	2
118	Search for bioactive natural products from medicinal plants of Bangladesh. Journal of Natural Medicines, 2010, 64, 393-401.	2.3	17
119	Isolation and structure elucidation of flavonoid glycosides from Solanum verbascifolium. Phytochemistry Letters, 2010, 3, 88-92.	1.2	9
120	Isolation of new fuzanins, carbamate-containing natural products, from Kitasatospora sp. IFM10917. Journal of Antibiotics, 2010, 63, 385-388.	2.0	8
121	Practical Total Synthesis of Fuligocandins A and B. Synlett, 2010, 2010, 2498-2502.	1.8	10
122	A Bisindole Alkaloid with Hedgehog Signal Inhibitory Activity from the MyxomycetePerichaena chrysosperma. Journal of Natural Products, 2010, 73, 1711-1713.	3.0	26
123	Activity of Mangosteen Xanthones and Teleocidin A-2 in Death Receptor Expression Enhancement and Tumor Necrosis Factor Related Apoptosis-Inducing Ligand Assays. Journal of Natural Products, 2010, 73, 452-455.	3.0	18
124	Dehydrofuligoic Acid, a New Yellow Pigment Isolated from the Myxomycete Fuligo septica f. flava. Heterocycles, 2010, 82, 839.	0.7	8
125	Izumiphenazines Aâ^'C: Isolation and Structure Elucidation of Phenazine Derivatives from <i>Streptomyces</i> sp. IFM 11204. Journal of Natural Products, 2010, 73, 1999-2002.	3.0	75
126	Constituents of Amoora cucullata with TRAIL resistance-overcoming activity. Organic and Biomolecular Chemistry, 2010, 8, 3696.	2.8	35

#	Article	IF	CITATIONS
127	Terpenoids and a Flavonoid Clycoside from <i>Acacia pennata</i> Leaves as Hedgehog/GLI-Mediated Transcriptional Inhibitors. Journal of Natural Products, 2010, 73, 995-997.	3.0	59
128	Novel tryptophan metabolites, chromoazepinone A, B and C, produced by a blocked mutant of Chromobacterium violaceum, the biosynthetic implications and the biological activity of chromoazepinone A and B. Organic and Biomolecular Chemistry, 2010, 8, 3157.	2.8	7
129	Search for Bioactive Natural Products Targeting Cancer-Related Signaling Pathways. Yuki Gosei Kagaku Kyokaishi/Journal of Synthetic Organic Chemistry, 2009, 67, 1094-1104.	0.1	20
130	Cadinane Sesquiterpenes from Curcuma parviflora. Journal of Natural Products, 2009, 72, 782-783.	3.0	8
131	Fuligoic acid, a new yellow pigment with a chlorinated polyene–pyrone acid structure isolated from the myxomycete Fuligo septica f. flava. Tetrahedron Letters, 2009, 50, 3189-3190.	1.4	16
132	Isolation of new carbamate- or pyridine-containing natural products, fuzanins A, B, C, and D from Kitasatospora sp. IFM10917. Tetrahedron, 2009, 65, 369-373.	1.9	45
133	Cassaine diterpenoid dimers isolated from Erythrophleum succirubrum with TRAIL-resistance overcoming activity. Tetrahedron Letters, 2009, 50, 4658-4662.	1.4	25
134	Death receptor 5 targeting activity-guided isolation of isoflavones from Millettia brandisiana and Ardisia colorata and evaluation of ability to induce TRAIL-mediated apoptosis. Bioorganic and Medicinal Chemistry, 2009, 17, 1181-1186.	3.0	22
135	Death receptor 5 promoter-enhancing compounds isolated from Catimbium speciosum and their enhancement effect on TRAIL-induced apoptosis. Bioorganic and Medicinal Chemistry, 2009, 17, 6748-6754.	3.0	32
136	The first Hes1 dimer inhibitors from natural products. Bioorganic and Medicinal Chemistry Letters, 2009, 19, 5778-5781.	2.2	23
137	New Wnt/βâ€Catenin Signaling Inhibitors Isolated from <i>Eleutherine palmifolia</i> . Chemistry - an Asian Journal, 2009, 4, 540-547.	3.3	57
138	A Method for the Rapid Discovery of Naturally Occurring Products by Proteins Immobilized on Magnetic Beads and Reverse Affinity Chromatography. Chemistry - an Asian Journal, 2009, 4, 1802-1808.	3.3	20
139	Cardenolide Glycosides of <i>Thevetia peruviana</i> and Triterpenoid Saponins of <i>Sapindus emarginatus</i> as TRAIL Resistance-Overcoming Compounds. Journal of Natural Products, 2009, 72, 1507-1511.	3.0	18
140	Structure of New Monoterpene Glycoside from Sibiraea angustata RCHD. and Its Anti-obestic Effect. Chemical and Pharmaceutical Bulletin, 2009, 57, 294-297.	1.3	18
141	Cribrarione C, a Naphthoquinone Pigment from the Myxomycete Cribraria meylanii. Chemical and Pharmaceutical Bulletin, 2009, 57, 894-895.	1.3	12
142	Flavonoids from Eupatorium odoratum with Death Receptor 5 Promoter Enhancing Activity. Heterocycles, 2009, 77, 1379.	0.7	13
143	Quinic acid esters from <i>Pluchea indica</i> with collagenase, MMPâ€2 and MMPâ€9 inhibitory activities. Phytotherapy Research, 2008, 22, 264-266.	5.8	21
144	Studies on search for bioactive natural products targeting TRAIL signaling leading to tumor cell apoptosis. Medicinal Research Reviews, 2008, 28, 688-714.	10.5	84

#	Article	IF	CITATIONS
145	Naturally Occurring Smallâ€Molecule Inhibitors of Hedgehog/GLIâ€Mediated Transcription. ChemBioChem, 2008, 9, 1082-1092.	2.6	121
146	A novel sesquiterpenoid dimer parviflorene F induces apoptosis by up-regulating the expression of TRAIL-R2 and a caspase-dependent mechanism. Bioorganic and Medicinal Chemistry, 2008, 16, 1756-1763.	3.0	39
147	Hedgehog/GLI-mediated transcriptional inhibitors from Zizyphus cambodiana. Bioorganic and Medicinal Chemistry, 2008, 16, 9420-9424.	3.0	56
148	Determination of absolute stereochemistry, total synthesis, and evaluation of peptides from the myxomycete Physarum melleum. Bioorganic and Medicinal Chemistry Letters, 2008, 18, 95-98.	2.2	19
149	Efficient Synthesis of Chromone and Flavonoid Derivatives with Diverse Heterocyclic Units. Chemistry - an Asian Journal, 2008, 3, 2056-2064.	3.3	15
150	Acylated Triterpenoid Saponins from <i>Schima noronhae</i> and Their Cell Growth Inhibitory Activity. Journal of Natural Products, 2008, 71, 918-921.	3.0	21
151	Tyrosine Derivatives Isolated from <i>Streptomyces</i> sp. IFM 10937 in a Screening Program for TRAIL-Resistance-Overcoming Activity. Journal of Natural Products, 2008, 71, 1963-1966.	3.0	34
152	Synthesis and Evaluation of Isomers of Melleumin B from the Myxomycete Physarum melleum; an Approach for Wnt Signal Inhibitor Trusted in Nature. Heterocycles, 2008, 76, 1425.	0.7	9
153	Constituents from <i>Hoya parasitica</i> and their Cell Growth Inhibitory Activity. Planta Medica, 2008, 74, 760-763.	1.3	13
154	Mangiferin Identified in a Screening Study Guided by Neuraminidase Inhibitory Activity. Planta Medica, 2007, 73, 1195-1196.	1.3	18
155	First isolation of sesquiterpenes and flavonoids fromZingiber spectabileand identification of zerumbone as the major cell growth inhibitory component. Natural Product Research, 2007, 21, 1242-1247.	1.8	23
156	Brandisianins Aâ^'F, Isoflavonoids Isolated from <i>Millettia brandisiana</i> in a Screening Program for Death-Receptor Expression Enhancement Activity. Journal of Natural Products, 2007, 70, 1910-1914.	3.0	38
157	Synthesis, determination of stereochemistry, and evaluation of new bisindole alkaloids from the myxomycete Arcyria ferruginea: An approach for Wnt signal inhibitor. Bioorganic and Medicinal Chemistry Letters, 2007, 17, 4254-4257.	2.2	50
158	A novel natural compound, a cycloanthranilylproline derivative (Fuligocandin B), sensitizes leukemia cells to apoptosis induced by tumor necrosis factor–related apoptosis-inducing ligand (TRAIL) through 15-deoxy-î"12, 14 prostaglandin J2 production. Blood, 2007, 110, 1664-1674.	1.4	55
159	Lignan glycosides and flavonoids from Saraca asoca with antioxidant activity. Journal of Natural Medicines, 2007, 61, 480-482.	2.3	65
160	Secoiridoid components from Jasminum grandiflorum. Phytochemistry, 2007, 68, 1718-1721.	2.9	24
161	Parviflorene J, a Cytotoxic Sesquiterpene Dimer with a New Rearranged Skeleton from Curcuma parviflora. Heterocycles, 2007, 72, 649.	0.7	15
162	New Kehokorins and Trichiols from the Myxomycete Trichia favoginea. Heterocycles, 2007, 71, 1807.	0.7	14

#	Article	IF	CITATIONS
163	Cell Growth and Cell Cycle Inhibitory Activities of 20-Epidiosgenyl Saponin from Calamus insignis. Heterocycles, 2007, 74, 931.	0.7	0
164	Bisindole Alkaloids from MyxomycetesArcyriadenudataandArcyriaobvelata. Journal of Natural Products, 2006, 69, 1252-1254.	3.0	27
165	Diterpenes fromLeucasasperalnhibiting Prostaglandin-Induced Contractions. Journal of Natural Products, 2006, 69, 988-994.	3.0	37
166	Prostaglandin inhibitory and antioxidant components of Cistus laurifolius, a Turkish medicinal plant. Journal of Ethnopharmacology, 2006, 108, 371-378.	4.1	63
167	Nine Constituents Including Six Xanthone-Related Compounds Isolated from Two Ascomycetes, Gelasinospora santi-florii and Emericella quadrilineata, Found in a Screening Study Focused on Immunomodulatory Activity. Chemical and Pharmaceutical Bulletin, 2006, 54, 550-553.	1.3	39
168	Dihydroflavonol BB-1, an extract of natural plant Blumea balsamifera, abrogates TRAIL resistance in leukemia cells. Blood, 2006, 107, 679-688.	1.4	65
169	Steroidal saponins from Calamus insignis, and their cell growth and cell cycle inhibitory activities. Bioorganic and Medicinal Chemistry, 2006, 14, 659-665.	3.0	24
170	Kehokorins A–C, novel cytotoxic dibenzofurans isolated from the myxomycete Trichia favoginea var. persimilis. Tetrahedron Letters, 2006, 47, 1505-1508.	1.4	44
171	Trichiol and 3-epitrichiol acetate, novel cytotoxic sterols with an unprecedented 2,6-dioxabicyclo[2.2.2]octan-3-one ring system from the myxomycete Trichia favoginea var. persimilis. Tetrahedron Letters, 2006, 47, 4351-4354.	1.4	3
172	Flavonoids and sesquiterpenoids, constituents from Eupatorium capillifolium, found in a screening study guided by cell growth inhibitory activity. Journal of Natural Medicines, 2006, 60, 325-328.	2.3	5
173	First isolation of three diterpenes as naturally-occurring compounds from Sindora siamensis. Journal of Natural Medicines, 2006, 61, 77-79.	2.3	7
174	Flavonoids from Sonneratia caseolaris. Journal of Natural Medicines, 2006, 60, 264-265.	2.3	49
175	A new lignan from Aphanamixis polystachya. Journal of Natural Medicines, 2006, 60, 258-260.	2.3	36
176	Tryptophan 4-Aminocinnamamides from a Myxomycete Fuligo aurea. Heterocycles, 2006, 69, 463.	0.7	7
177	New Bisindole Alkaloids Isolated from Myxomycetes Arcyria cinerea and Lycogala epidendrum. Chemical and Pharmaceutical Bulletin, 2005, 53, 594-597.	1.3	23
178	Eleven New 2-Pyrones from a Fungi Imperfecti, Trichurus terrophilus, Found in a Screening Study Guided by Immunomodulatory Activity. Chemical and Pharmaceutical Bulletin, 2005, 53, 923-929.	1.3	14
179	Melleumin A, a novel peptide lactone isolated from the cultured myxomycete Physarum melleum. Tetrahedron Letters, 2005, 46, 267-271.	1.4	25
180	Total synthesis of TT-1 (rasfonin), an α-pyrone-containing natural product from a fungus Trichurus terrophilus. Tetrahedron, 2005, 61, 1827-1833.	1.9	16

#	Article	IF	CITATIONS
181	Cytotoxic dimeric sesquiterpenoids from Curcuma parviflora: isolation of three new parviflorenes and absolute stereochemistry of parviflorenes A, B, D, F, and G. Tetrahedron, 2005, 61, 6700-6706.	1.9	16
182	Pyrroloiminoquinone pigments from Didymium iridis. Biochemical Systematics and Ecology, 2005, 33, 323-325.	1.3	12
183	New cytotoxic bisindole alkaloids with protein tyrosine kinase inhibitory activity from a myxomycete Lycogala epidendrum. Bioorganic and Medicinal Chemistry Letters, 2005, 15, 2776-2780.	2.2	32
184	Isolation of Bioactive Natural Products from Myxomycetes. Medicinal Chemistry, 2005, 1, 575-590.	1.5	17
185	Sesquiterpenoids and Plasmin-Inhibitory Flavonoids from Blumea balsamifera. Journal of Natural Products, 2005, 68, 447-449.	3.0	41
186	Collagenase Inhibitory Quinic Acid Esters fromIpomoeapes-caprae. Journal of Natural Products, 2005, 68, 794-796.	3.0	46
187	Isolation of Austroinulin Possessing Cell Cycle Inhibition Activity fromBlumea glomerataand Revision of Its Absolute Configuration. Planta Medica, 2004, 70, 1170-1173.	1.3	16
188	Parviflorenes B–F, novel cytotoxic unsymmetrical sesquiterpene-dimers with three backbone skeletons from Curcuma parviflora. Tetrahedron, 2004, 60, 10817-10824.	1.9	31
189	New chlorogenin hexasaccharide isolated from Agave fourcroydes with cytotoxic and cell cycle inhibitory activities. Bioorganic and Medicinal Chemistry, 2004, 12, 3841-3845.	3.0	47
190	Tubiferal A, a backbone-rearranged triterpenoid lactone isolated from the myxomycete Tubifera dimorphotheca, possessing reversal of drug resistance activity. Tetrahedron, 2004, 60, 9835-9839.	1.9	20
191	Immunomodulatory Constituents from an Ascomycete,Chaetomiumseminudum. Journal of Natural Products, 2004, 67, 98-102.	3.0	72
192	Six Immunosuppressive Features from an Ascomycete, Zopfiella ion gicaudata, Found in a Screening Study Monitored by Immunomodulatory Activity. Chemical and Pharmaceutical Bulletin, 2004, 52, 1005-1008.	1.3	80
193	Cycloanthranilylproline-Derived Constituents from a Myxomycete Fuligo candida. Chemical and Pharmaceutical Bulletin, 2004, 52, 368-370.	1.3	38
194	Dihydroflavonol BB-1, an Extract of Natural Plant Blumea Balsamifera, Abrogates TRAIL Resistance in Adult T-Cell Leukemia/Lymphoma Blood, 2004, 104, 2290-2290.	1.4	0
195	Development of Unexploited Microbial Resources: Natural Products from Slime Molds. ChemInform, 2003, 34, no.	0.0	Ο
196	Cribrarione A, a New Antimicrobial Naphthoquinone Pigment from a Myxomycete Cribraria purpurea ChemInform, 2003, 34, no.	0.0	0
197	Physarigins A—C, Three New Yellow Pigments from a Cultured Myxomycete Physarum rigidum ChemInform, 2003, 34, no.	0.0	0
198	Absolute stereochemistry of TT-1 (rasfonin), an α-pyrone-containing natural product from a fungus, Trichurus terrophilus. Tetrahedron Letters, 2003, 44, 8427-8431.	1.4	17

#	Article	IF	CITATIONS
199	Parviflorene A, a novel cytotoxic unsymmetrical sesquiterpene–dimer constituent from Curcuma parviflora. Tetrahedron Letters, 2003, 44, 2327-2329.	1.4	19
200	Physarigins A–C, three new yellow pigments from a cultured myxomycete Physarum rigidum. Tetrahedron Letters, 2003, 44, 4479-4481.	1.4	19
201	Cribrarione A, a new antimicrobial naphthoquinone pigment from a myxomycete Cribraria purpurea. Tetrahedron, 2003, 59, 3433-3435.	1.9	25
202	Isolation of bisindole alkaloids that inhibit the cell cycle from Myxomycetes Arcyria ferruginea and Tubifera casparyi. Bioorganic and Medicinal Chemistry Letters, 2003, 13, 2879-2881.	2.2	41
203	Sterol and Triterpenoid Constituents ofVerbenalittoraliswith NGF-Potentiating Activity. Journal of Natural Products, 2003, 66, 696-698.	3.0	47
204	Cribrarione B, a New Naphthoquinone Pigment from the Myxomycete Cribraria cancellata. Journal of Natural Products, 2003, 66, 1611-1612.	3.0	23
205	Dihydrolindbladiones, Three New Naphthoquinone Pigments from a MyxomyceteLindbladia tubulina. Journal of Natural Products, 2003, 66, 999-1001.	3.0	19
206	Discovery of New Macrolides from Marine Organisms. , 2003, , 57-98.		0
207	Bahiensol, a New Glycerolipid from a Cultured Myxomycete Didymium bahiense var. bahiense. Chemical and Pharmaceutical Bulletin, 2003, 51, 612-613.	1.3	15
208	A New Iridoid Glycoside with Nerve Growth Factor-Potentiating Activity, Gelsemiol 6'-trans-Caffeoyl-1-glucoside, from Verbena littoralis. Chemical and Pharmaceutical Bulletin, 2003, 51, 1103-1105.	1.3	26
209	Prenylated Xanthones from Garcinia xanthochymus. Chemical and Pharmaceutical Bulletin, 2003, 51, 1332-1334.	1.3	27
210	Six New Constituents from an Ascomycete, Chaetomium quadrangulatum, Found in a Screening Study Focused on Monoamine Oxidase Inhibitory Activity Chemical and Pharmaceutical Bulletin, 2003, 51, 247-251.	1.3	26
211	Separation of Leucas aspera, a Medicinal Plant of Bangladesh, Guided by Prostaglandin Inhibitory and Antioxidant Activities. Chemical and Pharmaceutical Bulletin, 2003, 51, 595-598.	1.3	144
212	Wrightiamines A and B, Two New Cytotoxic Pregnane Alkaloids from Wrightia javanica. Chemical and Pharmaceutical Bulletin, 2003, 51, 737-739.	1.3	16
213	Littorachalcone, a New Enhancer of NGF-Mediated Neurite Outgrowth, from Verbena littoralis. Chemical and Pharmaceutical Bulletin, 2003, 51, 872-874.	1.3	29
214	Development of Unexploited Microbial Resources: Natural Products from Slime Molds Yuki Gosei Kagaku Kyokaishi/Journal of Synthetic Organic Chemistry, 2003, 61, 152-163.	0.1	6
215	Lindbladione and Related Naphthoquinone Pigments from a Myxomycete <i>Lindbladia tubulina</i> . Chemical and Pharmaceutical Bulletin, 2002, 50, 1126-1127.	1.3	25
216	Five New Chromones Possessing Monoamine Oxidase Invibitor Activity from an Ascomycete, Chaetomium quadrangulatum Chemical and Pharmaceutical Bulletin, 2002, 50, 330-336.	1.3	41

#	Article	IF	CITATIONS
217	New Onoceranoid Triterpene Constituents from Lansium domesticum. Journal of Natural Products, 2002, 65, 1709-1711.	3.0	41
218	Alkenylphenol and alkenylsalicylic acid from Mangifera caesia. Biochemical Systematics and Ecology, 2002, 30, 475-478.	1.3	5
219	5α,8α-Epidioxysterol sulfate from a diatom Odontella aurita. Phytochemistry, 2002, 61, 359-360.	2.9	9
220	Littoralisone, a Novel Neuritogenic Iridolactone Having an Unprecedented Heptacyclic Skeleton Including Four- and Nine-Membered Rings Consisting of Glucose fromVerbena littoralis. Journal of Organic Chemistry, 2001, 66, 2165-2167.	3.2	45
221	New Multiflorane-Type Triterpenoid Acids fromSandoricum indicum. Journal of Natural Products, 2001, 64, 1243-1245.	3.0	19
222	Immunomodulatory Constituents from an Ascomycete,Eupenicillium crustaceum, and Revised Absolute Structure of Macrophorin D. Journal of Natural Products, 2001, 64, 1234-1237.	3.0	50
223	Dendocarbins Aâ^'N, New Drimane Sesquiterpenes from the Nudibranch Dendrodoris carbunculosa. Journal of Natural Products, 2001, 64, 726-731.	3.0	43
224	Laboratory Culture of the Myxomycetes:  Formation of Fruiting Bodies of Didymium bahiense and Its Plasmodial Production of Makaluvamine A. Journal of Natural Products, 2001, 64, 108-110.	3.0	56
225	Analgesic Components of Saposhnikovia Root (Saposhnikovia divaricata) Chemical and Pharmaceutical Bulletin, 2001, 49, 154-160.	1.3	121
226	Immunomodulatory Constituents from an Ascomycete, Emericella aurantio-brunnea Chemical and Pharmaceutical Bulletin, 2000, 48, 1436-1441.	1.3	33
227	Colopsinols D and E, New Polyhydroxyl Linear Carbon Chain Compounds from Marine Dinoflagellate Amphidinium sp Chemical and Pharmaceutical Bulletin, 2000, 48, 1447-1451.	1.3	22
228	Bioactive secondary metabolites from Okinawan sponges and tunicates. Studies in Natural Products Chemistry, 2000, 23, 185-231.	1.8	5
229	Casein kinase II inhibitors isolated form two Brazilian plants Hymenaea parvifolia and Wulffia baccata. Bioorganic and Medicinal Chemistry Letters, 1999, 9, 2157-2160.	2.2	10
230	Colopsinols B and C, new long chain polyhydroxy compounds from cultured marine dinoflagellate Amphidinium sp Journal of the Chemical Society Perkin Transactions 1, 1999, , 3483-3487.	0.9	24
231	Amphidinolide B, a powerful activator of actomyosin ATPase enhances skeletal muscle contraction. Biochimica Et Biophysica Acta - General Subjects, 1999, 1427, 24-32.	2.4	13
232	Marine Natural Products and Marine Chemical Ecology. , 1999, , 415-649.		27
233	Colopsinol A, a Novel Polyhydroxyl Metabolite from Marine DinoflagellateAmphidiniumsp Journal of Organic Chemistry, 1999, 64, 1478-1482.	3.2	41
234	Monoamine Oxidase Inhibitory Naphthoquinone and/or Naphthalene Dimers from Lemuni Hitam (Diospyros sp.), a Malaysian Herbal Medicine Chemical and Pharmaceutical Bulletin, 1999, 47, 1473-1476.	1.3	9

#	Article	IF	CITATIONS
235	Bioactive products from marine micro- and macro-organisms. Pure and Applied Chemistry, 1999, 71, 1123-1126.	1.9	62
236	Studies on stereochemistry of theonezolides A â^¼ C: Determination of absolute stereochemistry of the C-4 â^¼ C-17 fragment. Tetrahedron, 1998, 54, 4819-4826.	1.9	7
237	9-Methyl-7-bromoeudistomin D induces Ca2+ release from cardiac sarcoplasmic reticulum. European Journal of Pharmacology, 1998, 357, 261-265.	3.5	5
238	New polyketide peroxides from Okinawan marine sponge Plakortis sp Tetrahedron, 1998, 54, 2041-2048.	1.9	20
239	Luteophanols B and C, new polyhydroxyl compounds from marine dinoflagellate Amphidinium sp Tetrahedron, 1998, 54, 14455-14464.	1.9	59
240	New Cyclic Polyketide Peroxides from Okinawan Marine SpongePlakortissp Journal of Natural Products, 1998, 61, 1427-1429.	3.0	11
241	9-Methyl-7-bromoeudistomin D, a powerful caffeine-like Ca2+ releaser, acts on Ca2+-induced Ca2+ release channels of cardiac sarcoplasmic reticulum. The Japanese Journal of Pharmacology, 1998, 76, 204.	1.2	0
242	Studies on the Stereochemistry of Theonezolides A-C: Elucidation of the Relative Configurations of 1,3-Diol Moieties of the C-4Ã`C-17 Fragment. Heterocycles, 1998, 49, 39.	0.7	7
243	Amphidinolides: Unique Macrolides from Marine Dinoflagellates. Heterocycles, 1997, 44, 543.	0.7	95
244	Topsentinols A-J, New Sterols with Highly Branched Side Chains from Marine Sponge Topsentia sp Chemical and Pharmaceutical Bulletin, 1997, 45, 1435-1438.	1.3	27
245	Taurospongin A, a Novel Acetylenic Fatty Acid Derivative Inhibiting DNA Polymerase β and HIV Reverse Transcriptase from SpongeHippospongiasp Journal of Organic Chemistry, 1997, 62, 3831-3836.	3.2	67
246	Luteophanol A, a New Polyhydroxyl Compound from Symbiotic Marine DinoflagellateAmphidiniumsp Journal of Organic Chemistry, 1997, 62, 3820-3823.	3.2	70
247	Studies on the macrolides from marine dinoflagellate Amphidinium sp.: Structures of amphidinolides R and S and a succinate feeding experiment. Tetrahedron, 1997, 53, 7827-7832.	1.9	36
248	Studies on the Stereochemistry of Amphidinolides: Synthesis of a Diastereomer of the C-1-C-9 Fragment of Amphidinolide C Chemical and Pharmaceutical Bulletin, 1996, 44, 1819-1822.	1.3	31
249	Total synthesis of pseudodistomin C, a sphingosine-related piperidine alkaloid from tunicate Pseudodistoma kanoko. Tetrahedron, 1996, 52, 4573-4580.	1.9	28
250	Amphidinolide Q, a novel 12-membered macrolide from the cultured marine dinoflagellate Amphidinium sp. Tetrahedron Letters, 1996, 37, 1449-1450.	1.4	32
251	Absolute stereochemistry of penaresidins A and B. Tetrahedron Letters, 1996, 37, 6775-6776.	1.4	44
252	The mode of rabbit platelet shape change and aggregation induced by theonezolide-A, a novel polyketide macrolide, isolated from the Okinawan marine sponge <i>Theonella</i> sp Canadian Journal of Physiology and Pharmacology, 1996, 74, 193-199.	1.4	17

#	Article	IF	CITATIONS
253	Sphingosine-related Marine Alkaloids: Cyclic Amino Alcohols. Heterocycles, 1996, 42, 943.	0.7	46
254	The mode of rabbit platelet shape change and aggregation induced by theonezolide-A, a novel polyketide macrolide, isolated from the Okinawan marine sponge <i>Theonella</i> sp Canadian Journal of Physiology and Pharmacology, 1996, 74, 193-199.	1.4	0
255	Three new oxylipins related to 3,6-dioxo-4-docosenoic acid from Okinawan marine sponges, Plakortis spp. Tetrahedron, 1995, 51, 5979-5986.	1.9	31
256	Structure-activity relationship of bromoeudistomin D, a powerful Ca2+ releaser in skeletal muscle sarcoplasmic reticulum. European Journal of Pharmacology, 1995, 288, 285-293.	2.6	15
257	Revised Structure of Pseudodistomin A, a Piperidine Alkaloid Isolated from the Okinawan Tunicate Pseudodistoma kanoko. Journal of Natural Products, 1995, 58, 804-806.	3.0	27
258	Pseudodistomin C, a new piperidine alkaloid with unusual absolute configuration from the Okinawan tunicate Pseudodistoma kanoko. Journal of Organic Chemistry, 1995, 60, 6941-6945.	3.2	49
259	Amphidinolides O and P, novel 15-membered macrolides from the dinoflagellate Amphidinium sp.: Analysis of the relative stereochemistry and stable solution conformation. Journal of Organic Chemistry, 1995, 60, 6062-6066.	3.2	43
260	Isolation of Apo-9'-fucoxanthinone from the Cultured Marine Dinoflagellate Amphidinium sp Journal of Natural Products, 1995, 58, 1097-1099.	3.0	28
261	Biosynthetic studies of amphidinolide J: explanation of the generation of the unusual odd-numbered macrocyclic lactone. Journal of the Chemical Society Chemical Communications, 1995, , 1639.	2.0	27
262	4, 6â€Dibromoâ€3â€hydroxycarbazole (an analogue of caffeineâ€like Ca ²⁺ releaser), a novel type of inhibitor of Ca ²⁺ â€induced Ca ²⁺ release in skeletal muscle sarcoplasmic reticulum. British Journal of Pharmacology, 1995, 114, 941-948.	5.4	10
263	Stereochemistry of the Modified Tryptophan Residues Contained in Konbamide and Keramamide a Isolated from Marine Sponges of the Genus <i>Theonella</i> . Natural Product Research, 1994, 4, 293-296.	0.4	12
264	Convolutamides A â^¼ F, novel γ-lactam alkaloids from the marine bryozoan Amathia convoluta. Tetrahedron, 1994, 50, 10201-10206.	1.9	65
265	Isolation and structure of shimofuridins B â^¼ G from the Okinawan marine tunicate Aplidium multiplicatum. Tetrahedron, 1994, 50, 8651-8656.	1.9	23
266	Isolation and structures of Theonezolides B and C from the Okinawan marine sponge Theonella sp Tetrahedron, 1994, 50, 8355-8362.	1.9	27
267	Amphidinin A, a novel amphidinolide-related metabolite from the cultured marine dinoflagellate Amphidinium sp Tetrahedron Letters, 1994, 35, 7049-7050.	1.4	29
268	Absolute stereochemistry of amphidinolide B. Tetrahedron Letters, 1994, 35, 8241-8242.	1.4	42
269	Two New Tryptophan-Derived Alkaloids from the Okinawan Marine Sponge Aplysina Sp Journal of Natural Products, 1994, 57, 1008-1011.	3.0	50
270	Shimofuridin A, a nucleoside derivative embracing an acylfucopyranoside unit isolated from the Okinawan marine tunicate Aplidium multiplicatum. Journal of Organic Chemistry, 1994, 59, 255-257.	3.2	32

#	Article	IF	CITATIONS
271	Amphidinolide N, a novel 26-membered macrolide with remarkably potent cytotoxicity from the cultured marine dinoflagellate Amphidinium sp Journal of the Chemical Society Chemical Communications, 1994, , 1455.	2.0	35
272	Nepheliosyne A, New C47 Acetylenic Acid from the Okinawan Marine Sponge Xestospongia sp Journal of Natural Products, 1994, 57, 1300-1303.	3.0	20
273	Plakoridine A, a new tyramine-containing pyrrolidine alkaloid from the Okinawan marine sponge Plakortis sp Journal of Organic Chemistry, 1994, 59, 3712-3713.	3.2	33
274	Amphidinolide M, a Novel 29-Membered Macrolide from the Cultured Marine Dinoflagellate Amphidinium sp Journal of Organic Chemistry, 1994, 59, 4698-4700.	3.2	15
275	Two new polycyclic aromatic alkaloids from the Okinawan marine sponge Biemna sp Tetrahedron, 1993, 49, 8337-8342.	1.9	30
276	Manzamenones G and H, new dimeric fatty-acid derivatives from the Okinawan marine sponge Plakortis sp Tetrahedron, 1993, 49, 5955-5960.	1.9	21
277	Untenone A, a new cyclopentenone from the Okinawan marine sponge Plakortis sp. corresponding to the dienophile in the biosynthesis of manzamenones. Tetrahedron Letters, 1993, 34, 3749-3750.	1.4	44
278	Bioactive metabolites of symbiotic marine microorganisms. Chemical Reviews, 1993, 93, 1753-1769.	47.7	344
279	Keruffaride, a new all-ciscyclopentanepentol-containing metabolite from the okinawan marine sponge luffariella sp Journal of the Chemical Society Chemical Communications, 1993, , 79.	2.0	7
280	Amphidinolide K, a new 19-membered macrolide from the cultured symbiotic dinoflagellate Amphidinium sp. Journal of Organic Chemistry, 1993, 58, 6928-6929.	3.2	28
281	Amphidinolide J: A cytotoxic macrolide from the marine dinoflagellate Amphidinium sp. Determination of the absolute stereochemistry. Journal of Organic Chemistry, 1993, 58, 2645-2646.	3.2	39
282	Theonezolide a: a novel polyketide natural product from the Okinawan marine sponge Theonella sp. Journal of the American Chemical Society, 1993, 115, 6661-6665.	13.7	47
283	Keruffaride: Structure Revision and Isolation from Plural Genera of Okinawan Marine Sponges. Journal of Natural Products, 1993, 56, 1856-1860.	3.0	26
284	Luffariolides F and G, New Manoalide Derivatives from the Okinawan Marine Sponge Luffariella Sp Journal of Natural Products, 1993, 56, 436-439.	3.0	26
285	Biemnasterol, a New Cytotoxic Sterol with the Rare 22,25-Diene Side Chain, Isolated from the Marine Sponge Biemna sp Journal of Natural Products, 1993, 56, 2016-2018.	3.0	17
286	The Stereochemistry of Variabilin, an Antimicrobial Sesterterpene Isolated from Marine Sponges. Natural Product Research, 1993, 3, 189-192.	0.4	15
287	Inhibition of Mammalian Topoisomerase I by Xestoquinone and Halenaquinone. Bioscience, Biotechnology and Biochemistry, 1993, 57, 330-331.	1.3	10
288	New Sesterterpenes with Nerve Growth Factor Synthesis-Stimulating Activity from the Okinawan Marine Sponge Hyrtios sp Chemical and Pharmaceutical Bulletin, 1993, 41, 2190-2191.	1.3	20

#	Article	IF	CITATIONS
289	Chapter 2 Marine Alkaloids II. Alkaloids: Chemistry and Pharmacology, 1992, , 41-124.	0.2	10
290	SO-75R1, a new mutactimycin derivative produced by Nocardia brasiliensis Journal of Antibiotics, 1992, 45, 995-997.	2.0	22
291	Agelasine G, a New Antileukemic Alkaloid from the Okinawan Marine Sponge Agelas sp Chemical and Pharmaceutical Bulletin, 1992, 40, 766-767.	1.3	55
292	Purealidins E-G, New Bromotyrosine Alkaloids from the Okinawan Marine Sponge Psammaplysilla purea. Journal of Natural Products, 1992, 55, 1325-1327.	3.0	23
293	Niphatesines E–H, new pyridine alkaloids from the Okinawan marine sponge Niphates sp Journal of the Chemical Society Perkin Transactions 1, 1992, , 1291-1294.	0.9	39
294	Ircinals A and B from the Okinawan marine sponge Ircinia sp.: plausible biogenetic precursors of manzamine alkaloids. Journal of Organic Chemistry, 1992, 57, 2480-2483.	3.2	161
295	Manzamenones A-F from the Okinawan marine sponges Plakortis Sp.: novel dimeric fatty acid derivatives possessing a bicyclo[4.3.0]nonane skeleton. Journal of Organic Chemistry, 1992, 57, 5255-5260.	3.2	48
296	Luffariolides A-E, new cytotoxic sesterterpenes from the Okinawan marine sponge Luffariella sp. Journal of Organic Chemistry, 1992, 57, 3503-3507.	3.2	44
297	New Pentacyclic Compounds from the Okinawan Marine Sponge Xestospongia sapra. Journal of Natural Products, 1992, 55, 994-998.	3.0	29
298	Theoneberine: the first brominated benzyltetrahydroprotoberberine alkaloid from the Okinawan marine sponge Theonella sp. Journal of Organic Chemistry, 1992, 57, 6680-6682.	3.2	30
299	Keramamide F, a new thiazole-containing peptide from the Okinawan marine sponge Theonella sp. Journal of Organic Chemistry, 1992, 57, 5540-5542.	3.2	69
300	Aplysidine, a new nucleoside from the Okinawan marine sponge Aplysina Sp Tetrahedron, 1992, 48, 7145-7148.	1.9	22
301	Pseudodistomin B: Revised structure and first total synthesis. Tetrahedron Letters, 1992, 33, 7389-7390.	1.4	29
302	Plakotenin, a new cytotoxic carboxylic acid from the okinawan marine sponge plakortis Sp Tetrahedron Letters, 1992, 33, 2579-2580.	1.4	27
303	Purealidin D, a new pyridine alkaloid from the okinawan marine sponge psammaplysilla purea. Tetrahedron Letters, 1992, 33, 2597-2598.	1.4	30
304	New congeners of swinholides from the okinawan marine sponge Theonella sp Journal of the Chemical Society Perkin Transactions 1, 1991, , 3185.	0.9	37
305	Keramamide A, a novel peptide from the Okinawan marine sponge Theonella sp Journal of the Chemical Society Perkin Transactions 1, 1991, , 2609.	0.9	55
306	Amphidinolides G and H: new potent cytotoxic macrolides from the cultured symbiotic dinoflagellate Amphidinium sp. Journal of Organic Chemistry, 1991, 56, 5221-5224.	3.2	87

#	Article	IF	CITATIONS
307	Konbamide, a novel peptide with calmoduiin antagonistic activity from the Okinawan marine sponge Theonella sp Journal of the Chemical Society Chemical Communications, 1991, , 1050-1052.	2.0	54
308	New congeners of bistheonellides from Okinawan marine sponges of the genus Theonella. Journal of the Chemical Society Perkin Transactions 1, 1991, , 2379.	0.9	20
309	Cystodytins D-I, New Cytotoxic Tetracyclic Aromatic Alkaloids from the Okinawan Marine Tunicate Cystodytes dellechiajei. Journal of Natural Products, 1991, 54, 1634-1638.	3.0	52
310	1H- and 13C-nmr Spectral Investigation on Amphidinolide A, an Antileukemic Marine Macrolide. Journal of Natural Products, 1991, 54, 1435-1439.	3.0	26
311	Keramamides B .apprx. D, novel peptides from the Okinawan marine sponge Theonella sp. Journal of the American Chemical Society, 1991, 113, 7812-7813.	13.7	70
312	Manzacidins A-C, novel tetrahydropyrimidine alkaloids from the Okinawan marine sponge Hymeniacidon sp. Journal of Organic Chemistry, 1991, 56, 4574-4576.	3.2	136
313	Penaresidin A and B, two novel azetidine alkaloids with potent actomyosin ATPase-activating activity from the Okinawan marine sponge Penares sp Journal of the Chemical Society Perkin Transactions 1, 1991, , 1135-1137.	0.9	114
314	Amphidinolide F, a new cytotoxic macrolide from the marine dinoflagellate Amphidinium sp Journal of Antibiotics, 1991, 44, 1259-1261.	2.0	68
315	Novel antibiotics, furaquinocins C, D, E, F, G and H Journal of Antibiotics, 1991, 44, 390-395.	2.0	60
316	A new antibiotic, okicenone. I. Taxonomy, fermentation, isolation and biological characteristics Journal of Antibiotics, 1991, 44, 814-818.	2.0	9
317	A new antibiotic, okicenone. II. Physico-chemical properties and structure elucidation Journal of Antibiotics, 1991, 44, 819-823.	2.0	12
318	Eudistomidins E and F, new β-carboline alkaloids from the okinawan marine tunicate Eudistoma glaucus. Tetrahedron Letters, 1991, 32, 3539-3542.	1.4	42
319	Purealidin A, a new cytotoxic bromotyrosine-derived alkaloid from the Okinawan marine spongePsammaplysilla purea. Experientia, 1991, 47, 299-300.	1.2	38
320	Stelliferins A–F, new antineoplastic isomalabaricane triterpenes from the Okinawan marine sponge Jaspis stellifera Tetrahedron, 1991, 47, 2181-2194.	1.9	56
321	lejimalides C and D, new antineoplastic 24-membered macrolide sulfates from the okinawan marine tunicate Eudistoma cf. rigida. Tetrahedron Letters, 1991, 32, 797-798.	1.4	65
322	Revised structures of prianosins C and D, antineoplastic alkaloids from the Okinawan marine sponge Prianos melanos. Tetrahedron Letters, 1991, 32, 1227-1228.	1.4	49
323	Purealidins B and C, new bromotyrosine alkaloids from the okinawan marine sponge psammaplysilla purea. Tetrahedron, 1991, 47, 6617-6622.	1.9	193
324	1-(α-Diazobenzyl)pyrene: A Reagent for Photolabile and Fluorescent Protection of Carboxyl Groups of Amino Acids and Peptides. Synlett, 1991, 1991, 35-36.	1.8	17

#	Article	IF	CITATIONS
325	Novel antibiotics, furaquinocins A and B. Taxonomy, fermentation, isolation and physico-chemical and biological characteristics Journal of Antibiotics, 1990, 43, 247-252.	2.0	61
326	Ageliferins, potent actomyosin Atpase activators from the Okinawan marine sponge sp. Tetrahedron, 1990, 46, 5579-5586.	1.9	122
327	Hyrtiosins A and B, new indole alkaloids from the Okinawan marine sponge Hyrtios erecta. Tetrahedron, 1990, 46, 7699-7702.	1.9	88
328	Rigidin, a novel alkaloid with calmodulin antagonistic activity from the okinawan marine tunicate Eudistoma cf. rigida. Tetrahedron Letters, 1990, 31, 4617-4620.	1.4	68
329	Niphatesines A–D, new antineoplastic pyridine alkaloids from the okinawan marine sponge Niphates sp Journal of the Chemical Society Perkin Transactions 1, 1990, , 3301-3303.	0.9	43
330	Amphidinolide E, a novel antileukemic 19-membered macrolide from the cultured symbiotic dinoflagellate Amphidinium sp. Journal of Organic Chemistry, 1990, 55, 3421-3423.	3.2	52
331	Biosynthesis of furaquinocins A and B. Journal of Organic Chemistry, 1990, 55, 1132-1133.	3.2	59
332	Bioactive Substances from Marine Organisms. Journal of Fiber Science and Technology, 1990, 46, P283-P287.	0.0	0
333	9-Methyl-7-bromoeudistomin D, a potent inducer of calcium release from sarcoplasmic reticulum of skeletal muscle. Experientia, 1989, 45, 782-783.	1.2	31
334	Structures of novel antibiotics, furaquinocins A and B. Tetrahedron Letters, 1989, 30, 7427-7430.	1.4	33
335	Cytotoxic Macrolides from a Cultured Marine Dinoflagellate of the Genus Amphidinium. Journal of Natural Products, 1989, 52, 1036-1041.	3.0	50
336	Hymenosulphate, a novel sterol sulphate with Ca-releasing activity from the cultured marine haptophyte Hymenomonas sp. Journal of the Chemical Society Perkin Transactions 1, 1989, , 101.	0.9	24
337	Novel cytocidal antibiotics, glucopiericidinols A1 and A2 Taxonomy, fermentation, isolation, structure elucidation and biological characteristics Journal of Antibiotics, 1989, 42, 1734-1740.	2.0	16
338	Symbioramide, a novel Ca2+-ATPase activator from the cultured dinoflagellateSymbiodinium sp Experientia, 1988, 44, 800-802.	1.2	51
339	Metachromins A and B, novel antineoplastic sesquiterpenoids from the Okinawan sponge Hippospongia cf. metachromia. Journal of Organic Chemistry, 1988, 53, 2855-2858.	3.2	55
340	Amphidinolide C: the first twenty-five membered macrocyclic lactone with potent antineoplastic activity from the cultured dinoflagellate Amphidinium sp. Journal of the American Chemical Society, 1988, 110, 490-494.	13.7	101
341	Bitter principles of Ailanthus altissima Swingle. Studies on oxidative degradation of the A-ring of ailanthone derivatives Chemical and Pharmaceutical Bulletin, 1987, 35, 3011-3015.	1.3	4
342	Amphidinolide-B, a novel macrolide with potent antineoplastic activity from the marine dinoflagellate Amphidinium sp Journal of the Chemical Society Chemical Communications, 1987, , 1127.	2.0	45

#	Article	IF	CITATIONS
343	Pseudodistomins A and B, novel antineoplastic piperdine alkaloids with calmodulin antagonistic activity from the Okinawan tunicate Pseudodistoma kanoko. Journal of Organic Chemistry, 1987, 52, 450-453.	3.2	79
344	Prianosin A, a novel antileukemic alkaloid from the okinawan marine sponge Prianos melanos. Tetrahedron Letters, 1987, 28, 4939-4942.	1.4	102
345	Scytophycins, cytotoxic and antimycotic agents from the cyanophyte Scytonema pseudohofmanni. Journal of Organic Chemistry, 1986, 51, 5300-5306.	3.2	210
346	Amphidinolide-A, a novel antineoplastic macrolide from the marine dinoflagellate Amphidinium sp Tetrahedron Letters, 1986, 27, 5755-5758.	1.4	83
347	Bitter Principles ofAilanthus altissimaSWINGLE. Conversion of Ailanthone into Shinjulactone C. Bulletin of the Chemical Society of Japan, 1985, 58, 2357-2360.	3.2	9
348	Constituents of Seeds ofBrucea javanica. Structures of New Bitter Principles, Yadanziolides A, B, C, Yadanziosides F, I, J, and L Bulletin of the Chemical Society of Japan, 1985, 58, 2673-2679.	3.2	35
349	Structure Determination of a New Bitter Principle, Shinjulactone L, fromAilanthus altissima. Bulletin of the Chemical Society of Japan, 1985, 58, 2723-2724.	3.2	15
350	Structures of New Quassinoid Glycosides, Yadanziosides A, B, C, D, E, G, H, and New Quassinoids, Dehydrobrusatol and Dehydrobruceantinol fromBrucea javanica(L.) MERR. Bulletin of the Chemical Society of Japan, 1985, 58, 2680-2686.	3.2	39
351	Constituents of Seeds ofAilanthus altissimaSWINGLE. Isolation and Structures of Shinjuglycosides A, B, C, and D. Bulletin of the Chemical Society of Japan, 1984, 57, 2496-2501.	3.2	25
352	SHINJULACTONE F, A NEW BITTER PRINCIPLE WITH A 5Î ² H-PICRASANE SKELETON FROMAILANTHUS ALTISSIMASWINGLE. Chemistry Letters, 1984, 13, 555-556.	1.3	9
353	Structure Determination of Bitter Principles ofAilanthus altissima.Structures of Shinjulactones F, I, J, and K. Bulletin of the Chemical Society of Japan, 1984, 57, 2885-2892.	3.2	18
354	Shinjulactones G and H, New Bitter Principles ofAilanthus altissimaSWINGLE. Bulletin of the Chemical Society of Japan, 1984, 57, 2013-2014.	3.2	19
355	Structure Determination of Bitter Principles ofAilanthus altissima. Structures of Shinjulactones B, D, and E. Bulletin of the Chemical Society of Japan, 1984, 57, 2484-2489.	3.2	25
356	Structures of yadanziolides A, B, and C, new bitter principles from Brucea javanica Chemical and Pharmaceutical Bulletin, 1984, 32, 4698-4701.	1.3	20
357	New quassinoid glycosides, yadanziosides A-H, from Brucea javanica Chemical and Pharmaceutical Bulletin, 1984, 32, 4702-4705.	1.3	21
358	Conversion of ailanthone into shinjulactone C through an ionic [4+2] cycloaddition reaction. Tetrahedron Letters, 1983, 24, 4843-4846.	1.4	10
359	Constituents of the Root Bark ofAilanthus altissimaSWINGLE. Isolation and X-Ray Crystal Structures of Shinjudilactone and Shinjulactone C and Conversion of Ailanthone into Shinjudilactone. Bulletin of the Chemical Society of Japan, 1983, 56, 3683-3693.	3.2	38
360	Structure Determination of Bitter Principles inAilanthus altissima.Structure of Shinjulactone A and Revised Structure of Ailanthone. Bulletin of the Chemical Society of Japan, 1983, 56, 3694-3698.	3.2	30

#	Article	IF	CITATIONS
361	Structures of shinjulactones D and E, new bitter principles of Ailanthus altissima Swingle Chemical and Pharmaceutical Bulletin, 1983, 31, 2179-2182.	1.3	7
362	Conversion of ailanthone into shinjudilactone, a backbone-rearranged picrasane Chemical and Pharmaceutical Bulletin, 1982, 30, 1917-1920.	1.3	8
363	ON THE STRUCTURE OF AILANTHONE, A BITTER PRINCIPLE FROMAILANTHUS ALTISSIMA. Chemistry Letters, 1982, 11, 661-662.	1.3	15
364	Shinjulactone C, a new quassinoid ith a 1α, 12α:5α, 13α-dicyclo-9β-picrasane skeleton from SWINGLE. Tetrahedron Letters, 1982, 23, 1205-1206.	1.4	18
365	SHINJUDILACTONE, A NEW BITTER PRINCIPLE FROMAILANTHUS ALTISSIMASWINGLE. Chemistry Letters, 1981, 10, 1597-1598.	1.3	18