

Qinghua Tao

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3798270/publications.pdf>

Version: 2024-02-01

40
papers

1,626
citations

361413

20
h-index

330143

37
g-index

40
all docs

40
docs citations

40
times ranked

1980
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 1 | Cryo-EM structure of the nuclear ring from <i>Xenopus laevis</i> nuclear pore complex. <i>Cell Research</i> , 2022, 32, 349-358. | 12.0 | 19 |
| 2 | Structure of the cytoplasmic ring of the <i>Xenopus laevis</i> nuclear pore complex. <i>Science</i> , 2022, 376, . | 12.6 | 44 |
| 3 | HMCES modulates the transcriptional regulation of nodal/activin and BMP signaling in mESCs. <i>Cell Reports</i> , 2022, 40, 111038. | 6.4 | 1 |
| 4 | Phase separation of Axin organizes the β -catenin destruction complex. <i>Journal of Cell Biology</i> , 2021, 220, . | 5.2 | 59 |
| 5 | The cytokine FAM3B/PANDER is an FGFR ligand that promotes posterior development in <i>Xenopus</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2021, 118, . | 7.1 | 5 |
| 6 | Lysosomal degradation of the maternal dorsal determinant Hwa safeguards dorsal body axis formation. <i>EMBO Reports</i> , 2021, 22, e53185. | 4.5 | 7 |
| 7 | Molecular architecture of the luminal ring of the <i>Xenopus laevis</i> nuclear pore complex. <i>Cell Research</i> , 2020, 30, 532-540. | 12.0 | 51 |
| 8 | Structure of the cytoplasmic ring of the <i>Xenopus laevis</i> nuclear pore complex by cryo-electron microscopy single particle analysis. <i>Cell Research</i> , 2020, 30, 520-531. | 12.0 | 51 |
| 9 | Expression of the hormonal FGF co-receptor <i>Klotho beta</i> in the <i>Xenopus laevis</i> model. <i>Cell Biology International</i> , 2019, 43, 207-213. | 3.0 | 0 |
| 10 | Compound C induces protective autophagy in human cholangiocarcinoma cells via Akt/mTOR-independent pathway. <i>Journal of Cellular Biochemistry</i> , 2018, 119, 5538-5550. | 2.6 | 26 |
| 11 | Maternal Huluwa dictates the embryonic body axis through β -catenin in vertebrates. <i>Science</i> , 2018, 362, . | 12.6 | 52 |
| 12 | c-Myc promotes cholangiocarcinoma cells to overcome contact inhibition via the mTOR pathway. <i>Oncology Reports</i> , 2017, 38, 2498-2506. | 2.6 | 11 |
| 13 | KDM3A-mediated demethylation of histone H3 lysine 9 facilitates the chromatin binding of Neurog2 during neurogenesis. <i>Development (Cambridge)</i> , 2017, 144, 3674-3685. | 2.5 | 26 |
| 14 | Conserved gene regulatory module specifies lateral neural borders across bilaterians. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E6352-E6360. | 7.1 | 24 |
| 15 | The RNF146 E3 ubiquitin ligase is required for the control of Wnt signaling and body pattern formation in <i>Xenopus</i> . <i>Mechanisms of Development</i> , 2017, 147, 28-36. | 1.7 | 5 |
| 16 | The MLL/Setd1b methyltransferase is required for the Spemann's organizer gene activation in <i>Xenopus</i> . <i>Mechanisms of Development</i> , 2016, 142, 1-9. | 1.7 | 8 |
| 17 | Ascl1 represses the mesendoderm induction in <i>Xenopus</i> . <i>Acta Biochimica Et Biophysica Sinica</i> , 2016, 48, 1006-1015. | 2.0 | 0 |
| 18 | NF2/Merlin is required for the axial pattern formation in the <i>Xenopus laevis</i> embryo. <i>Mechanisms of Development</i> , 2015, 138, 305-312. | 1.7 | 7 |

| # | ARTICLE | IF | CITATIONS |
|----|---|------|-----------|
| 19 | Sebox regulates mesoderm formation in early amphibian embryos. <i>Developmental Dynamics</i> , 2015, 244, 1415-1426. | 1.8 | 6 |
| 20 | A novel role for ASCL1 in the regulation of mesendoderm formation via HDAC-dependent antagonism of VegT function. <i>Development (Cambridge)</i> , 2015, 143, 492-503. | 2.5 | 17 |
| 21 | Small C-terminal Domain Phosphatase 3 Dephosphorylates the Linker Sites of Receptor-regulated Smads (R-Smads) to Ensure Transforming Growth Factor β^2 (TGF β^2)-mediated Germ Layer Induction in <i>Xenopus</i> Embryos. <i>Journal of Biological Chemistry</i> , 2015, 290, 17239-17249. | 3.4 | 6 |
| 22 | Organizer-derived Bmp2 is required for the formation of a correct Bmp activity gradient during embryonic development. <i>Nature Communications</i> , 2014, 5, 3766. | 12.8 | 30 |
| 23 | NEDD4L regulates convergent extension movements in <i>Xenopus</i> embryos via Disheveled-mediated non-canonical Wnt signaling. <i>Developmental Biology</i> , 2014, 392, 15-25. | 2.0 | 29 |
| 24 | Maternal Mga is required for Wnt signaling and organizer formation in the early <i>Xenopus</i> embryo. <i>Acta Biochimica Et Biophysica Sinica</i> , 2012, 44, 939-947. | 2.0 | 3 |
| 25 | Regulation of Classical Cadherin Membrane Expression and F-Actin Assembly by Alpha-Catenins, during <i>Xenopus</i> Embryogenesis. <i>PLoS ONE</i> , 2012, 7, e38756. | 2.5 | 6 |
| 26 | Transmembrane Protein 198 Promotes LRP6 Phosphorylation and Wnt Signaling Activation. <i>Molecular and Cellular Biology</i> , 2011, 31, 2577-2590. | 2.3 | 37 |
| 27 | N- and E-cadherins in <i>Xenopus</i> are specifically required in the neural and non-neural ectoderm, respectively, for F-actin assembly and morphogenetic movements. <i>Development (Cambridge)</i> , 2009, 136, 1327-1338. | 2.5 | 110 |
| 28 | Wnt5a and Wnt11 interact in a maternal Dkk1-regulated fashion to activate both canonical and non-canonical signaling in <i>Xenopus</i> axis formation. <i>Development (Cambridge)</i> , 2008, 135, 3719-3729. | 2.5 | 131 |
| 29 | Wnt11/ β^2 -catenin signaling in both oocytes and early embryos acts through LRP6-mediated regulation of axin. <i>Development (Cambridge)</i> , 2007, 134, 503-513. | 2.5 | 86 |
| 30 | G-protein-coupled signals control cortical actin assembly by controlling cadherin expression in the early <i>Xenopus</i> embryo. <i>Development (Cambridge)</i> , 2007, 134, 2651-2661. | 2.5 | 32 |
| 31 | Jun NH2-terminal kinase (JNK) prevents nuclear beta-catenin accumulation and regulates axis formation in <i>Xenopus</i> embryos. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 16313-16318. | 7.1 | 60 |
| 32 | Lysophosphatidic acid signaling controls cortical actin assembly and cytoarchitecture in <i>Xenopus</i> embryos. <i>Development (Cambridge)</i> , 2005, 132, 805-816. | 2.5 | 26 |
| 33 | A novel G protein-coupled receptor, related to GPR4, is required for assembly of the cortical actin skeleton in early <i>Xenopus</i> embryos. <i>Development (Cambridge)</i> , 2005, 132, 2825-2836. | 2.5 | 19 |
| 34 | Maternal Wnt11 Activates the Canonical Wnt Signaling Pathway Required for Axis Formation in <i>Xenopus</i> Embryos. <i>Cell</i> , 2005, 120, 857-871. | 28.9 | 454 |
| 35 | The role of maternal CREB in early embryogenesis of <i>Xenopus laevis</i> . <i>Developmental Biology</i> , 2003, 261, 337-352. | 2.0 | 17 |
| 36 | The <i>Xenopus</i> noggin promoter drives roof-plate specific transcription. <i>NeuroReport</i> , 2003, 14, 2163-2166. | 1.2 | 6 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 37 | Repression through a distal TCF-3 binding site restricts <i>Xenopus myf-5</i> expression in gastrula mesoderm. <i>Mechanisms of Development</i> , 2002, 115, 79-89. | 1.7 | 27 |
| 38 | The roles of three signaling pathways in the formation and function of the Spemann Organizer. <i>Development (Cambridge)</i> , 2002, 129, 4027-4043. | 2.5 | 116 |
| 39 | An interferon regulatory factor-like binding element restricts <i>Xmyf-5</i> expression in the posterior somites during <i>Xenopus</i> myogenesis. <i>FEBS Letters</i> , 2001, 505, 47-52. | 2.8 | 12 |
| 40 | Analysis for the dorsalization potency of the animal blastomeres of the 16 cell stage <i>Xenopus</i> embryo. <i>Science Bulletin</i> , 1999, 44, 538-540. | 1.7 | 0 |