## Carmen Ghilardi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3849821/publications.pdf

Version: 2024-02-01

623734	752698
20 14	20
ions h-ind	lex g-index
22 22	1291
itations times r	anked citing authors
	20 14 ions h-ind

#	Article	IF	CITATIONS
1	The DNA-PK Inhibitor AZD7648 Sensitizes Patient-Derived Ovarian Cancer Xenografts to Pegylated Liposomal Doxorubicin and Olaparib Preventing Abdominal Metastases. Molecular Cancer Therapeutics, 2022, 21, 555-567.	4.1	11
2	PGC1 $\hat{l}\pm\hat{l}^2$ Expression Predicts Therapeutic Response to Oxidative Phosphorylation Inhibition in Ovarian Cancer. Cancer Research, 2022, 82, 1423-1434.	0.9	14
3	VEGF pathway inhibition potentiates PARP inhibitor efficacy in ovarian cancer independent of BRCA status. Journal of Hematology and Oncology, 2021, 14, 186.	17.0	27
4	A novel L1CAM isoform with angiogenic activity generated by NOVA2-mediated alternative splicing. ELife, 2019, $8$ , .	6.0	38
5	Contribution of tumor endothelial cells to drug resistance: anti-angiogenic tyrosine kinase inhibitors act as p-glycoprotein antagonists. Angiogenesis, 2017, 20, 233-241.	7.2	22
6	Thrombospondinâ $\in$ 1 is part of a Slugâ $\in$ independent motility and metastatic program in cutaneous melanoma, in association with $\langle scp \rangle VEGFR \langle scp \rangle $ a $\in$ 1 and $\langle scp \rangle FGF \langle scp \rangle $ a $\in$ 2. Pigment Cell and Melanoma Research, 2015, 28, 73-81.	3.3	45
7	Sunitinib prevents cachexia and prolongs survival of mice bearing renal cancer by restraining STAT3 and MuRF-1 activation in muscle. Oncotarget, 2015, 6, 3043-3054.	1.8	38
8	Trypsinogen 4 boosts tumor endothelial cells migration through proteolysis of tissue factor pathway inhibitor-2. Oncotarget, 2015, 6, 28389-28400.	1.8	13
9	Patient-Derived Ovarian Tumor Xenografts Recapitulate Human Clinicopathology and Genetic Alterations. Cancer Research, 2014, 74, 6980-6990.	0.9	110
10	Inhibition of SIRT2 Potentiates the Anti-motility Activity of Taxanes: Implications for Antineoplastic Combination Therapies. Neoplasia, 2012, 14, 846-IN16.	5.3	28
11	Regulator of G-protein signaling 5 (RGS5) protein: a novel marker of cancer vasculature elicited and sustained by the tumor's proangiogenic microenvironment. Cellular and Molecular Life Sciences, 2012, 69, 1167-1178.	5.4	40
12	Protease-activated receptor-1 (PAR-1) promotes the motility of human melanomas and is associated to their metastatic phenotype. Clinical and Experimental Metastasis, 2010, 27, 43-53.	3.3	18
13	Dual Targeting of Tumor and Endothelial Cells by Gonadotropin-Releasing Hormone Agonists to Reduce Melanoma Angiogenesis. Endocrinology, 2010, 151, 4643-4653.	2.8	15
14	Identification of novel vascular markers through gene expression profiling of tumor-derived endothelium. BMC Genomics, 2008, 9, 201.	2.8	56
15	Gene expression correlating with response to paclitaxel in ovarian carcinoma xenografts. Molecular Cancer Therapeutics, 2004, 3, 111-21.	4.1	46
16	Anticancer Therapy with Angiogenesis Inhibitors. Tumori, 2001, 87, 14-16.	1.1	1
17	p73 overexpression increases VEGF and reduces thrombospondin-1 production: implications for tumor angiogenesis. Oncogene, 2001, 20, 7293-7300.	5.9	51
18	Posttranscriptional Stimulation of Endothelial Cell Matrix Metalloproteinases 2 and 1 by Endothelioma Cells. Experimental Cell Research, 2000, 258, 384-394.	2.6	43