

Yonghe Li

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3896325/publications.pdf>

Version: 2024-02-01

20
papers

1,456
citations

471509

17
h-index

677142

22
g-index

23
all docs

23
docs citations

23
times ranked

2522
citing authors

#	ARTICLE	IF	CITATIONS
1	Multi-targeted therapy of cancer by niclosamide: A new application for an old drug. <i>Cancer Letters</i> , 2014, 349, 8-14.	7.2	303
2	Niclosamide Suppresses Cancer Cell Growth By Inducing Wnt Co-Receptor LRP6 Degradation and Inhibiting the Wnt/ β -Catenin Pathway. <i>PLoS ONE</i> , 2011, 6, e29290.	2.5	187
3	Low-Density Lipoprotein Receptor-Related Protein 1 Promotes Cancer Cell Migration and Invasion by Inducing the Expression of Matrix Metalloproteinases 2 and 9. <i>Cancer Research</i> , 2009, 69, 879-886.	0.9	160
4	Low-Density Lipoprotein Receptor Family. <i>Molecular Neurobiology</i> , 2001, 23, 53-68.	4.0	124
5	Silibinin inhibits Wnt/ β -catenin signaling by suppressing Wnt co-receptor LRP6 expression in human prostate and breast cancer cells. <i>Cellular Signalling</i> , 2012, 24, 2291-2296.	3.6	105
6	Effect of Niclosamide on Basal-like Breast Cancers. <i>Molecular Cancer Therapeutics</i> , 2014, 13, 800-811.	4.1	78
7	LRP6 expression promotes cancer cell proliferation and tumorigenesis by altering β -catenin subcellular distribution. <i>Oncogene</i> , 2004, 23, 9129-9135.	5.9	75
8	Niclosamide and its analogs are potent inhibitors of Wnt/ β -catenin, mTOR and STAT3 signaling in ovarian cancer. <i>Oncotarget</i> , 2016, 7, 86803-86815.	1.8	64
9	Role of Wnt Co-Receptor LRP6 in Triple Negative Breast Cancer Cell Migration and Invasion. <i>Journal of Cellular Biochemistry</i> , 2017, 118, 2968-2976.	2.6	51
10	Mesd binds to mature LDL-receptor-related protein-6 and antagonizes ligand binding. <i>Journal of Cell Science</i> , 2005, 118, 5305-5314.	2.0	49
11	Mesd Is a Universal Inhibitor of Wnt Coreceptors LRP5 and LRP6 and Blocks Wnt/ β -Catenin Signaling in Cancer Cells. <i>Biochemistry</i> , 2010, 49, 4635-4643.	2.5	43
12	In vitro Invasiveness of Human Breast Cancer Cells Is Promoted by Low Density Lipoprotein Receptor-Related Protein. <i>Invasion & Metastasis</i> , 1998, 18, 240-251.	0.5	32
13	Receptor-Associated Protein Facilitates Proper Folding and Maturation of the Low-Density Lipoprotein Receptor and Its Class 2 Mutants. <i>Biochemistry</i> , 2002, 41, 4921-4928.	2.5	32
14	Discovery of a novel inhibitor of kinesin-like protein KIFC1. <i>Biochemical Journal</i> , 2016, 473, 1027-1035.	3.7	32
15	Degradation of the LDL receptor class 2 mutants is mediated by a proteasome-dependent pathway. <i>Journal of Lipid Research</i> , 2004, 45, 1084-1091.	4.2	30
16	Mesd is a general inhibitor of different Wnt ligands in Wnt/LRP signaling and inhibits PC-3 tumor growth in vivo. <i>FEBS Letters</i> , 2011, 585, 3120-3125.	2.8	26
17	Discovery of novel frizzled-7 inhibitors by targeting the receptor's transmembrane domain. <i>Oncotarget</i> , 2017, 8, 91459-91470.	1.8	26
18	Striking differences of LDL receptor-related protein 1B expression in mouse and human. <i>Biochemical and Biophysical Research Communications</i> , 2005, 333, 868-873.	2.1	14

#	ARTICLE	IF	CITATIONS
19	The C-Terminal Region Mesd Peptide Mimics Full-Length Mesd and Acts as an Inhibitor of Wnt/ β 2-Catenin Signaling in Cancer Cells. PLoS ONE, 2013, 8, e58102.	2.5	12
20	LRP5, a multifunctional cell surface receptor. Current Opinion in Lipidology, 2004, 15, 361-363.	2.7	3