## Mogens GrÃ, nvold

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3905661/publications.pdf

Version: 2024-02-01

168 papers 6,818 citations

45 h-index 74163 75 g-index

179 all docs

179 docs citations

179 times ranked 7977 citing authors

#	Article	IF	CITATIONS
1	The development of the EORTC QLQ-C15-PAL: A shortened questionnaire for cancer patients in palliative care. European Journal of Cancer, 2006, 42, 55-64.	2.8	502
2	Replication and validation of higher order models demonstrated that a summary score for the EORTC QLQ-C30 is robust. Journal of Clinical Epidemiology, 2016, 69, 79-88.	5 <b>.</b> 0	306
3	Psychological distress and fatigue predicted recurrence and survival in primary breast cancer patients. Breast Cancer Research and Treatment, 2007, 105, 209-219.	2.5	268
4	General population normative data for the EORTC QLQ-C30 health-related quality of life questionnaire based on 15,386 persons across 13 European countries, Canada and the Unites States. European Journal of Cancer, 2019, 107, 153-163.	2.8	259
5	Chronic pain and other sequelae in longâ€ŧerm breast cancer survivors: Nationwide survey in Denmark. European Journal of Pain, 2009, 13, 478-485.	2.8	204
6	Quality of life in patients with benign thyroid disorders. A review. European Journal of Endocrinology, 2006, 154, 501-510.	3.7	184
7	Thresholds for clinical importance were established to improve interpretation of the EORTC QLQ-C30 in clinical practice and research. Journal of Clinical Epidemiology, 2020, 118, 1-8.	5.0	184
8	International standards for the analysis of quality-of-life and patient-reported outcome endpoints in cancer randomised controlled trials: recommendations of the SISAQOL Consortium. Lancet Oncology, The, 2020, 21, e83-e96.	10.7	180
9	Validity and reliability of the novel thyroid-specific quality of life questionnaire, ThyPRO. European Journal of Endocrinology, 2010, 162, 161-167.	3.7	163
10	Analysing data from patient-reported outcome and quality of life endpoints for cancer clinical trials: a start in setting international standards. Lancet Oncology, The, 2016, 17, e510-e514.	10.7	158
11	The Thyroid-Related Quality of Life Measure ThyPRO Has Good Responsiveness and Ability to Detect Relevant Treatment Effects. Journal of Clinical Endocrinology and Metabolism, 2014, 99, 3708-3717.	3.6	123
12	Self-assessment in cancer patients referred to palliative care. Cancer, 2002, 94, 512-520.	4.1	119
13	Symptomatology of cancer patients in palliative care. European Journal of Cancer, 2002, 38, 788-794.	2.8	113
14	Symptoms and problems in a nationally representative sample of advanced cancer patients. Palliative Medicine, 2009, 23, 491-501.	3.1	112
15	Disease-Specific as Well as Generic Quality of Life Is Widely Impacted in Autoimmune Hypothyroidism and Improves during the First Six Months of Levothyroxine Therapy. PLoS ONE, 2016, 11, e0156925.	2.5	109
16	Differential item functioning (DIF) analyses of health-related quality of life instruments using logistic regression. Health and Quality of Life Outcomes, 2010, 8, 81.	2.4	92
17	Cancer caregiving tasks and consequences and their associations with caregiver status and the caregiver's relationship to the patient: a survey. BMC Cancer, 2014, 14, 541.	2.6	86
18	Development of a Short Version of the Thyroid-Related Patient-Reported Outcome ThyPRO. Thyroid, 2015, 25, 1069-1079.	4.5	82

#	Article	IF	CITATIONS
19	Construct validity of a revised Physical Activity Scale and testing by cognitive interviewing. Scandinavian Journal of Public Health, 2010, 38, 707-714.	2.3	80
20	The Female Sexual Function Index (FSFI)â€"A Systematic Review of Measurement Properties. Journal of Sexual Medicine, 2019, 16, 640-660.	0.6	80
21	Prognostic value of patient-reported outcomes from international randomised clinical trials on cancer: a systematic review. Lancet Oncology, The, 2019, 20, e685-e698.	10.7	80
22	Which Domains of Thyroid-Related Quality of Life Are Most Relevant? Patients and Clinicians Provide Complementary Perspectives. Thyroid, 2007, 17, 647-654.	4.5	78
23	The use of EORTC measures in daily clinical practiceâ€"A synopsis of a newly developed manual. European Journal of Cancer, 2016, 68, 73-81.	2.8	75
24	A prospective evaluation of Dignity Therapy in advanced cancer patients admitted to palliative care. Palliative Medicine, 2014, 28, 448-458.	3.1	74
25	A longitudinal study of palliative care. Cancer, 2005, 103, 1747-1755.	4.1	73
26	Health-Related Quality of Life in EORTC clinical trials — 30 years of progress from methodological developments to making a real impact on oncology practice. European Journal of Cancer, Supplement, 2012, 10, 141-149.	2.2	73
27	The EORTC CAT Coreâ€"The computer adaptive version of the EORTC QLQ-C30 questionnaire. European Journal of Cancer, 2018, 100, 8-16.	2.8	68
28	Advance care planning in patients with advanced cancer: A 6-country, cluster-randomised clinical trial. PLoS Medicine, 2020, 17, e1003422.	8.4	68
29	Danish population-based reference data for the EORTC QLQ-C30: associations with gender, age and morbidity. Quality of Life Research, 2014, 23, 2183-2193.	3.1	67
30	Randomised clinical trial of early specialist palliative care plus standard care versus standard care alone in patients with advanced cancer: The Danish Palliative Care Trial. Palliative Medicine, 2017, 31, 814-824.	3.1	67
31	Statistical analysis of patient-reported outcome data in randomised controlled trials of locally advanced and metastatic breast cancer: a systematic review. Lancet Oncology, The, 2018, 19, e459-e469.	10.7	66
32	Improving a newly developed patient-reported outcome for thyroid patients, using cognitive interviewing. Quality of Life Research, 2008, 17, 1009-1017.	3.1	65
33	The relationship between overall quality of life and its subdimensions was influenced by culture: analysis of an international database. Journal of Clinical Epidemiology, 2008, 61, 788-795.	5.0	65
34	What Facilitates "Patient Empowerment―in Cancer Patients During Follow-Up: A Qualitative Systematic Review of the Literature. Qualitative Health Research, 2018, 28, 292-304.	2.1	63
35	The interaction between informal cancer caregivers and health care professionals: a survey of caregivers' experiences of problems and unmet needs. Supportive Care in Cancer, 2015, 23, 1719-1733.	2.2	62
36	The chronic autoimmune thyroiditis quality of life selenium trial (CATALYST): study protocol for a randomized controlled trial. Trials, 2014, 15, 115.	1.6	60

#	Article	lF	CITATIONS
37	Is Thyroid Autoimmunity per se a Determinant of Quality of Life in Patients with Autoimmune Hypothyroidism?. European Thyroid Journal, 2012, 1, 186-192.	2.4	59
38	Health care utilisation and characteristics of long-term breast cancer survivors: Nationwide survey in Denmark. European Journal of Cancer, 2009, 45, 625-633.	2.8	56
39	Expert conference on cancer pain assessment and classification-the need for international consensus: working proposals on international standards. BMJ Supportive and Palliative Care, 2011, 1, 281-287.	1.6	56
40	Use of item response theory to develop a shortened version of the EORTC QLQ-C30 emotional functioning scale. Quality of Life Research, 2004, 13, 1683-1697.	3.1	55
41	Quality-of-Life Impairments Persist Six Months After Treatment of Graves' Hyperthyroidism and Toxic Nodular Goiter: A Prospective Cohort Study. Thyroid, 2016, 26, 1010-1018.	4.5	55
42	Quality of Life in Patients with Benign Nontoxic Goiter: Impact of Disease and Treatment Response, and Comparison with the General Population. Thyroid, 2015, 25, 284-291.	4.5	54
43	Comparing higher order models for the EORTC QLQ-C30. Quality of Life Research, 2012, 21, 1607-1617.	3.1	52
44	Cross-cultural validity of the thyroid-specific quality-of-life patient-reported outcome measure, ThyPRO. Quality of Life Research, 2015, 24, 769-780.	3.1	52
45	Do advanced cancer patients in Denmark receive the help they need? A nationally representative survey of the need related to 12 frequent symptoms/problems. Psycho-Oncology, 2013, 22, 1724-1730.	2.3	49
46	Establishing anchor-based minimally important differences (MID) with the EORTC quality-of-life measures: a meta-analysis protocol. BMJ Open, 2018, 8, e019117.	1.9	45
47	Patient empowerment: a systematic review of questionnaires measuring empowerment in cancer patients. Acta Oncol $\tilde{A}^3$ gica, 2017, 56, 156-165.	1.8	44
48	Advance care planning $\hat{a} \in \hat{a}$ a multi-centre cluster randomised clinical trial: the research protocol of the ACTION study. BMC Cancer, 2016, 16, 264.	2.6	43
49	Moving forward toward standardizing analysis of quality of life data in randomized cancer clinical trials. Clinical Trials, 2018, 15, 624-630.	1.6	42
50	Confirmatory factor analysis of the thyroid-related quality of life questionnaire ThyPRO. Health and Quality of Life Outcomes, 2014, 12, 126.	2.4	41
51	A systematic review of the measurement properties of the Body Image Scale (BIS) in cancer patients. Supportive Care in Cancer, 2018, 26, 1715-1726.	2.2	41
52	Item response theory was used to shorten EORTC QLQ-C30 scales for use in palliative care. Journal of Clinical Epidemiology, 2006, 59, 36-44.	5.0	40
53	Testing the feasibility of the Dignity Therapy interview: adaptation for the Danish culture. BMC Palliative Care, 2010, 9, 21.	1.8	40
54	Anxiety and Depression Are More Prevalent in Patients with Graves' Disease than in Patients with Nodular Goitre. European Thyroid Journal, 2014, 3, 173-178.	2.4	40

#	Article	lF	CITATIONS
55	Development of computerized adaptive testing (CAT) for the EORTC QLQ-C30 physical functioning dimension. Quality of Life Research, 2011, 20, 479-490.	3.1	38
56	Interpreting European Organisation for Research and Treatment for Cancer Quality of life Questionnaire core 30 scores as minimally importantly different for patients with malignant melanoma. European Journal of Cancer, 2018, 104, 169-181.	2.8	38
57	Establishing the European Norm for the health-related quality of life domains of the computer-adaptive test EORTC CAT Core. European Journal of Cancer, 2019, 107, 133-141.	2.8	38
58	Effects of melatonin on physical fatigue and other symptoms in patients with advanced cancer receiving palliative care: A doubleâ€blind placeboâ€controlled crossover trial. Cancer, 2015, 121, 3727-3736.	4.1	37
59	Health-related quality of life in melanoma patients: Impact of melanoma-related limb lymphoedema. European Journal of Cancer, 2017, 85, 122-132.	2.8	37
60	The EORTC computer-adaptive tests measuring physical functioning and fatigue exhibited high levels of measurement precision and efficiency. Journal of Clinical Epidemiology, 2013, 66, 330-339.	5.0	33
61	Updated EORTC QLQ-C30 general population norm data for Germany. European Journal of Cancer, 2020, 137, 161-170.	2.8	32
62	Is admittance to specialised palliative care among cancer patients related to sex, age and cancer diagnosis? A nation-wide study from the Danish Palliative Care Database (DPD). BMC Palliative Care, 2017, 16, 21.	1.8	31
63	Factors associated with Danish cancer patients' return to work. A report from the population-based study †The Cancer Patient's World'. Cancer Epidemiology, 2012, 36, 222-229.	1.9	29
64	Few items in the thyroid-related quality of life instrument ThyPRO exhibited differential item functioning. Quality of Life Research, 2014, 23, 327-338.	3.1	29
65	Development of an item bank for the EORTC Role Functioning Computer Adaptive Test (EORTC RF-CAT). Health and Quality of Life Outcomes, 2016, 14, 72.	2.4	29
66	Development of an item bank for computerized adaptive test (CAT) measurement of pain. Quality of Life Research, 2016, 25, 1-11.	3.1	29
67	Minimally important differences for interpreting European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire Core 30 scores in patients with head and neck cancer. Head and Neck, 2020, 42, 3141-3152.	2.0	28
68	Breast cancer patients on adjuvant chemotherapy report a wide range of problems not identified by health-care staff. Breast Cancer Research and Treatment, 2007, 103, 185-195.	2.5	27
69	The EORTC emotional functioning computerized adaptive test: phases l–III of a crossâ€cultural item bank development. Psycho-Oncology, 2014, 23, 397-403.	2.3	27
70	Development and implementation of PROgmatic: A clinical trial management system for pragmatic multi-centre trials, optimised for electronic data capture and patient-reported outcomes. Clinical Trials, 2014, 11, 344-354.	1.6	26
71	Conceptualizing patient empowerment in cancer follow-up by combining theory and qualitative data. Acta Oncol $ ilde{A}^3$ gica, 2017, 56, 232-238.	1.8	26
72	Cross-sectorial cooperation and supportive care in general practice: cancer patients' experiences. Family Practice, 2011, 28, 532-540.	1.9	24

#	Article	IF	CITATIONS
73	The initial development of the $\hat{a}\in$ Cancer Caregiving Tasks, Consequences and Needs Questionnaire $\hat{a}\in$ (CaTCoN). Acta Oncol $\hat{A}^3$ gica, 2012, 51, 1009-1019.	1.8	23
74	Chemotherapy versus ovarian ablation as adjuvant therapy for breast cancer: impact on health-related quality of life in a randomized trial. Breast Cancer Research and Treatment, 2006, 98, 275-284.	2.5	22
75	Psychometric evaluation of the EORTC computerized adaptive test (CAT) fatigue item pool. Quality of Life Research, 2013, 22, 2443-2454.	3.1	22
76	Is specialized palliative cancer care associated with use of antineoplastic treatment at the end of life? A population-based cohort study. Palliative Medicine, 2018, 32, 1509-1517.	3.1	22
77	Minimally important differences for interpreting the EORTC QLQ 30 in patients with advanced colorectal cancer treated with chemotherapy. Colorectal Disease, 2020, 22, 2278-2287.	1.4	22
78	The validity and reliability of the â€~Cancer Caregiving Tasks, Consequences and Needs Questionnaire' (CaTCoN). Acta Oncológica, 2014, 53, 966-974.	1.8	21
79	Danish Palliative Care Database. Clinical Epidemiology, 2016, Volume 8, 637-643.	3.0	21
80	Prevalence and possible predictors of sexual dysfunction and self-reported needs related to the sexual life of advanced cancer patients. Acta Oncol $\tilde{A}^3$ gica, 2019, 58, 769-775.	1.8	21
81	International validation of the EORTC CAT Core: a new adaptive instrument for measuring core quality of life domains in cancer. Quality of Life Research, 2020, 29, 1405-1417.	3.1	21
82	An emotional functioning item bank of 24 items for computerized adaptive testing (CAT) was established. Journal of Clinical Epidemiology, 2016, 70, 90-100.	5.0	20
83	A new model of early, integrated palliative care: palliative rehabilitation for newly diagnosed patients with non-resectable cancer. Supportive Care in Cancer, 2019, 27, 3291-3300.	2.2	20
84	Methodological issues in the assessment of health-related quality of life in palliative care trials. Acta Anaesthesiologica Scandinavica, 1999, 43, 948-953.	1.6	19
85	Early, integrated palliative rehabilitation improves quality of life of patients with newly diagnosed advanced cancer: The Pal-Rehab randomized controlled trial. Palliative Medicine, 2021, 35, 1344-1355.	3.1	19
86	Detailed statistical analysis plan for the Danish Palliative Care Trial (DanPaCT). Trials, 2014, 15, 376.	1.6	18
87	Focusing on Core Patient-Reported Outcomes in Cancer Clinical Trials—Letter. Clinical Cancer Research, 2016, 22, 5617-5617.	7.0	18
88	Admittance to specialized palliative care (SPC) of patients with an assessed need: a study from the Danish palliative care database (DPD). Acta Oncolųgica, 2017, 56, 1210-1217.	1.8	18
89	Validation of the European Organisation for Research and Treatment of Cancer Quality of Life Questionnaire Core 30 Summary Score in Patients With Hematologic Malignancies. Value in Health, 2019, 22, 1303-1310.	0.3	18
90	A randomised, multicentre clinical trial of specialised palliative care plus standard treatment versus standard treatment alone for cancer patients with palliative care needs: the Danish palliative care trial (DanPaCT) protocol. BMC Palliative Care, 2013, 12, 37.	1.8	17

#	Article	IF	Citations
91	A parallel-group randomized clinical trial of individually tailored, multidisciplinary, palliative rehabilitation for patients with newly diagnosed advanced cancer: the Pal-Rehab study protocol. BMC Cancer, 2017, 17, 560.	2.6	17
92	Determining minimal important change for the thyroid-related quality of life questionnaire ThyPRO. Endocrine Connections, 2021, 10, 316-324.	1.9	17
93	Validation of the Danish version of the disease specific instrument EORTC QLQ-CR38 to assess Health-related quality of life in patients with colorectal cancer. Health and Quality of Life Outcomes, 2012, 10, 150.	2.4	16
94	A crossâ€cultural convergent parallel mixed methods study of what makes a cancerâ€related symptom or functional health problem clinically important. Psycho-Oncology, 2018, 27, 548-555.	2.3	16
95	Age, cancer site and gender associations with symptoms and problems in specialised palliative care: a large, nationwide, register-based study. BMJ Supportive and Palliative Care, 2019, , bmJspcare-2019-001880.	1.6	16
96	Quality of life in patients with diffuse large B-cell lymphoma treated with dose-dense chemotherapy is only affected temporarily. Leukemia and Lymphoma, 2011, 52, 400-408.	1.3	15
97	Cancer patients' evaluation of communication: a report from the population-based study †The Cancer Patient's World'. Supportive Care in Cancer, 2013, 21, 235-244.	2.2	15
98	Health-related quality of life in early breast cancer. Danish Medical Bulletin, 2010, 57, B4184.	0.3	15
99	The relatives' perspective on advanced cancer care in Denmark. A cross-sectional survey. Supportive Care in Cancer, 2012, 20, 3179-3188.	2.2	14
100	Development of a lack of appetite item bank for computer-adaptive testing (CAT). Supportive Care in Cancer, 2015, 23, 1541-1548.	2.2	14
101	How Should Thyroid-Related Quality of Life Be Assessed? Recalled Patient-Reported Outcomes Compared to Here-and-Now Measures. Thyroid, 2018, 28, 1561-1570.	4.5	13
102	Measuring health-related quality of life in colorectal cancer patients: systematic review of measurement properties of the EORTC QLQ-CR29. Supportive Care in Cancer, 2019, 27, 2395-2412.	2.2	13
103	Exploratory analyses of the Danish Palliative Care Trial (DanPaCT): a randomized trial of early specialized palliative care plus standard care versus standard care in advanced cancer patients. Supportive Care in Cancer, 2020, 28, 2145-2155.	2.2	13
104	Thresholds for clinical importance were defined for the European Organisation for Research and Treatment of Cancer Computer Adaptive Testing Coreâ€"an adaptive measure of core quality of life domains in oncology clinical practice and research. Journal of Clinical Epidemiology, 2020, 117, 117-125.	5.0	12
105	Minimally important differences for interpreting European Organisation for Research and Treatment of Cancer (EORTC) Quality of life Questionnaire core 30 scores in patients with ovarian cancer. Gynecologic Oncology, 2020, 159, 515-521.	1.4	12
106	Minimally important differences for the EORTC QLQ-C30 in prostate cancer clinical trials. BMC Cancer, 2021, 21, 1083.	2.6	12
107	Quality of life and symptoms in patients with malignant diseases admitted to a comprehensive cancer centre. Supportive Care in Cancer, 2014, 22, 1843-1849.	2.2	11
108	Comparing three different approaches to the measurement of needs concerning fatigue in patients with advanced cancer. Quality of Life Research, 2015, 24, 2231-2238.	3.1	11

#	Article	IF	CITATIONS
109	Psychometric evaluation of an item bank for computerized adaptive testing of the EORTC QLQ-C30 cognitive functioning dimension in cancer patients. Quality of Life Research, 2017, 26, 2919-2929.	3.1	11
110	A systematic review of the measurement properties of the European Organisation for Research and Treatment of Cancer In-patient Satisfaction with Care Questionnaire, the EORTC IN-PATSAT32. Supportive Care in Cancer, 2018, 26, 2551-2560.	2.2	11
111	Does the agreement of patient and physician assessments of health related quality of life in palliative care depend on patient characteristics?. Palliative Medicine, 2007, 21, 289-294.	3.1	10
112	Measuring the quality of end-of-life care: Development, testing, and cultural validation of the Danish version of Views of Informal Carers' Evaluation of Services–Short Form. Palliative Medicine, 2018, 32, 804-814.	3.1	10
113	Patients with Benign Thyroid Diseases Experience an Impaired Sex Life. Thyroid, 2018, 28, 1261-1269.	4.5	10
114	Patient-reported outcomes (PROs) in lung cancer: Experiences from a nationwide feasibility study. Lung Cancer, 2019, 128, 67-73.	2.0	10
115	Which symptoms and problems do advanced cancer patients admitted to specialized palliative care report in addition to those included in the EORTC QLQ-C15-PAL? A register-based national study. Supportive Care in Cancer, 2020, 28, 1725-1735.	2.2	10
116	Methylphenidate as Needed for Fatigue in Patients With Advanced Cancer. A Prospective, Double-Blind, and Placebo-Controlled Study. Journal of Pain and Symptom Management, 2020, 60, 992-1002.	1.2	9
117	Clinical research in cancer palliative care: a metaresearch analysis. BMJ Supportive and Palliative Care, 2020, 10, 249-258.	1.6	9
118	Evaluating the Thresholds for Clinical Importance of the EORTC QLQ-C15-PAL in Patients Receiving Palliative Treatment. Journal of Palliative Medicine, 2021, 24, 397-404.	1.1	9
119	Using method triangulation to validate a new instrument (CPWQ-com) assessing cancer patients' satisfaction with communication. Cancer Epidemiology, 2012, 36, 29-35.	1.9	8
120	Cohabitation Status Influenced Admittance to Specialized Palliative Care for Cancer Patients: A Nationwide Study from the Danish Palliative Care Database. Journal of Palliative Medicine, 2019, 22, 164-172.	1.1	8
121	Effect of the Carer Support Needs Assessment Tool intervention (CSNAT-I) in the Danish specialised palliative care setting: a stepped-wedge cluster randomised controlled trial. BMJ Supportive and Palliative Care, 2020, , bmjspcare-2020-002467.	1.6	8
122	Testing for differences in multiple quality of life dimensions: generating hypotheses from the experience of hospital staff. Quality of Life Research, 1998, 7, 479-486.	3.1	7
123	Satisfaction with information provided to Danish cancer patients: Validation and survey results. Patient Education and Counseling, 2013, 93, 239-247.	2.2	7
124	Conversion of standard retrospective patient-reported outcomes to momentary versions: cognitive interviewing reveals varying degrees of momentary compatibility. Quality of Life Research, 2018, 27, 1065-1076.	3.1	7
125	Adaptation and cross-cultural validation of the Spanish version of the Thyroid-Related Quality-of-Life Patient-Reported Outcome questionnaire. Endocrinologia, Diabetes Y NutriciÓn, 2018, 65, 500-507.	0.3	7
126	Nausea in advanced cancer: relationships between intensity, burden, and the need for help. Supportive Care in Cancer, 2019, 27, 265-273.	2.2	7

#	Article	IF	CITATIONS
127	Size and composition of family networks of decedents: A nationwide register-based study. Palliative Medicine, 2021, 35, 1652-1662.	3.1	7
128	Effectiveness of a nurse-delivered (FOCUS+) and a web-based (iFOCUS) psychoeducational intervention for people with advanced cancer and their family caregivers (DIAdIC): study protocol for an international randomized controlled trial. BMC Palliative Care, 2021, 20, 193.	1.8	7
129	Patient reported outcome data as performance indicators in surgically treated lung cancer patients. Lung Cancer, 2019, 130, 143-148.	2.0	6
130	Patient-reported symptoms and problems at admission to specialized palliative care improved survival prediction in 30,969 cancer patients: A nationwide register-based study. Palliative Medicine, 2020, 34, 795-805.	3.1	6
131	Functional impairment, symptom severity, and overall quality of life in patients with advanced lung or colorectal cancer in six European countries: baseline findings from the ACTION study. Supportive Care in Cancer, 2021, 29, 5797-5810.	2.2	6
132	Burnout in physicians: a survey of the Danish society for palliative medicine. BMJ Supportive and Palliative Care, 2024, 14, 52-55.	1.6	6
133	Minimally important differences of EORTC QLQ-C30 scales in patients with lung cancer or malignant pleural mesothelioma – Interpretation guidance derived from two randomized EORTC trials. Lung Cancer, 2022, 167, 65-72.	2.0	6
134	Regional and age differences in specialised palliative care for patients with pancreatic cancer. BMC Palliative Care, 2021, 20, 192.	1.8	6
135	Clustering of EORTC QLQ-C30 health-related quality of life scales across several cancer types: Validation study. European Journal of Cancer, 2022, 170, 1-9.	2.8	6
136	Response to Commentary on Comparing translations of the EORTC QLQ-C30 using differential item functioning analyses. Quality of Life Research, 2006, 15, 1119-1120.	3.1	5
137	An evaluation of the response category translations of the EORTC QLQ-C30 questionnaire. Quality of Life Research, 2013, 22, 1483-1490.	3.1	5
138	Development of an item bank for computerized adaptive testing of self-reported cognitive difficulty in cancer patients. Neuro-Oncology Practice, 2017, 4, 189-196.	1.6	5
139	Predictors of the Extent of Agreements between Terminally ill Cancer Patients and their Family Caregivers in Taiwan. Quality of Life Research, 2006, 15, 405-406.	3.1	4
140	How does pain experience relate to the need for pain relief? A secondary exploratory analysis in a large sample of cancer patients. Supportive Care in Cancer, 2016, 24, 4187-4195.	2.2	4
141	Change in health-related quality of life over 1Âmonth in cancer patients with high initial levels of symptoms and problems. Quality of Life Research, 2016, 25, 2669-2674.	3.1	4
142	Improving information to caregivers of cancer patients: the Herlev Hospital Empowerment of Relatives through More and Earlier information Supply (HERMES) randomized controlled trial. Supportive Care in Cancer, 2020, 28, 939-950.	2.2	4
143	Development and content validation of a questionnaire measuring patient empowerment in cancer follow-up. Quality of Life Research, 2020, 29, 2253-2274.	3.1	4
144	Symptoms of advanced cancer in palliative medicine: a longitudinal study. BMJ Supportive and Palliative Care, 2023, 13, e415-e427.	1.6	4

#	Article	IF	CITATIONS
145	Process, content, and experiences of delivering the Carer Support Needs Assessment Tool Intervention (CSNAT-I) in the Danish specialised palliative care setting. Supportive Care in Cancer, 2022, 30, 377-387.	2.2	4
146	Assessing Emotional Functioning with Increased Power: Relative Validity of a Customized EORTC Short Form in the International ACTION Trial. Value in Health, 2019, 22, 92-98.	0.3	3
147	Content validation of the EORTC QLQ-C15-PAL with advanced cancer patients and health care professionals from palliative care services in Chile. BMC Palliative Care, 2020, 19, 81.	1.8	3
148	Investigating the response scale of the EORTC QLQ-C30 in German cancer patients and a population survey. Health and Quality of Life Outcomes, 2021, 19, 235.	2.4	2
149	Response to Letter to the Editor: Improving information to caregivers of cancer patients: the Herlev Hospital Empowerment of Relatives through More and Earlier information Supply (HERMES) randomized controlled trial. Supportive Care in Cancer, 2020, 28, 417-417.	2.2	1
150	Early specialised palliative care: interventions, symptoms, problems. BMJ Supportive and Palliative Care, 2021, 11, 444-453.	1.6	1
151	Development and Psychometric Evaluation of an Item Bank for Computerized Adaptive Testing of the EORTC Insomnia Dimension in Cancer Patients (EORTC CAT-SL). Applied Research in Quality of Life, 2021, 16, 827-844.	2.4	1
152	The quality of end of life care for Danish cancer patients who have received specialized palliative: a national survey using the Danish version of VOICES-SF. Supportive Care in Cancer, 2022, 30, 3593.	2.2	1
153	Advance care planningâ€"family carer psychological distress and involvement in decision making: the ACTION trial. BMJ Supportive and Palliative Care, 2023, 13, e807-e811.	1.6	1
154	Another look at floor and ceiling effects in the EORTC QLQ 30 Physical Functioning subscale and possible solutions. Cancer, 2022, 128, 2384-2385.	4.1	1
155	957 CHRONIC PAIN AND OTHER SEQUELAE IN BREAST CANCER SURVIVORS: A NATIONWIDE STUDY IN DENMARK. European Journal of Pain, 2006, 10, S248-S248.	2.8	O
156	OP55â€Evaluating costs of advance care planning; results from the international ACTION study. , 2019, , .		0
157	Similar levels of symptoms and problems were found among patients referred to specialized palliative care by general practitioners and hospital physicians: A nationwide register-based study of 31,139 cancer patients. Palliative Medicine, 2020, 34, 1118-1126.	3.1	O
158	Should analyses of large, national palliative care data sets with patient reported outcomes (PROs) be restricted to services with high patient participation? A register-based study. BMC Palliative Care, 2020, 19, 89.	1.8	0
159	Reply to: Olanzapine: Sancho Panza for clinicians who care for patients with advanced cancer. Supportive Care in Cancer, 2020, 28, 1539-1539.	2.2	O
160	Coverage of symptomatic toxicities from the common terminology criteria for adverse events (CTCAE) framework within the european organization for research and treatment of cancer (EORTC) item library Journal of Clinical Oncology, 2021, 39, e24117-e24117.	1.6	0
161	O-76â€First results from the action study. BMJ Supportive and Palliative Care, 2015, 5, A24.2-A24.	1.6	0
162	Title is missing!. , 2020, 17, e1003422.		0

#	Article	IF	Citations
163	Title is missing!. , 2020, 17, e1003422.		О
164	Title is missing!. , 2020, 17, e1003422.		0
165	Title is missing!. , 2020, 17, e1003422.		O
166	Title is missing!. , 2020, 17, e1003422.		0
167	Title is missing!. , 2020, 17, e1003422.		O
168	Title is missing!. , 2020, 17, e1003422.		0