

Baoyun Xia

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/39123/publications.pdf>

Version: 2024-02-01

15
papers

948
citations

933447

10
h-index

996975

15
g-index

15
all docs

15
docs citations

15
times ranked

1326
citing authors

#	ARTICLE	IF	CITATIONS
1	Tobacco-Specific Nitrosamines (NNAL, NNN, NAT, and NAB) Exposures in the US Population Assessment of Tobacco and Health (PATH) Study Wave 1 (2013–2014). <i>Nicotine and Tobacco Research</i> , 2021, 23, 573-583.	2.6	30
2	Concentrations of Cotinine and 4-(Methylnitrosamino)-1-(3-Pyridyl)-1-Butanol (NNAL) in U.S. Non-Daily Cigarette Smokers. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2021, 30, 1165-1174.	2.5	10
3	Sensitive Quantification of Nicotine in Bronchoalveolar Lavage Fluid by Acetone Precipitation Combined With Isotope-Dilution Liquid Chromatography-Tandem Mass Spectrometry. <i>ACS Omega</i> , 2021, 6, 13962-13969.	3.5	7
4	Associations between Biomarkers of Exposure and Lung Cancer Risk among Exclusive Cigarette Smokers in the Golestan Cohort Study. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 7349.	2.6	5
5	Vitamin E Acetate in Bronchoalveolar-Lavage Fluid Associated with EVALI. <i>New England Journal of Medicine</i> , 2020, 382, 697-705.	27.0	510
6	Biomarkers of Exposure among USA Adult Hookah Users: Results from Wave 1 of the Population Assessment of Tobacco and Health (PATH) Study (2013–2014). <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 6403.	2.6	7
7	Evaluation of Tobacco Smoke and Diet as Sources of Exposure to Two Heterocyclic Aromatic Amines for the U.S. Population: NHANES 2013–2014. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020, 29, 103-111.	2.5	5
8	Biomarkers of Exposure among Adult Smokeless Tobacco Users in the Population Assessment of Tobacco and Health Study (Wave 1, 2013–2014). <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020, 29, 659-667.	2.5	18
9	Opiate and Tobacco Use and Exposure to Carcinogens and Toxicants in the Golestan Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2020, 29, 650-658.	2.5	23
10	Urinary Biomarkers of Carcinogenic Exposure among Cigarette, Waterpipe, and Smokeless Tobacco Users and Never Users of Tobacco in the Golestan Cohort Study. <i>Cancer Epidemiology Biomarkers and Prevention</i> , 2019, 28, 337-347.	2.5	34
11	Evaluation of Bronchoalveolar Lavage Fluid from Patients in an Outbreak of E-cigarette, or Vaping, Product Use–Associated Lung Injury – 10 States, August–October 2019. <i>Morbidity and Mortality Weekly Report</i> , 2019, 68, 1040-1041.	15.1	159
12	High-throughput and sensitive analysis of urinary heterocyclic aromatic amines using isotope-dilution liquid chromatography–tandem mass spectrometry and robotic sample preparation system. <i>Analytical and Bioanalytical Chemistry</i> , 2016, 408, 8149-8161.	3.7	10
13	Assessing exposure to tobacco-specific carcinogen NNK using its urinary metabolite NNAL measured in US population: 2011–2012. <i>Journal of Exposure Science and Environmental Epidemiology</i> , 2016, 26, 249-256.	3.9	43
14	Quantitative analysis of five tobacco-specific <i>N</i> -nitrosamines in urine by liquid chromatography–atmospheric pressure ionization tandem mass spectrometry. <i>Biomedical Chromatography</i> , 2014, 28, 375-384.	1.7	35
15	Validation of a LC-MS/MS Method for Quantifying Urinary Nicotine, Six Nicotine Metabolites and the Minor Tobacco Alkaloids—Anatabine and Anabasine—in Smokers' Urine. <i>PLoS ONE</i> , 2014, 9, e101816.	2.5	52