

César González-Blanch

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3918994/publications.pdf>

Version: 2024-02-01

70
papers

3,181
citations

172457

29
h-index

161849

54
g-index

75
all docs

75
docs citations

75
times ranked

3962
citing authors

#	ARTICLE	IF	CITATIONS
1	Transdiagnostic group cognitive behavioural therapy for emotional disorders in primary care: the results of the PsicAP randomized controlled trial. <i>Psychological Medicine</i> , 2022, 52, 3336-3348.	4.5	20
2	Exploring the structure of the GAD-7 scale in primary care patients with emotional disorders: A network analysis approach. <i>Journal of Clinical Psychology</i> , 2022, 78, 283-297.	1.9	8
3	Psychometric properties of the Mindful Attention Awareness Scale (MAAS) in a first-episode psychosis sample.. <i>Psychological Assessment</i> , 2022, 34, 188-196.	1.5	6
4	The impact of persistent social anxiety on social functioning and health-related quality of life in young people with remitted first-episode psychosis. <i>Microbial Biotechnology</i> , 2022, 16, 868-874.	1.7	2
5	The mediating role of emotion regulation in transdiagnostic cognitive behavioural therapy for emotional disorders in primary care: Secondary analyses of the PsicAP randomized controlled trial. <i>Journal of Affective Disorders</i> , 2022, 303, 206-215.	4.1	10
6	Cost-effectiveness and cost-utility evaluation of individual vs. group transdiagnostic psychological treatment for emotional disorders in primary care (PsicAP-Costs): a multicentre randomized controlled trial protocol. <i>BMC Psychiatry</i> , 2022, 22, 99.	2.6	6
7	Emotion Regulation as a Moderator of Outcomes of Transdiagnostic Group Cognitive-Behavioral Therapy for Emotional Disorders. <i>Behavior Therapy</i> , 2022, 53, 628-641.	2.4	5
8	Cognitive variables associated with depressive and anxiety symptoms in patients with cancer: A five-year follow-up study. <i>Psycho-Oncology</i> , 2022, 31, 798-805.	2.3	3
9	Validity of Brief Scales for Assessing Transdiagnostic Cognitive Emotion Regulation Strategies in Primary Care Patients With Emotional Disorders. <i>European Journal of Psychological Assessment</i> , 2021, 37, 86-94.	3.0	12
10	Mental health interventions for healthcare workers during the first wave of COVID-19 pandemic in Spain. <i>Revista De PsiquiatrÃ­a Y Salud Mental</i> , 2021, 14, 83-89.	1.8	27
11	Factors differentiating panic disorder with and without comorbid symptoms of depression. <i>Psychology and Psychotherapy: Theory, Research and Practice</i> , 2021, 94, 523-540.	2.5	1
12	Moderators and predictors of treatment outcome in transdiagnostic group cognitive-behavioral therapy for primary care patients with emotional disorders. <i>Depression and Anxiety</i> , 2021, 38, 757-767.	4.1	5
13	The Horyzons project: a randomized controlled trial of a novel online social therapy to maintain treatment effects from specialist first-episode psychosis services. <i>World Psychiatry</i> , 2021, 20, 233-243.	10.4	62
14	Moderators of cognitive insight outcome in metacognitive training for first-episode psychosis. <i>Journal of Psychiatric Research</i> , 2021, 141, 104-110.	3.1	2
15	The role of cognitive factors in differentiating individuals with somatoform disorders with and without depression. <i>Journal of Psychosomatic Research</i> , 2021, 148, 110573.	2.6	1
16	Latent structure and factor invariance of somatic symptoms in the patient health questionnaire (PHQ-15). <i>Journal of Affective Disorders</i> , 2020, 261, 21-29.	4.1	25
17	Promoting Psychosocial Functioning and Recovery in Schizophrenia Spectrum and Other Psychotic Disorders. , 2020, , 419-445.		1
18	The role of social relatedness and self-beliefs in social functioning in first-episode psychosis: Are we overestimating the contribution of illness-related factors?. <i>European Psychiatry</i> , 2020, 63, e92.	0.2	5

#	ARTICLE	IF	CITATIONS
19	Disability and perceived stress in primary care patients with major depression. <i>Psicothema</i> , 2020, 32, 167-175.	0.9	3
20	Cognitive factors related to distress in patients recently diagnosed with cancer. <i>Psycho-Oncology</i> , 2019, 28, 1987-1994.	2.3	5
21	HORYZONS trial: protocol for a randomised controlled trial of a moderated online social therapy to maintain treatment effects from first-episode psychosis services. <i>BMJ Open</i> , 2019, 9, e024104.	1.9	65
22	Factorial invariance of a computerized version of the GAD-7 across various demographic groups and over time in primary care patients. <i>Journal of Affective Disorders</i> , 2019, 252, 114-121.	4.1	53
23	A computerized version of the Patient Health Questionnaire-4 as an ultra-brief screening tool to detect emotional disorders in primary care. <i>Journal of Affective Disorders</i> , 2018, 234, 247-255.	4.1	45
24	The factor structure of the Medical Outcomes Studyâ€“Social Support Survey: A comparison of different models in a sample of recently diagnosed cancer patients. <i>Journal of Psychosomatic Research</i> , 2018, 108, 32-38.	2.6	22
25	Domain-specific associations between disability and depression, anxiety, and somatization in primary care patients. <i>Psychiatry Research</i> , 2018, 269, 596-601.	3.3	9
26	The association between different domains of quality of life and symptoms in primary care patients with emotional disorders. <i>Scientific Reports</i> , 2018, 8, 11180.	3.3	44
27	Enhancing social functioning in young people at Ultra High Risk (UHR) for psychosis: A pilot study of a novel strengths and mindfulness-based online social therapy. <i>Schizophrenia Research</i> , 2018, 202, 369-377.	2.0	99
28	Factor structure and measurement invariance across various demographic groups and over time for the PHQ-9 in primary care patients in Spain. <i>PLoS ONE</i> , 2018, 13, e0193356.	2.5	67
29	Terapia cognitivo-conductual transdiagñ³stica en Atenci³n Primaria: un contexto ideal. <i>Revista De Psicoterapia</i> , 2018, 29, 37-52.	0.0	3
30	Predictors of functional recovery in first-episode psychosis: A systematic review and meta-analysis of longitudinal studies. <i>Clinical Psychology Review</i> , 2017, 58, 59-75.	11.4	192
31	Modelo jerÁrquico de diagn³stico y derivaci³n de los trastornos mentales comunes en centros de atenci³n primaria. Una propuesta a partir del ensayo clÁnico PsicAP. <i>Ansiedad Y Estrés</i> , 2017, 23, 124-129.	0.2	21
32	Impacto econ³mico y carga de los trastornos mentales comunes en Espa±a: una revisi³n sistemÁtica y crÁtica. <i>Ansiedad Y Estrés</i> , 2017, 23, 118-123.	0.2	27
33	Tratamiento psicol³gico de los trastornos emocionales en atenci³n primaria: fundamentos te³ricos y empÁricos del estudio PsicAP. <i>Ansiedad Y Estrés</i> , 2017, 23, 91-98.	0.2	4
34	Online peer-to-peer support in youth mental health: seizing the opportunity. <i>Epidemiology and Psychiatric Sciences</i> , 2016, 25, 123-126.	3.9	37
35	Beyond Clinical Remission in First Episode Psychosis: Thoughts on Antipsychotic Maintenance vs. Guided Discontinuation in the Functional Recovery Era. <i>CNS Drugs</i> , 2016, 30, 357-368.	5.9	67
36	Planning Abilities in Patients with Anorexia Nervosa Compared with Healthy Controls. <i>Clinical Neuropsychologist</i> , 2016, 30, 228-242.	2.3	7

#	ARTICLE	IF	CITATIONS
37	Longitudinal relationship between expressed emotion and cannabis misuse in young people with first-episode psychosis. <i>European Psychiatry</i> , 2015, 30, 20-25.	0.2	7
38	Online social media: New data, new horizons in psychosis treatment. <i>Schizophrenia Research</i> , 2015, 166, 345-346.	2.0	10
39	Social Functioning Trajectories of Young First-Episode Psychosis Patients with and without Cannabis Misuse: A 30-Month Follow-Up Study. <i>PLoS ONE</i> , 2015, 10, e0122404.	2.5	20
40	Online, social media and mobile technologies for psychosis treatment: A systematic review on novel user-led interventions. <i>Schizophrenia Research</i> , 2014, 156, 96-106.	2.0	311
41	Differential associations of cognitive insight components with pretreatment characteristics in first-episode psychosis. <i>Psychiatry Research</i> , 2014, 215, 308-313.	3.3	13
42	The relevance of cognitive, clinical and premorbid variables in predicting functional outcome for individuals with first-episode psychosis: A 3 year longitudinal study. <i>Psychiatry Research</i> , 2013, 209, 302-308.	3.3	54
43	Asociaci3n entre rumiaciones y malestar emocional en pacientes reci3n diagnosticados de c4ncer. <i>Psicooncologia</i> , 2013, 10, .	0.3	2
44	Identifying attentional deficits in people with first-episode psychosis with the Scale for the Assessment of Negative Symptoms attention subscale: is it possible?. <i>Comprehensive Psychiatry</i> , 2012, 53, 701-705.	3.1	5
45	A Digit Symbol Coding Task as a Screening Instrument for Cognitive Impairment in First-Episode Psychosis. <i>Archives of Clinical Neuropsychology</i> , 2011, 26, 48-58.	0.5	48
46	Review: family interventions reduce relapse or hospitalisation in people with schizophrenia. <i>Evidence-Based Mental Health</i> , 2011, 14, 115-115.	4.5	1
47	Effects of Family Psychoeducation on Expressed Emotion and Burden of Care in First-Episode Psychosis: A Prospective Observational Study. <i>Spanish Journal of Psychology</i> , 2010, 13, 389-395.	2.1	32
48	Prognostic value of cognitive functioning for global functional recovery in first-episode schizophrenia. <i>Psychological Medicine</i> , 2010, 40, 935-944.	4.5	60
49	First-episode schizophrenia patients neuropsychologically within the normal limits: Evidence of deterioration in speed of processing. <i>Schizophrenia Research</i> , 2010, 119, 18-26.	2.0	34
50	Cannabis use and cognitive functioning in first-episode schizophrenia patients. <i>Schizophrenia Research</i> , 2010, 124, 142-151.	2.0	52
51	Brief cognitive-behavioral therapy with fibromyalgia patients in routine care. <i>Comprehensive Psychiatry</i> , 2009, 50, 517-525.	3.1	27
52	Cognitive functioning and negative symptoms in first episode schizophrenia: Different patterns of correlates. <i>Neurotoxicity Research</i> , 2008, 14, 227-235.	2.7	13
53	Treatment of the first episode of schizophrenia: An update on pharmacologic and psychological interventions. <i>Current Psychiatry Reports</i> , 2008, 10, 202-209.	4.5	15
54	Epidemiological factors associated with treated incidence of first4episode non4effective psychosis in Cantabria: insights from the Clinical Programme on Early Phases of Psychosis. <i>Microbial Biotechnology</i> , 2008, 2, 178-187.	1.7	131

#	ARTICLE	IF	CITATIONS
55	1-year follow-up study of cognitive function in first-episode non-affective psychosis. <i>Schizophrenia Research</i> , 2008, 104, 165-174.	2.0	42
56	Lack of influence of COMT Val158Met genotype on cognition in first-episode non-affective psychosis. <i>Schizophrenia Research</i> , 2008, 102, 206-209.	2.0	15
57	Antipsychotic-Induced Weight Gain in Chronic and First-Episode Psychotic Disorders. <i>CNS Drugs</i> , 2008, 22, 547-562.	5.9	229
58	Non-pharmacological management of antipsychotic-induced weight gain: Systematic review and meta-analysis of randomised controlled trials. <i>British Journal of Psychiatry</i> , 2008, 193, 101-107.	2.8	272
59	Pretreatment predictors of cognitive deficits in early psychosis. <i>Psychological Medicine</i> , 2008, 38, 737-746.	4.5	24
60	Cannabis abuse is associated with decision-making impairment among first-episode patients with schizophrenia-spectrum psychosis. <i>Psychological Medicine</i> , 2008, 38, 1257-1266.	4.5	49
61	Preserved Orbitofrontal Function in First-Episode Schizophrenia. <i>Journal of Nervous and Mental Disease</i> , 2008, 196, 67-70.	1.0	4
62	Lack of association between clinical and cognitive change in first-episode psychosis: the first 6 weeks of treatment. <i>Canadian Journal of Psychiatry</i> , 2008, 53, 839-47.	1.9	11
63	Cognitive dysfunction in first-episode psychosis: the processing speed hypothesis. <i>British Journal of Psychiatry</i> , 2007, 191, s107-s110.	2.8	108
64	Reduced thalamic volume in first-episode non-affective psychosis: Correlations with clinical variables, symptomatology and cognitive functioning. <i>NeuroImage</i> , 2007, 35, 1613-1623.	4.2	66
65	Caudate nucleus volume and its clinical and cognitive correlations in first episode schizophrenia. <i>Schizophrenia Research</i> , 2007, 91, 87-96.	2.0	76
66	Cognitive dimensions in first-episode schizophrenia spectrum disorders. <i>Journal of Psychiatric Research</i> , 2007, 41, 968-977.	3.1	77
67	A 12-Week Randomized Clinical Trial to Evaluate Metabolic Changes in Drug-Naive, First-Episode Psychosis Patients Treated With Haloperidol, Olanzapine, or Risperidone. <i>Journal of Clinical Psychiatry</i> , 2007, 68, 1733-1740.	2.2	87
68	Cognitive functioning in the early course of first-episode schizophrenia spectrum disorders. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2006, 256, 364-371.	3.2	38
69	Attenuation of Antipsychotic-Induced Weight Gain With Early Behavioral Intervention in Drug-Naive First-Episode Psychosis Patients. <i>Journal of Clinical Psychiatry</i> , 2006, 67, 1253-1260.	2.2	152
70	Prefrontal cognitive functions in stabilized first-episode patients with schizophrenia spectrum disorders: A dissociation between dorsolateral and orbitofrontal functioning. <i>Schizophrenia Research</i> , 2005, 77, 279-288.	2.0	79